

## **PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Aqupharm 9 Ringer's Solution for Infusion

### **1. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Sodium Chloride 0.86% w/v  
Potassium Chloride 0.03% w/v  
Calcium Chloride Dihydrate 0.03% w/v

Approx. ion concentration in millimoles per litre:  
Sodium 147, Potassium 4, Calcium 2.2, Chloride 156.

### **3. PHARMACEUTICAL FORM**

Infusion solution

### **4. PACKAGE SIZE**

500ml  
1000ml

### **5. TARGET SPECIES**

Dogs and Cats

### **6. INDICATION(S)**

Sterile non-pyrogenic infusion solution for use in dogs and cats under veterinary supervision.

### **7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Administration: Read package leaflet before use. Do not use unless the solution is clear and free from particles. This product does not contain an antimicrobial preservative. For single use only. Wash hands after use.

### **8. WITHDRAWAL PERIOD**

### **9. SPECIAL WARNING(S), IF NECESSARY**

<User Warnings>

### **10. EXPIRY DATE**

## **11. SPECIAL STORAGE CONDITIONS**

Storage: Do not store above 25°C. Do not freeze.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority. Discard any remaining solution.

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]**

For animal treatment only. To be supplied only on veterinary prescription.

POM-V

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of sight and reach of children.

## **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Ecuphar NV, Legeweg 157-i, 8020 Oostkamp, Belgium

## **16. MARKETING AUTHORISATION NUMBER(S)**

Vm 32742/4023

## **17. MANUFACTURER’S BATCH NUMBER**

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Aqupharm 9 Ringer's Solution for Infusion

### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Sodium Chloride 0.86% w/v  
Potassium Chloride 0.03% w/v  
Calcium Chloride Dihydrate 0.03% w/v

Approx. ion concentration in millimoles per litre:  
Sodium 147, Potassium 4, Calcium 2.2, Chloride 156

### **3. PHARMACEUTICAL FORM**

Infusion solution

### **4. PACKAGE SIZE**

20 x 500ml  
10 x 1000ml

### **5. TARGET SPECIES**

Dogs and Cats

### **6. INDICATION(S)**

Sterile non-pyrogenic infusion solution for use in dogs and cats under veterinary supervision.

### **7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Administration: Read package leaflet before use. Do not use unless the solution is clear and free from particles. This product does not contain an antimicrobial preservative. For single use only. Wash hands after use.

### **8. WITHDRAWAL PERIOD**

### **9. SPECIAL WARNING(S), IF NECESSARY**

<User Warnings>

### **10. EXPIRY DATE**

## **11. SPECIAL STORAGE CONDITIONS**

Storage: Do not store above 25°C. Do not freeze.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority. Discard any remaining solution.

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]**

*For animal treatment only. To be supplied only on veterinary prescription.*

POM-V

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of sight and reach of children.

## **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Ecuphar NV, Legeweg 157-i, 8020 Oostkamp, Belgium

## **16. MARKETING AUTHORISATION NUMBER(S)**

Vm 32742/4023

## **17. MANUFACTURER’S BATCH NUMBER**

Lot:

**PACKAGE LEAFLET FOR:****1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing Authorisation Holder:

Ecuphar NV, Legeweg 157-i, 8020 Oostkamp, Belgium

Manufacturer responsible for batch release:

Infomed Fluids SRL, 50 Theodor Pallady blvd., District 3, 032266 Bucharest, Romania

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Aqupharm 9 Ringer's Solution for Infusion.

**3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS**

Presentation

Aqupharm No. 9 is a sterile, preservative free, solution for infusion, presented in a flexible pouch containing:

Sodium Chloride 0.86% w/v

Potassium Chloride 0.03% w/v

Calcium Chloride, dihydrate 0.03% w/v

Ions

Sodium 147.0mmol/l

Potassium 4.0mmol/l

Calcium 2.2mmol/l

Chloride 156.0mmol/l

The approximate ion concentration in millimoles per litre is sodium 147, potassium 4, calcium 2.2, chloride 156.

**4. INDICATION(S)**

Aqupharm No.9 is an isotonic solution used in dogs and cats for the treatment of dehydration with salt depletion and where there has been some intracellular potassium loss. In cases of persistent vomiting there are substantial losses of both hydrogen and chloride ions, resulting in an excess of intracellular sodium. Renal compensation leads to an increased potassium loss and consequent hypokalaemia. It is indicated for pyometra when associated with severe vomiting.

**5. CONTRAINDICATIONS**

Contra-indications:

Sodium overload may occur in cases with myocardial and renal damage. It should also be appreciated that in the period following surgical interference or severe trauma there may be an inability to excrete excessive sodium.

Undesirable effects:

Hypernatraemia (sodium overload) or an inability to excrete excessive sodium. Thrombosis of a chosen vein is always a possibility with intravenous infusion. If infusions is protracted then another vein should be selected after 12-24 hours.

## **6. ADVERSE REACTIONS**

## **7. TARGET SPECIES**

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Aquopharm No.9 should be prewarmed to 37°C to prevent hypothermia.

Remove outer bag and protective giving set inlet tab. Push cannula fully into giving set. Prime giving set. Perform venepuncture and immediately attach giving set. Adjust infusion rate as required. Delivery is from a closed circuit, it does not need an air inlet.

Giving sets should be changed every 24 hours.

The quantity of fluid and electrolyte for administration will consider existing deficits, maintenance needs and continuing losses.

The existing deficit is that which has been lost prior to examination. This must be estimated by evaluating the patient's history, making a physical examination and using laboratory aids. Maintenance therapy is to replace normal losses occurring via urine, faeces, respiratory tract and skin. As a general rule, maintenance therapy requires 50 ml/kg bodyweight/day. Continuing losses during a disease period should be estimated whenever possible, i.e., quantity of vomit, diarrhoea or blood loss.

The clinical response of the animal rather than formulae or equations should be used to guide fluid therapy. The intravenous route of administration is preferred. Indwelling intravenous catheters offer significant advantage in intravenous fluid therapy. Subcutaneous administration may be used for isotonic and non-irritating solutions.

The rate of administration should be considered with each individual patient. The aim should be to correct about half of the calculated deficit in the first 1-2 hours. As a general rule the following formula is the maximum satisfactory rate (less where cardiovascular or pulmonary disease exists).

Maximum rate = Body wt (kg) × 90 = ml fluid per hour

This rate should be slowed after the first hour and considerably slowed if no urine flow is established. Signs of over rapid administration include restlessness, moist lung sounds, tachycardia, tachypnoea, nasal discharge, coughing, vomiting and diarrhoea.

## **9. ADVICE ON CORRECT ADMINISTRATION**

## **10. WITHDRAWAL PERIOD(S)**

## **11. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C. Do not freeze. Keep out of sight and reach of children.

## **12. SPECIAL WARNING(S)**

### **Precautions and warnings:**

For animal treatment only. In evaluating an animal for possible fluid therapy the state of hydration, electrolyte balance, acid-base balance, renal function and caloric balance should be considered. Evaluation will be based on history, physical examination and laboratory testing.

Although The Solution contains potassium the quantity may be inadequate in the presence of intracellular potassium loss; where such deficiency is known to occur it may be necessary to give oral potassium supplements.

### Treatment of overdose:

#### Symptoms:

Associated signs of hypernatraemia include pronounced thirst, dry mucous membranes, constipation, hyperpyrexia, CNS disturbances, and ultimately convulsions. A plasma Na<sup>+</sup> concentration of >150mEq/l and a urine specific gravity of >1.030 indicate a hypernatraemic state.

### **Treatment of overdose:**

Injection of a diuretic.

### **User warning:**

Wash hands after Use.

Pharmaceutical precautions

This product does not contain an antimicrobial preservative. Single use only, any remaining solution should be discarded.

Interaction with other medicines:

Drugs should not be mixed in infusion containers or through the giving sets unless the components are of known compatibility. The user should refer to the manufacturer's literature for any drug substance which he or she proposes to co-administer, and also to the Appendix of Drug Incompatibilities in the current edition of The Veterinary Formulary. Aqupharm No. 9 is incompatible with Sodium bicarbonate intravenous solution and Noradrenaline acid tartrate.

Do not use this veterinary medical product after the expiry date stated on the label.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Dispose of any unused product and empty containers in accordance with guidance from your local waste authority.

### **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

August 2022

### **15. OTHER INFORMATION**

[POM-V]


UK authorised veterinary medicinal product.

To be supplied only on veterinary prescription.

Package quantities 500ml and 1000ml flexible pouches. Not all pack sizes may be marketed.

Pack sizes: Cardboard box containing 20 bags of 500ml solution for infusion, 10 bags of 1000ml solution for infusion.

*Marketing Authorisation Number: Vm 32742/4023*



Approved: 11 August 2022