

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING

UNITS: PIPETTE LABEL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Clear Wormer 20 mg Spot-on Solution for Cats & Kittens

Praziquantel

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Praziquantel 20 mg

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

20 mg

4. ROUTE(S) OF ADMINISTRATION

Spot-on

5. WITHDRAWAL PERIOD

6. BATCH NUMBER

Batch No:

7. EXPIRY DATE

EXP:

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS BLISTER

LABEL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Clear Wormer 20 mg Spot-on Solution for Cats & Kittens

Praziquantel

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Pets Choice Limited

3. EXPIRY DATE

EXP:

4. BATCH NUMBER

Batch No:

5. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only

MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO

PACKAGE LEAFLET, Carton Box/Blister card

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Pets Choice Ltd
Brentwood House
Lower Philips Road
Whitebirk Industrial Estate
Blackburn
Lancashire
BB1 5UD
UK

Manufacturer responsible for batch release:

Pets Choice Ltd
8 Wemberham Lane
Yatton
North Somerset
BS49 4BS
United Kingdom

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Clear Wormer 20 mg Spot-on Solution for Cats and Kittens

Praziquantel

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

Contains 20 mg praziquantel in 0.5 ml Spot-on solution.
Clear, colourless to pale amber solution

4. PHARMACEUTICAL FORM

Spot-on solution.

5. PACKAGE SIZE

2 x 0.5 ml 2 pipettes
4 x 0.5 ml 4 pipettes Multi Cats Pack

6. INDICATION(S)

Wormer treatment

Bob Martin Clear Wormer 20 mg Spot-on Solution for Cats & Kitten is an effective treatment against tapeworms affecting cats, including *Dipylidium caninum*, *Taenia taeniaformis*, *Echinococcus multilocularis*.

7. CONTRAINDICATIONS

Do not use on cats weighing less than 1 kg bodyweight.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

8. ADVERSE REACTIONS

On very rare occasions a transient local reaction such as scurf (flaking skin) or mild exudation (weeping) may be observed at the application site following treatment.

On very rare occasions salivation may occur if the cat licks the site of application immediately after treatment, as the product is bitter tasting. This is not a sign of intoxication and will disappear within a few minutes of treatment.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

9. TARGET SPECIES

Cats

10. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Suitable for cats and kittens over 1kg. For cats over 5kg additional pipettes will be required.

The minimum dose rate is 8 mg/kg bodyweight, which equates to 1 pipette per 2.5 kg bodyweight.

Bodyweight	Number of Tubes	Quantity of Praziquantel	mg/kg bw
1 - 2.5 kg	1	20 mg	8 - 20
>2.5 - 5 kg	2	40 mg	8 - 16
>5 kg	3	60 mg	maximum 12

Note: If your cat weighs more than 5kg apply an extra pipette per 2.5kg bodyweight.

11. ADVICE ON CORRECT ADMINISTRATION

Topical / spot-on

DO NOT administer Bob Martin Clear Wormer 20mg Spot-on Solution by mouth. Weigh your cat carefully before calculating the dose.



1. Remove a single pipette from the pack. Hold the pipette upright and allow any liquid in the neck of the pipette to drain into the main body. Continue holding the pipette upright and snap off the tip.



2. Part the coat on the back of the neck immediately behind the head until the skin is visible. If the cat is wearing a collar remove it first so that it does not restrict application. Correct application will minimise the chance for your cat to lick the product.



3. Tap the narrow part of the pipette to ensure the contents are within the application nozzle. Place the tip of the pipette on the skin and slowly squeeze the entire contents onto the skin. Each application will be absorbed into the skin. Allow the contents of each pipette to be absorbed before applying any additional pipettes.

To minimise the possibility of run-off after application of more than one pipette, it is advised that the applications should be performed slowly to allow absorption and that it may be advisable to allow the contents of the previous pipette to be absorbed before applying another.

Tapeworm: infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Flea control: flea infestations can be controlled by the regular use of effective flea control remedies.

Mice control: if cats roam and hunt, contact with, and consumption of, mice and subsequent re-infestation with *Taenia taeniaeformis* is impossible to prevent.

It is recommended to re-apply the product when signs of tapeworm infestation reappear or at monthly intervals.

12. WITHDRAWAL PERIOD

13. SPECIAL STORAGE PRECAUTIONS

Store away from food, drink and animal feeding stuffs.

Store below 25°C.

Store in a dry place in the original package.

Keep product in the outer carton.

Do not use this product after the expiry date which is stated on the label after 'EXP'.

The expiry date refers to the last day of that month.

14. SPECIAL WARNING(S)<User Warnings>

i) Special precautions for use in animals

For external use only

Do not use in cats weighing less than 1 kg bodyweight.

Care should be taken to avoid the contents of the pipette coming into contact with the eyes or mouth of the recipient animal.

Do not allow recently treated animals to groom each other.

ii) Special precautions to be taken by the person administering the medicinal product to animals

This product can be an irritant to the skin and eyes.

Care should be taken to prevent contact of the solution with the skin, eyes and mouth, including hand-to eye and hand-to mouth contact.

If accidental contact with the skin or eyes occurs, wash off any skin contamination with soap and water immediately. Rinse the affected eyes thoroughly with clean, fresh water.

In the event of skin or eye contact, seek medical advice if irritation persists and show the Doctor this package.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or label to the physician.

Laboratory studies with the excipient N-methyl-2-pyrrolidone in rabbits and rats have shown evidence of teratogenic, foetotoxic, maternotoxic and reprotoxic effects. Avoid direct contact with the product and application site.

Do not stroke or groom animals until the area of application is dry (typically around 1 hour after application).

Wash hands thoroughly after use.
Do not eat, drink or smoke during application.

Store away from food, drink or animal feedingstuffs.

iii) Other precautions

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority (e. g. experts or institutes of parasitology). If the cat has visited areas where Echinococcus spp. are prevalent, a veterinarian should be consulted.

Keep recently treated pets away from varnished, polished, plastic or leather surfaces.

There are no contra-indications against use during pregnancy and lactation. Overdosing can lead to slight skin reactions which disappear without treatment within a few days.

15. EXPIRY DATE

EXP End

16. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

17. DATE ON WHICH THE LABEL WAS LAST APPROVED

August 2023

18. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.
[Distribution category]

AVM-GSL

19. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

20. MARKETING AUTHORISATION NUMBER(S)

Vm 52797/4019

21. MANUFACTURER’S BATCH NUMBER

22. OTHER INFORMATION

Please note that this product is only effective against tapeworms and does not treat other parasitic worms, including roundworms.

EFFECTIVE WORM CONTROL

There is no permanent treatment for worms. It is recommended to re-apply the product when signs of tapeworm infestation re-appear or at monthly intervals. It is also important to remove sources of possible re-infection such as fleas and mice.

To safeguard your health and that of your pet please read this carton carefully before you apply the product to your pet. The directions for use can be found under this flap on the back of the carton, and should be read before each use.

If signs of disease persist or appear consult a veterinary surgeon.

HOW OFTEN TO USE:


Use the treatment record below to record when you last treated your cat and to indicate when the next treatment is due.

Treatment Calendar	Cat's name	
	First treatment	Second treatment
Date	Day 1	Due + 1 month

Made by Pet's Choice Limited in the UK under license from Martin & Martin Holdings

Bob Martin is a registered trademark of Martin & Martin Holdings.

Since 1892 Bob Martin has provided vet-quality healthcare solutions for happy, healthypets.



Approved: 09 August 2023