

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO PACKAGE LEAFLET

Blister card with box containing 1, 2 or 4 pipettes

4Fleas

Protector 60mg

Spot-on solution

For Cats & Kittens 1kg – 6kg

Kills flea eggs and larvae

Protects for 3 months

Prevents flea multiplication

Johnson's 4fleas protector 60mg spot-on solution for Cats & Kittens weighing between 1kg and 6kg. Easy spot-on administration. One vial prevents flea multiplication (*Ctenocephalides felis*) in dogs by inhibiting egg and larvae development for 3 months.



DIRECTIONS FOR USE: Do not remove vial from pack until ready for use. For external use only. Use one vial for each animal requiring treatment.

Apply Johnson's 4fleas protector 60mg spot-on solution for cats & kittens directly on to cat's skin at base of neck, expelling the contents by carefully squeezing bottom of vial 4 times. Care should be taken to maintain contact between vial and skin during application. A second application can be made once, 3 months after the first one, to give a further 3 months' protection.

If you consider that your dog needs repeat treatment after this period, consult your veterinary surgeon.

CONTRA-INDICATIONS AND SPECIAL WARNINGS: Do not use on pregnant or nursing females.

Do not use on kittens weighing less than 1kg, If your cat weighs more than 6kg use an alternative product or seek veterinary advice. Do not use if your cat is unwell or recovering from an illness. In rare cases, hair loss, skin flaking, itching, skin redness and irritation may occur at the application site. Generalised itching and hair loss has been reported. In very rare cases, increased salivation or vomiting has been observed after treatment.

FURTHER INFORMATION: If there are several animals in the same household, appropriate treatment should be carried out for each dog or cat, preferably at the same time. A flea elimination strategy should be aimed at all stages of the flea life-cycle. Cats treated with the Johnson's 4Fleas protector (10%) may be at risk of infestation by a new population of adult fleas from outside the household. Infestations occurring during the initial administration or appearing during the treatment period

may be eliminated with an additional appropriate product authorised for use on cats in order to kill adult fleas. Fleas from pets often infest pet's basket, bedding and regular resting areas, such as carpets and soft furnishings. Wash or change the dog's bedding and vacuum carpets and furnishings regularly, then use a suitable treatment for the household that kills fleas.

The effect of bathing or wetting immediately before or after treatment has not been examined. It is likely that efficacy will be reduced if the cat is wetted or shampooed immediately before or after treatment.

OPERATOR WARNINGS: Avoid contact with skin and eyes. In case of accidental contact, rinse thoroughly with water. In case of eye irritation, seek medical attention. Wash hands after use.

DISPOSAL ADVICE: Dispose of empty packaging and any remaining product in the household refuse.

The product should not be allowed to enter watercourses as this may be dangerous for fish and other aquatic organisms.

**KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.
FOR EXTERNAL USE ONLY.**

Contents: 1 vial containing 60mg of Pyriproxyfen in 0.6ml of solution.

AVM-GSL

Vm. 17902/4012

UK authorised veterinary medicinal product

Marketing authorisation holder: Alfamed S.A.S. 06517 Carros Cedex-France

Manufactured by Virbac S.A. 06516 Carros Cedex-France

Johnson's Veterinary Products Ltd,

Sutton Coldfield, West Midlands UK B75 7DF.

Tel: 0121 378 1684 e-mail: info@johnsons-vet.com

Approved: 10/11/2017



