

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Carton Box**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Doxytab vet. Flav. 15 mg Tablets for dogs and cats  
doxycycline

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains  
Active substance:  
Doxycycline 15 mg  
(as doxycycline hyclate 17.3 mg)

**3. PHARMACEUTICAL FORM**

Tablet.

**4. PACKAGE SIZE**

3 x 10 tablets  
5 x 10 tablets  
10 x 10 tablets  
1 x 30 tablets  
5 x 30 tablets  
10 x 30 tablets

**5. TARGET SPECIES**

Dogs and cats.

**6. INDICATIONS**

**7. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**8. WITHDRAWAL PERIOD**

**9. SPECIAL WARNINGS, IF NECESSARY**

Read the package leaflet before use.

People with known hypersensitivity to tetracyclines should avoid contact with this product.

Children are particularly at risk of intoxication. Take care that children do not come into contact with the product.

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Keep the blister in the outer carton. Any remaining portions of divided tablets should be returned in the opened blister and given at the next administration.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

CP Pharma Handelsgesellschaft mbH  
Ostlandring 13  
31303 Burgdorf  
Germany

**16. MARKETING AUTHORISATION NUMBER**

Vm 20916/4035

**17. MANUFACTURER'S BATCH NUMBER**

Batch number:

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

Aluminium/ PVC/ PE/PVDC blister pack

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Doxytab vet. Flav. 15 mg Tablets for dogs and cats  
doxycycline

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

CP-Pharma Handelsgesellschaft mbH

**3. EXPIRY DATE**

EXP {month/year}

**4. BATCH NUMBER**

Batch number:

**5. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

**B. PACKAGE LEAFLET**

## PACKAGE LEAFLET:

**Doxytab vet. Flav. 15 mg Tablets for dogs and cats [RMS and other CMS]  
Doxytab vet. 15 mg Tablets for dogs and cats [Denmark, Sweden, Finland,  
France, Ireland and UK]  
Cepedox vet. 15 mg Tablets for dogs and cats [Spain]**

### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder and manufacturer responsible for batch release:

CP Pharma Handelsgesellschaft mbH  
Ostlandring 13  
31303 Burgdorf  
Germany

### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Doxytab vet. Flav. 15 mg Tablets for dogs and cats  
doxycycline

### 3. STATEMENT OF THE ACTIVE SUBSTANCE

Each tablet contains:

#### Active substance:

Doxycycline	15 mg
(as doxycycline hyclate	17.3 mg)

Yellow with brown spots, round and convex tablet with a cross-shaped break line on one side. The tablets can be divided into 2 or 4 equal parts.

### 4. INDICATIONS

Treatment of the following conditions caused by bacteria sensitive to doxycycline:

#### Dogs:

Rhinitis caused by *Bordetella bronchiseptica* and *Pasteurella* spp.;  
Bronchopneumonia caused by *Bordetella* spp. and *Pasteurella* spp.;  
Interstitial nephritis caused by *Leptospira* spp.;

#### Cats:

Respiratory infections caused by *Bordetella bronchiseptica*, *Chlamydophila felis*,  
*Pasteurella* spp.

## 5. CONTRAINDICATIONS

Do not use in known cases of hypersensitivity to tetracyclines or to any of the excipients.

## 6. ADVERSE REACTIONS

Gastrointestinal disorders such as vomiting, diarrhoea and oesophagitis have been reported as side effects following long term doxycycline therapy.

In very young animals discoloration of the teeth may occur by the formation of a tetracycline-calcium phosphate complex.

Hypersensitivity reactions, photosensitivity and in exceptional cases photodermatitis may occur after exposure to intense daylight.

Retardation of skeletal growth of young animals (reversible upon discontinuation of therapy).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Dogs and cats.

## 8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

### *Method of administration*

Oral use.

### *Dosage*

The general recommended dose is 10 mg doxycycline per kg per day. The daily dose may be split into two administrations per day (i.e. 5 mg/kg bodyweight twice daily).

The majority of routine cases are expected to respond after between 5 and 7 days of therapy. Therapy should continue for 2 to 3 days beyond the clinical cure for acute infections. In chronic or refractory cases, a longer course of therapy, up to 14 days, may be required.


In dogs with interstitial nephritis due to leptospirosis, treatment for 14 days is recommended.

In cats with *C. felis* infections, it is recommended to administer treatment for a period of 28 days in order to ensure elimination of the organism.

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing.

The most appropriate tablet strength should be used in order to minimize divided tablets to be kept until the next dosing. Tablets can be divided into 2 or 4 equal parts to ensure accurate dosing. Place the tablet on a flat surface, with its scored side facing up and the convex (rounded) side facing the surface.





2 equal parts: press down with your thumbs on both sides of the tablet.

4 equal parts: press down with your thumb in the middle of the tablet.

## 9. ADVICE ON CORRECT ADMINISTRATION

Tablets should be administered with food.

Return any divided tablets to the blister pack. Divided tablets should be used at the next administration. Any divided tablets remaining after the last administration of the product should be discarded.

## 10. WITHDRAWAL PERIOD

Not applicable.

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Keep the blister in the outer carton. Any remaining portions of divided tablets should be returned in the opened blister and given at the next administration.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and blister after EXP.

## 12. SPECIAL WARNINGS

Special warnings for each target species:

None.

Special precautions for use in animals

The product should be administered with caution to animals with dysphagia or diseases accompanied with vomiting, since administration of doxycycline hyclate tablets has been associated with oesophageal erosion.

In order to reduce the likelihood of oesophageal irritation as well as other gastrointestinal side effects, the product should be administered together with food. Special care should be taken when administering the product to animals with liver disease, since increases in hepatic enzymes have been documented in some animals after doxycycline treatment.

The product should be administered with caution to young animals, since tetracyclines as a class may cause permanent discolouration of the teeth, when administered during tooth development. However, human literature indicates that

doxycycline is less likely than other tetracyclines to cause these abnormalities, due to its reduced ability to chelate calcium.

As tablets are flavoured store tablets out of reach of the animals in order to avoid accidental ingestion.

Due to the likely variability (time, geographical) in the occurrence of resistance of bacteria for doxycycline, bacteriological sampling and susceptibility testing are recommended. Official, national and regional antimicrobial policies should be taken into account when the product is used.

Use of the product deviating from the instructions given in this package leaflet may increase the prevalence of bacteria resistant to doxycycline and may decrease the effectiveness of treatment with other tetracyclines, due to the potential for cross-resistance.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product may cause hypersensitivity reactions. People with known hypersensitivity to tetracyclines should avoid contact with the veterinary medicinal product.

If you develop symptoms following exposure such as skin rash, seek medical advice immediately and show the package leaflet to the physician. This product may cause serious gastrointestinal effects if ingested, especially by children. To avoid accidental ingestion, unused tablet parts should be returned to the open blister space and inserted back into the carton that should be stored in a safe place out of sight and reach of children. In case of accidental ingestion, seek medical advice and show the package leaflet or the label to the physician.

Wash hands after use.

#### Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Tetracyclines as a class can retard foetal skeletal development (fully reversible) and cause discolouration of the deciduous teeth. However, evidence from human literature suggests that doxycycline is less likely to cause these abnormalities than other tetracyclines. Use only according to the benefit/risk assessment by the responsible veterinarian.

#### Interaction with other medicinal products and other forms of interaction:

Do not administer concurrently with bactericidal antibiotics such as penicillins and cephalosporins.

Oral absorbents and substances containing multivalent cations such as antacids and iron salts should not be used from 3 hours before to 3 hours after the administration of doxycycline as they reduce doxycycline availability. The half-life of doxycycline is reduced by concurrent administration of antiepileptic drugs such as phenobarbital and phenytoin.

#### Overdose (symptoms, emergency procedures, antidotes):

In cases of overdose no symptoms are to be expected other than those mentioned as adverse events.

#### Incompatibilities:

Not applicable.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

### **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

### **15. OTHER INFORMATION**

Aluminium - PVC/PE/PVDC blister

Package sizes:

Cardboard box of 3 blisters of 10 tablets  
Cardboard box of 5 blisters of 10 tablets  
Cardboard box of 10 blisters of 10 tablets  
Cardboard box of 1 blister of 30 tablets  
Cardboard box of 5 blisters of 30 tablets  
Cardboard box of 10 blisters of 30 tablets  
Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

#### **België/Belgique/Belgien**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

#### **Magyarország**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

#### **Danmark**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

#### **Österreich**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

#### **Deutschland**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

#### **Nederland**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**España**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**France**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**Ireland**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**Italia**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**Polska**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**Portugal**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**United Kingdom**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**Suomi/Finland**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

**Sverige**

CP-Pharma Handelsgesellschaft mbH  
Ostlandring 13  
DE- 31303 Burgdorf  
Tel.: +49 5136 60660  
info@cp-pharma.de.

Approved 14 July 2020

