

PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Cardboard box}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Vetbromide 600 mg tablets

2. STATEMENT OF ACTIVE SUBSTANCES

Each tablet contains:

Active substances:

Potassium bromide 600 mg

3. PACKAGE SIZE

60 tablets

120 tablets

4. TARGET SPECIES

Dogs



5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Oral use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Store below 30 °C.

After piercing a blister, replace unused tablet parts into the blister and place the blister back into the carton. Remaining tablet portions should be given at the next administration.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

DOMES PHARMA

14. MARKETING AUTHORISATION NUMBERS

Vm 54982/4008

15. BATCH NUMBER

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS { PVC/PVDC/Aluminium blisters }**

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Vetbromide



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

600 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Vetbromide 600 mg tablets for dogs

2. Composition

Each tablet contains:

Active substances:

Potassium bromide 600 mg

Round, white tablet with 2 scored lines on each side. Tablets can be divided into 2 or 4 equal parts.

3. Target species

Dogs



4. Indications for use

An antiepileptic agent for use in the control of idiopathic epileptic seizures, either as a single agent or as an adjunct to phenobarbital in the control of refractory cases of idiopathic epilepsy.

5. Contraindications

Do not use in dogs with severe renal insufficiency.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

6. Special warnings

Special warnings:

The concentration of bromide in serum, the clinical response and the therapeutic effect of administration of the veterinary medicinal product may vary between individuals (see section 8). The presence of cluster seizures/status epilepticus, due to the severity of the seizure activity, is often associated with poor response to anti-epileptic treatment. In these cases, remission (seizure freedom) may be difficult to achieve.

For dogs with normal hepatic function, phenobarbital is generally considered the first-choice antiepileptic drug. However, potassium bromide can be recommended as

alternative, especially in dogs with hepatic dysfunction or in dogs with concurrent disorders requiring life-long administration of potentially hepatotoxic medications, since potassium bromide is not metabolised in the liver.

A high chloride intake can increase the elimination of bromide (see section 6 Interaction). An increase in the dog's salt intake may require an adjustment in bromide dose. The salt content of a dog's diet during the treatment period should be maintained at a stable level. It is advisable not to change the dog's diet during therapy.

Special precautions for safe use in the target species:

Do not abruptly discontinue therapy as this may precipitate seizures.

This veterinary medicinal product should be used with caution in dogs with mild or moderate renal insufficiency, since excretion of bromide is reduced (see also section 5). To prevent bromide accumulation and a relative overdose of bromide (see section 6 Overdose), administer a reduced dose and monitor the serum bromide concentration closely (see section 8).

A reduction in chloride intake (low sodium diet) can increase the likelihood of adverse reactions or bromide intoxication (see section 6 Interaction and Overdose).

Close monitoring for adverse reactions is advisable at higher serum bromide concentrations.

Administration on an empty stomach may induce vomiting.

Dogs weighing less than 10 kg cannot be accurately dosed with the recommended starting dose for adjunctive treatment of 15 mg/kg twice daily, as the minimum dose achievable by division of the veterinary medicinal product is 150 mg (see section 8).

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product may cause eye-irritation. Avoid hand-to-eye contact. If the veterinary medicinal product comes into contact with the eyes, rinse immediately and thoroughly with clean water.

This veterinary medicinal product may be harmful upon ingestion, and cause adverse effects such as nausea and vomiting. Avoid oral ingestion including hand-to-mouth contact. To avoid accidental ingestion, particularly by a child, unused tablet parts should be returned to the open blister space and inserted back into the carton. Store in a closed cabinet. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands thoroughly, immediately after breaking or handling tablets.

To the physician:

An intravenous administration of isotonic sodium chloride (0.9%) will rapidly eliminate bromide ions in humans.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Laboratory studies have not produced any evidence of adverse effects of potassium bromide on reproduction at non-maternotoxic doses. Use only according to the benefit-risk assessment by the responsible veterinarian.

Potassium bromide crosses the placental barrier. Since bromide may be excreted into milk, monitor suckling puppies for somnolence/sedative effects; if necessary, consider early weaning, or an artificial suckling method.

Interaction with other medicinal products and other forms of interaction:

Due to the competition between chloride ions and bromide ions for reabsorption by the kidneys, any major change in chloride intake can modify serum bromide concentrations which are directly correlated to treatment efficacy and the occurrence of adverse reactions. A reduction in chloride intake (low sodium diet) can cause a rise in serum bromide levels and increase the likelihood of bromide intoxication (see section 6 Overdose). An increase in chloride intake (high salt diet) can cause a fall in serum bromide levels which could lead to seizures. Where possible, the diet of treated dogs should therefore not be altered. Seek veterinary advice before making any change to the dog's diet.

On biochemistry profiles serum chloride concentrations are often falsely elevated because the assays cannot distinguish between chloride and bromide ions.

Loop diuretics such as furosemide can increase bromide excretion and lower the efficacy of the treatment (risk of recurrence of seizures) if the dose is not adjusted. Administration of fluids or drug formulations containing chloride can lower serum bromide concentrations.

Bromide is synergistic with other GABA-ergic drugs such as phenobarbital.

Overdose:

Clinical signs of bromide toxicity (e.g. ataxia, somnolence) can occur in dogs with renal insufficiency or when a very high dose of bromide is administered. If overdose is suspected, the dosage should be reduced immediately, with close monitoring of serum bromide concentrations in order to establish an appropriate therapeutic concentration. Dose and serum bromide levels at which intolerance is observed vary between dogs. In cases of overdose requiring medical attention, administer 0.9% sodium chloride solution intravenously to reduce serum bromide concentrations.

7. Adverse events

Dogs:

Very common (>1 animal / 10 animals treated):

Polyphagia (increased appetite) (with or without weight gain)¹, polydipsia (increased thirst) (with or without polyuria – excessive urine production)¹,

Hind limb weakness¹, ataxia (incoordination, stumbling gait)¹, sedation¹

Loose stools¹, diarrhoea¹, vomiting¹

Common (1 to 10 animals / 100 animals treated):

Apathy (lack of interest, listless)¹, depression¹, hyperexcitation¹, aggression¹

Snoring (abnormal)¹, cough¹

Loss of appetite¹

Urinary incontinence¹ and/or urination (nocturnal)¹

Uncommon (1 to 10 animals / 1 000 animals treated):

Skin disorders¹

Undetermined frequency:

High pancreatic-specific lipase (cPLi)²

Low thyroxine (T4)³, Low free thyroxine (FT4)³

¹ These adverse reactions may disappear after the first stage of treatment but may also persist in dogs on higher doses of therapy. In these cases, symptoms usually disappear following a reduction in dose. If the dog appears too sedated, assess the serum concentrations of bromide and, if applicable, phenobarbital to determine whether the dose of either should be reduced.

If potassium bromide dose is reduced, serum bromide concentrations should be monitored in order to ensure that they fall within the therapeutic range.

² Although pancreatitis has been suggested to occur in association with the administration of bromide and/or phenobarbital, there is no conclusive evidence of a direct causal relationship between bromide administration and the development of pancreatitis in dogs.

³ Treating dogs with potassium bromide can cause a decrease in T4 plasma concentration, although this is not necessarily clinically relevant.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, routes and method of administration

Oral use.

Administer twice daily with food in order to reduce the risk of gastrointestinal irritation.

In dogs with severe and frequent seizures or when a dog is being switched rapidly from phenobarbital to potassium bromide, a loading dose of 60 mg/kg bodyweight

twice daily, for 5 days (equivalent to a total daily dose of 120 mg/kg) can be administered in order to quickly reach therapeutic serum concentrations.

The maintenance dose should be titrated to the individual dog as the required dosage and therapeutic serum bromide concentration may vary between animals and depends on the nature and severity of the underlying disease.

Monotherapy:

The recommended starting dose is 30 mg/kg bodyweight twice daily (equivalent to a total daily dose of 60 mg/kg).

Adjunctive treatment, in combination with phenobarbital:

The recommended starting dose is 15 mg/kg bodyweight twice daily (equivalent to a total daily dose of 30 mg/kg). Use in dogs with a bodyweight of less than 10 kg should be subject to a risk/benefit assessment, see section 6.

At the beginning of treatment, bromide serum concentrations should be checked regularly, e.g. 1 week and 1 month after the loading period and three months after treatment initiation at maintenance dosage. Therapeutic serum levels vary between 1000 mg/L to 3000 mg/L when potassium bromide is used as monotherapy and between 800 mg/L and 2000 mg/L, when used as adjunctive treatment. Close monitoring for side effects is advisable, particularly when serum bromide concentrations have reached the upper limit of the therapeutic range for monotherapy.

It is recommended to administer at least half of the initial starting dose to dogs with mild or moderate renal insufficiency, with more frequent monitoring of serum bromide levels (see section 6).

If the clinical response is not satisfactory or if adverse reactions occur, the dose may be adjusted based on the dog's serum bromide levels. Serum concentrations should be measured after each dose adjustment once steady state serum levels have been reached (typically 3 months after a change), unless earlier evaluation is necessary. Long term monitoring of serum bromide concentrations should be performed as clinically justified by the individual case.

9. Advice on correct administration

Not applicable.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Store below 30 °C.

After piercing a blister, replace unused tablet parts into the blister and place the blister back into the carton. Remaining tablet portions should be given at the next administration.

Do not use this veterinary medicinal product after the expiry date which is stated on the blister and carton after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Marketing authorisation numbers:

Vm 54982/4008

Pack sizes:

Cardboard box containing 60 or 120 tablets (4 or 8 blisters with 15 tablets each)

Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder:

Domes Pharma
3 Rue Andre Citroën
63430 Pont-Du-Chateau
France

Manufacturer responsible for batch release:

EUROPHARTECH
34 rue Henri Matisse
63370 Lempdes
France

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

TVM UK Animal Health Ltd.
Building B, Kirtlington Business centre,
Kirtlington OX5 3JA
United Kingdom
Tel 0800 038 5868
Email: help@tvm-uk.com

17. Other information

POM-V

Gavin Hall
Approved: 05 March 2026