

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARTON BOX

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Fipronil 268 mg spot-on solution for large dogs
Fipronil

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Contains: 268mg Fipronil

3. PHARMACEUTICAL FORM

Spot-on solution

4. PACKAGE SIZE

1 pipette, 2*, 3*, 4*, 5* or 6* pipettes
[*Delete as appropriate]

5. TARGET SPECIES

Dog

6. INDICATION(S)

For the treatment of flea (*Ctenocephalides spp.*) and tick (*Rhipicephalus sanguineus* and *Ixodes ricinus*) infestations.

Bob Martin Clear Spot on has a persistent insecticidal efficacy for up to 8 weeks against fleas (*Ctenocephalides spp.*) and a persistent acaricidal efficacy for up to 4 weeks against ticks (*Rhipicephalus sanguineus*, *Ixodes ricinus* and *Dermacentor reticulatus*).

Fleas will be killed within 24 h. Ticks will usually be killed within 48 h after contact with Fipronil. However, if ticks of some species (*Dermacentor reticulatus*) are already present when the product is applied, all of the ticks may not be killed within the first 48 hours.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD), where this has been previously diagnosed by a veterinary surgeon.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

For external use only.

Animals should be weighed accurately prior to treatment.

One pipette per dog of 20 – 40kg.

For dogs over 60 kg, use two pipettes of 2.68 ml.

Please read the package leaflet before use for full instructions, including user warnings.



1. Remove the pipette from the blister packaging. Hold the pipette in an upright position and snap off the top section of the cap, thereby opening the pipette.
2. Part your dog's coat until the skin is visible. Place the tip of the pipette directly against the bared skin and squeeze gently. Empty approximately half the contents at each of two points along the dog's back, preferably at the base of the head and between the shoulder blades. Squeeze the pipette several times to ensure dosing is complete. Avoid applying the solution onto the fur and do not rub into the skin.
3. Application of the solution as directed minimises the possibility that the animal will lick the solution off. Do not allow pets to lick each other after treatment.

8. WITHDRAWAL PERIOD

9. SPECIAL WARNING(S), IF NECESSARY

Please read the package leaflet before use for full instructions, including user warnings.

Do not use on rabbits!

10. EXPIRY DATE

EXP: end {month/year}

11. SPECIAL STORAGE CONDITIONS

Store below 25°C, in a dry place in the original packaging.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of waste materials in accordance with local requirements. Fipronil may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only.
For external use only.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Pets Choice Limited
Brentwood House
Lower Philips Road
Whitebirk Industrial Estate
Blackburn
Lancashire
BB1 5UD
United Kingdom

16. MARKETING AUTHORISATION NUMBER(S)

Vm 52797/5007

17. MANUFACTURER’S BATCH NUMBER

Batch no: {number}

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

PVC BLISTER WITH FOIL SEAL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Fipronil 268 mg spot-on solution for large dogs
Fipronil

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Pets Choice Limited
Brentwood House
Lower Philips Road
Whitebirk Industrial Estate
Blackburn
Lancashire
BB1 5UD
United Kingdom

3. EXPIRY DATE

EXP: end {month/year}

4. BATCH NUMBER

Batch No. {number}

5. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

PIPETTE LABEL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Fipronil 268 mg spot-on solution for large dogs

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

268 mg.

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

Not applicable – single dose pipette.

4. ROUTE(S) OF ADMINISTRATION

Spot-on

5. WITHDRAWAL PERIOD

6. BATCH NUMBER

Batch no: {number}

7. EXPIRY DATE

EXP: end {month/year}

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PACKAGE LEAFLET
Bob Martin Fipronil

Spot-on solution for small, medium, large and extra large dogs

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER
AND OF THE MANUFACTURING AUTHORISATION HOLDER
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Pets Choice Limited
Brentwood House
Lower Philips Road
Whitebirk Industrial Estate
Blackburn
Lancashire
BB1 5UD
United Kingdom

Manufacturer for batch release:

Pets Choice Ltd
8 Wemberham Lane
Yatton
North Somerset
BS49 4BS
United Kingdom

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin Fipronil 67 mg spot-on solution for small dogs
Bob Martin Fipronil 134 mg spot-on solution for medium dogs
Bob Martin Fipronil 268 mg spot-on solution for large dogs
Bob Martin Fipronil 402 mg spot-on solution for extra large dogs
Fipronil

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Dog	Weight (kg)	Unit dose	Fipronil	Excipients: Butylhydroxanisole	Excipients: Butylhydroxytoluene
Small dogs	2 - 10	0.67 ml	67 mg	0.134 mg/pipette	0.067 mg/ pipette
Medium dogs	10-20	1.34 ml	134 mg	0.268 mg/ pipette	0.134 mg/ pipette
Large dogs	20-40	2.68 ml	268 mg	0.536 mg/ pipette	0.268 mg/ pipette
Extra large dogs	40-60	4.02 ml	402 mg	0.804 mg/ pipette	0.402 mg/ pipette

4. INDICATION(S)

For the treatment of flea (*Ctenocephalides spp.*) and tick (*Rhipicephalus sanguineus* and *Ixodes ricinus*) infestations.

Bob Martin Clear Spot on has a persistent insecticidal efficacy for up to 8 weeks against fleas (*Ctenocephalides spp.*) and a persistent acaricidal efficacy for up to 4 weeks against ticks (*Rhipicephalus sanguineus*, *Ixodes ricinus* and *Dermacentor reticulatus*).

Fleas will be killed within 24 h. Ticks will usually be killed within 48 h after contact with Fipronil. However, if ticks of some species (*Dermacentor reticulatus*) are already present when the product is applied, all of the ticks may not be killed within the first 48 hours.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD), where this has been previously diagnosed by a veterinary surgeon.

5. CONTRAINDICATIONS

Do not use on puppies less than 2 months old and /or weighing less than 2kg for small dogs, 10 kg for medium dogs, 20 kg for large dogs and 40kg for extra large dogs, respectively.

Do not use on sick (e.g. systemic diseases, fever...) or convalescent animals.

Do not use in rabbits, as adverse reactions and even death could occur.

This product is specifically developed for dogs. Do not use in cats, as this could lead to overdosing.

Do not use on animals with known hypersensitivity to the active substance or to any of the excipients.

6. ADVERSE REACTIONS

If licking occurs, a brief period of hypersalivation may be observed due mainly to the nature of the carrier.

Among the extremely rare suspected adverse reactions, transient cutaneous reactions on the application site (skin discoloration, local alopecia, pruritus, erythema) and general pruritus or alopecia have been reported after use.

Exceptionally, hypersalivation, reversible neurologic symptoms (hyperesthesia,


depression, nervous symptoms), vomiting or respiratory symptoms have been observed after use.

7. TARGET SPECIES

Dog.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

External use only. Administer by topical application to the skin 1 pipette per animal, or if over 60kg, 2 pipettes of 2.68ml.



Remove the pipette from the blister packaging. Hold the pipette in an upright position and snap off the top section of the cap, thereby opening the pipette.

Part your dog's coat until the skin is visible. Place the tip of the pipette directly against the bared skin and squeeze gently. Empty approximately half the contents at each of two points along the dog's back, preferably at the base of the head and between the shoulder blades. Squeeze the pipette several times to ensure dosing is complete. Avoid applying the solution onto the fur and do not rub into the skin.

Application of the solution as directed minimises the possibility that the animal will lick the solution off. Do not allow pets to lick each other after treatment.

9. ADVICE ON CORRECT ADMINISTRATION

Application of the solution as directed minimises the possibility that the animal will lick the solution off. Do not allow pets to lick each other after treatment.

Care should be taken to avoid excessive wetting of the hair with the product since this will cause a sticky appearance of hairs at the treatment spot. However, should this occur, it will usually disappear within 24-48 hours post application. Crystals may be seen on the hair and slight scaling may occur within 24-48 hours after application.

Treatment schedule:

For optimal control of flea and/or tick infestation the treatment schedule can be based on the local epidemiological situation.

The minimum interval between two treatments is 4 weeks.

10. WITHDRAWAL PERIOD

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.
Store below 25°C, in a dry place in the original packaging.

12. SPECIAL WARNING(S)

Avoid frequent swimming or shampooing the animal because the maintenance of effectiveness of the product in these cases has not been tested.

The product does not prevent ticks from attaching to the animals. If the animal has been treated prior to exposure to the ticks, the ticks will be killed in the first 24-48 hours after attachment. This will usually be prior to engorgement, minimising but not excluding the risk of transmission of diseases. Once dead, ticks will often drop off the animal, but any remaining ticks may be removed with a gentle twist.

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.

For optimal control of flea infestation in multi-pet household, all dogs and cats in the household should be treated with a suitable insecticide.

When treating infestations, all in-contact dogs should be treated with an appropriate product at the same time.

When used as part of a strategy for the treatment of Flea Allergy Dermatitis, monthly applications to the allergic patient and to other dogs and cats in the household are recommended.

Special precautions for use

i) Special precautions for use in animals

Animals should be weighed accurately prior to treatment.

Avoid contact with the animal's eyes. In the case of accidental eye contact, immediately and thoroughly flush the eyes with water.

It is important to make sure that the product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment.

Do not allow pets to lick each other after treatment

Do not apply the product on wounds or damaged skin.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product can cause mucous membrane and eye irritation. Therefore, contact between the product and the mouth or eyes should be avoided.

In the case of accidental eye contact, immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the physician.

Do not smoke, drink or eat during application.

Avoid contents coming into contact with the fingers. If this occurs, wash hands with soap and water. Wash hands after use.

Animals or people with a known hypersensitivity to fipronil or any of the other ingredients should avoid contact with the product.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Keep pipettes in the original packaging and dispose of used pipettes immediately.

iii) Environmental warning

Fipronil may adversely affect aquatic organisms. Dogs should not be allowed to swim in water courses for 2 days after application.

iv) Other precautions

The product may have adverse effects on painted, varnished or other household surfaces or furnishings.

Use during pregnancy and lactation

Laboratory studies using fipronil have not shown any evidence of teratogenic or embryotoxic effect. Studies have not been carried out with this product in pregnant and lactating bitches. Use in pregnancy and lactation only in accordance with professional veterinary advice and a benefit/risk assessment.

Interaction with other medicinal products and other forms of interaction

None known.

Overdose (symptoms, emergency procedures, antidotes), if necessary

No adverse effects were observed in target animal safety studies in 2 month-old puppies, growing dogs and dogs weighing about 2 kg treated with 5 times the therapeutic dose once a month for 3 consecutive months. The risk of adverse effects (see section 6) may increase in cases of overdose.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Fipronil may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

July 2021

15. OTHER INFORMATION

Blister cards or boxes of 1, 2, 3, 4, 5 or 6 pipettes.

Not all pack sizes may be marketed.

For animal treatment only

Approved 07 July 2021

A handwritten signature in black ink, appearing to read "A. Hunter.", is positioned below the approval date.