

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Plastic box of 10 bottles of lyophilisate and 10 bottles of solvent; Plastic box of 50 bottles of lyophilisate and 50 bottles of solvent}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Purevax RCP lyophilisate and solvent for suspension for injection

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Per dose of 1 ml or 0.5 ml:

FHV (F2 strain)	$\geq 10^{4.9}$ CCID <sub>50</sub>
FCV (431 and G1 strains)	$\geq 2.0$ ELISA U.
FPV (PLI IV)	$\geq 10^{3.5}$ CCID <sub>50</sub>

**3. PACKAGE SIZE**

Lyophilisate (10 x 1 dose) + solvent (10 x 1 ml)  
Lyophilisate (50 x 1 dose) + solvent (50 x 1 ml)  
Lyophilisate (10 x 1 dose) + solvent (10 x 0.5 ml)  
Lyophilisate (50 x 1 dose) + solvent (50 x 0.5 ml)

**4. TARGET SPECIES**

Cats.

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Subcutaneous use.

**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

Exp. {dd/mm/yyyy}  
Once reconstituted use immediately.

**9. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.  
Protect from light.  
Do not freeze.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Vetmedica GmbH

**14. MARKETING AUTHORISATION NUMBERS**

Vm 04491/5053

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

POM-V

Veterinary medicinal product subject to prescription

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING  
UNITS {Lyophilisate bottle}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Purevax RCP



**2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE  
SUBSTANCES**

1 dose

1 ml or 0.5 ml

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {dd/mm/yyyy}

**5. ROUTE(S) OF ADMINISTRATION**

**6. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING**  
**UNITS {Solvent bottle}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Purevax RCP solvent



**2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

1 ml or 0.5 ml

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {dd/mm/yyyy}

**5. ROUTE(S) OF ADMINISTRATION**

**6. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

## **PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Purevax RCP lyophilisate and solvent for suspension for injection

### **2. COMPOSITION**

Per dose of 1 ml or 0.5 ml:

#### **Active substances:**

##### **Lyophilisate:**

Attenuated feline rhinotracheitis herpesvirus (FHV F2 strain)	$\geq 10^{4.9}$ CCID <sub>50</sub> <sup>1</sup>
Inactivated feline calicivirus (FCV 431 and G1 strains) antigens	$\geq 2.0$ ELISA U.
Attenuated feline panleucopenia virus (PLI IV)	$10^{3.5}$ CCID <sub>50</sub> <sup>1</sup>

<sup>1</sup> cell culture infective dose 50%

##### **Solvent:**

Water for injections q.s. 1 ml or 0.5 ml

Lyophilisate: friable pellet, homogeneous from beige to white.

Solvent: clear colourless liquid.

### **3. TARGET SPECIES**

Cats.

### **4. INDICATIONS FOR USE**

Active immunisation of cats aged 8 weeks and older:

- against feline viral rhinotracheitis to reduce clinical signs,
- against calicivirus infection to reduce clinical signs,
- against feline panleucopenia to prevent mortality and clinical signs.

Onset of immunity: 1 week after primary vaccination course.

Duration of immunity: 1 year after primary vaccination course and 3 years after the last re-vaccination.

### **5. CONTRAINDICATIONS**

None.

### **6. SPECIAL WARNINGS**

Vaccinate healthy animals only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy and lactation:

Do not use during the whole pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:

Safety and efficacy data are available which demonstrate that this vaccine can be mixed and administered with Boehringer Ingelheim non-adjuvanted vaccine against feline leukaemia and/or administered the same day but not mixed with Boehringer Ingelheim adjuvanted vaccine against rabies.

Safety and efficacy data are available which demonstrate that this vaccine can be mixed and administered with Boehringer Ingelheim non-adjuvanted vaccine against rabies.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the products mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

Overdose (symptoms, emergency procedures, antidotes):

No adverse event other than those already mentioned in section on "Adverse events" have been observed, except hyperthermia that may exceptionally last 5 days.

Major incompatibilities:

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product and except those mentioned above.

## 7. ADVERSE EVENTS

Cats:

**Common (1 to 10 animals / 100 animals treated):**

Apathy, anorexia, and hyperthermia<sup>1</sup>.

Injection site reactions (pain, itching, oedema).<sup>2</sup>

**Uncommon (1 to 10 animals / 1 000 animals treated):**

Hypersensitivity reaction.<sup>3</sup>

**Very rare (<1 animal / 10 000 animals treated, including isolated reports):**

Emesis.<sup>4</sup>

<sup>1</sup> lasting usually for 1 or 2 days.

<sup>2</sup> slight pain at palpation, itching or limited oedema disappearing within 1 or 2 weeks at most.

<sup>3</sup> which may require appropriate symptomatic treatment.

<sup>4</sup> mostly within 24 to 48 hours.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package

leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

## **8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

Subcutaneous route.

After reconstitution of the lyophilisate with 0.5 ml or 1 ml of the solvent (depending on the presentation chosen), inject one dose of vaccine according to the following vaccination schedule:

Primary vaccination course:

- first injection: from 8 weeks of age,
- second injection: 3 to 4 weeks later.

Where high levels of maternal antibodies against rhinotracheitis, calicivirosis or panleucopenia components are expected to be present (e.g. in kittens of 9 to 12 weeks of age born from queens, which were vaccinated before pregnancy and/or with known or suspected previous exposure to the pathogen(s)), the primary vaccination course should be delayed until 12 weeks of age.

Revaccination:

- the first revaccination must be carried out one year after the primary vaccination course,
- subsequent revaccinations: at intervals of up to 3 years.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Reconstitute gently the vaccine in order to obtain a uniform suspension with limited foam formation.

Visual appearance after reconstitution: clear slightly yellow suspension.

## **10. WITHDRAWAL PERIODS**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C).

Protect from light.

Do not freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and the bottle after Exp.

Shelf life after reconstitution according to directions: use immediately.

## 12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater. Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## 13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

## 14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 04491/5053

Plastic box containing:

10 x 1 dose of lyophilisate and 10 x 1 ml of solvent or  
50 x 1 dose of lyophilisate and 50 x 1 ml of solvent or  
10 x 1 dose of lyophilisate and 10 x 0.5 ml of solvent or  
50 x 1 dose of lyophilisate and 50 x 0.5 ml of solvent.

Not all pack sizes may be marketed.

## 15. PID LINK (Do not print heading)

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

## 16. CONTACT DETAILS

Marketing authorisation holder:

Boehringer Ingelheim Vetmedica GmbH  
55216 Ingelheim/Rhein  
Germany

Manufacturer responsible for the batch release:

Boehringer Ingelheim Animal Health France SCS  
Laboratoire Porte des Alpes  
Rue de l'aviation  
69800 Saint-Priest  
France

Local representatives and contact details to report suspected adverse reactions:

United Kingdom (Great Britain)  
Boehringer Ingelheim Animal Health UK Ltd., United Kingdom  
Tel: + 44 1344 746957



## 17. OTHER INFORMATION

The product was shown to reduce excretion of feline calicivirus at onset of immunity and for one year after vaccination.

POM-V

For animal treatment only

*Gavin Hall*

Approved: 20 December 2024