

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**{CARDBOARD BOX}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Milprazon CHEWABLE 4 mg/10 mg film-coated tablets

*For memory stickers:*  
*milbemycin oxime/praziquantel*

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains: milbemycin oxime 4 mg and praziquantel 10 mg.

**3. PACKAGE SIZE**

2 tablets  
4 tablets  
48 tablets

**4. TARGET SPECIES**

Cats (small cats and kittens weighing at least 0.5 kg)



**5. INDICATIONS**

Flavoured broad spectrum anthelmintic

*For products not subject to veterinary prescription:*

Treatment of mixed infections by immature and adult tapeworms and roundworms

**6. ROUTES OF ADMINISTRATION**

Oral use.

*For products not subject to veterinary prescription:*

Dosage:

Body weight	Tablets
0.5 - 1 kg	½ tablet
> 1 - 2 kg	1 tablet

## 7. WITHDRAWAL PERIODS

## 8. EXPIRY DATE

Exp. {mm/yyyy}

Shelf life for halved tablets after first opening the immediate packaging: 6 months.

## 9. SPECIAL STORAGE PRECAUTIONS

Store in the original package in order to protect from moisture.  
Halved tablets should be stored below 25°C in the original blister and be used for the next administration.  
Keep the blister in the outer carton.

## 10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

## 11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

## 12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

## 13. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

## 14. MARKETING AUTHORISATION NUMBERS

Vm 01656/3079

**15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

{BLISTER}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Milprazon CHEWABLE



**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

4 mg/10 mg

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET

### 1. Name of the veterinary medicinal product

Milprazon CHEWABLE 4 mg/10 mg film-coated tablets for small cats and kittens weighing at least 0.5 kg

### 2. Composition

Each film-coated tablet contains:

#### Active substances:

Milbemycin oxime	4 mg
Praziquantel	10 mg

#### Excipients:

Titanium dioxide (E171): 0.21 mg  
Iron Oxide, yellow (E172): 0.18 mg  
Iron Oxide, red (E172): 0.02 mg

Brownish orange, oval, biconvex film-coated tablets with score line on one side.  
The tablets can be divided into halves.

### 3. Target species

Cats (small cats and kittens weighing at least 0.5 kg).



### 4. Indications for use

Treatment of mixed infections by immature and adult tapeworms and roundworms of the following species:

- Tapeworms:

*Dipylidium caninum*

*Taenia* spp.

*Echinococcus multilocularis*

- Roundworms:

*Ancylostoma tubaeforme*

*Toxocara cati*

Prevention of heartworm disease (*Dirofilaria immitis*) if concomitant treatment against tapeworms is indicated.



## **5. Contraindications**

Do not use in cats of less than 6 weeks of age and/or weighing less than 0.5 kg  
Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

## **6. Special warnings**

### Special warnings:

It is recommended to treat all the animals living in the same household concomitantly.

In order to effectively control worm infection local information about presence and susceptibility of the parasites (epidemiological information) and the living conditions of the cat should be taken into account, and it is recommended to seek professional (e.g. veterinary) advice.

When infection with tapeworm *D. caninum* is present, concomitant treatment against intermediate hosts, such as fleas and lice, should be considered to prevent re-infection.

Parasite resistance to any particular class of anthelmintic (drugs acting against worms) may develop following frequent, repeated use of an anthelmintic of that class.

### Special precautions for safe use in the target species:

No studies have been performed with severely debilitated cats or individuals with seriously compromised kidney or liver function. The veterinary medicinal product is not recommended for such animals or only according to a benefit/risk assessment by the responsible veterinarian.

Ensure cats and kittens weighing between 0.5 kg and  $\leq 2$  kg receive the appropriate tablet strength (4 mg milbemycin oxime/10 mg praziquantel) and the appropriate dose (1/2 or 1 tablet) for the corresponding weight band (1/2 tablet for cats weighing 0.5 to 1 kg; 1 tablet for cats weighing  $>1$  to 2 kg).

As the tablets are flavoured, they should be stored in a safe place out of the reach of animals.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental ingestion of a tablet by a child may be harmful. In order to prevent children from accessing the veterinary medicinal product, tablets should be administered and stored out of sight and reach of children.

Part tablets should be returned to the open blister pocket and inserted into the outer carton.

In the event of accidental ingestion of one or more tablets, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

### Other precautions:

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (WOAH), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority (e.g. experts or institutes of parasitology).

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Fertility:

Can be used in breeding animals.

Interaction with other medicinal products and other forms of interaction:

No interactions were observed when the recommended dose of the macrocyclic lactone selamectin was administered during treatment with milbemyacin oxime and praziquantel at the recommended dose. In the absence of further studies, caution should be taken in the case of concurrent use of the veterinary medicinal product and other macrocyclic lactones. Also, no such studies have been performed with reproducing animals.

Overdose:

In case of overdose, in addition to adverse events observed at the recommended dose (see "Adverse reactions"), drooling may be observed. This sign will usually disappear spontaneously within a day.

## **7. Adverse events**

Cats:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Hypersensitivity reaction; Systemic disorders (e.g. lethargy)*; Neurological disorders (e.g. ataxia (incoordination) and muscle tremors)*; Digestive tract disorders (e.g. emesis (vomiting) and diarrhoea)*.
--	--

\*Especially in young cats.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>  
e-mail: [adverse.events@vmd.gov.uk](mailto:adverse.events@vmd.gov.uk)

## **8. Dosage for each species, routes and method of administration**

Oral use.

To ensure a correct dosage, body weight should be determined as accurately as possible.

Minimum recommended dose rate: 2 mg of milbemyacin oxime and 5 mg of praziquantel per kg are given orally as a single dose.

Depending on the bodyweight of the cat, the practical dosing is as follows:

<b>Body weight</b>	<b>Film-coated tablets for small cats and kittens</b>
0.5 - 1 kg	½ tablet
more than 1 - 2 kg	1 tablet

### **9. Advice on correct administration**

The veterinary medicinal product should be administered with or after some food. Doing so ensures optimum protection against heartworm disease.

The veterinary medicinal product can be inserted into a programme for prevention of heartworm disease if at the same time treatment against tapeworms is indicated. For the prevention of heartworm disease: the veterinary medicinal product kills *Dirofilaria immitis* larvae up to one month after their transmission by mosquitoes. For regular prevention of heartworm disease the use of a monosubstance is preferred.

### **10. Withdrawal periods**

Not applicable.

### **11. Special storage precautions**

Keep out of the sight and reach of children.

Store in the original package in order to protect from moisture. This veterinary medicinal product does not require any special temperature storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the blister and carton after Exp. The expiry date refers to the last day of that month.

Shelf life for halved tablets after first opening the immediate packaging: 6 months. Halved tablets should be stored below 25°C in the original blister and be used for the next administration.

Keep the blister in the outer carton.

### **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

This veterinary medicinal product should not enter water courses as milbemycin oxime may be dangerous for fish and other aquatic organisms.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with

any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

### **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription (BG, CZ, DE, EE, ES, UK (NI), HR, HU, IE, IT, LV, PL, PT, RO, SK).

Veterinary medicinal product not subject to prescription (BE, LT, NL).

Veterinary medicinal product subject to prescription except for some pack sizes (FR, SI).

### **14. Marketing authorisation numbers and pack sizes**

Vm 01656/3079

Cardboard box with 1 blister of 2 tablets.

Cardboard box with 1 blister of 4 tablets.

Cardboard box with 12 blisters, each blister contains 4 tablets (total 48 tablets).

Not all pack sizes may be marketed.

### **15. PID LINK (Do not print heading)**

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

### **16. Contact details**

Marketing authorisation holder and manufacturer responsible for batch release and contact details to report suspected adverse reactions:

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

Manufacturer responsible for batch release:

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

KRKA - FARMA d.o.o., V. Holjevca 20/E, 10450 Jastrebarsko, Croatia

TAD Pharma GmbH, Heinz-Lohmann-Straße 5, 27472 Cuxhaven, Germany

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**17. Other information**

*Gavin Hall*  
Approved: 26 November 2024