PARTICULARS TO APPEAR ON IMMEDIATE PACKAGE
POLYETHLENE BACK PACK CONTAINER – 1 LITRE

1. NAME OF THE VETERINARY MEDICINAL PRODUCT
CLIK 50 mg/ml Pour-On Suspension

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES
Each ml contains:

Active substance:
Dicyclanil 50 mg/ml

Excipients:

Ponceau 4R (E124)	0.05 mg
Methyl parahydroxybenzoate (E218)	1.50 mg
Propyl parahydroxybenzoate (E216)	3.00 mg
Butylated hydroxytoluene (E321)	0.50 mg.

3. PACKAGE SIZE
0.8 litres

4. TARGET SPECIES
Sheep

5. INDICATION(S)

6. ROUTES OF ADMINISTRATION
Pour on.

7. WITHDRAWAL PERIODS
Meat and offal: 40 days.
Not authorised for use in animals producing milk for human consumption.

8. EXPIRY DATE
Exp. {mm/yyyy}
Once opened use within 1 year.
Once opened use by….

9. SPECIAL STORAGE PRECAUTIONS
Protect from frost.
Store in the original container.
Keep the container tightly closed.
Protect from direct sunlight.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”
Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”
For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”
Keep out of the sight and reach of children.

13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Elanco GmbH
Heinz-Lohmann Strasse 4
Groden
27472 Cuxhaven
Germany

14. MARKETING AUTHORISATION NUMBERS

Vm 52127/5093

15. BATCH NUMBER

Lot {number}

16. SPECIAL WARNING(S), IF NECESSARY

For external use only.
If you notice any serious effects or other effects not mentioned on this label, please inform your veterinary surgeon.
Special precautions for use in animals.
The product is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm. Established strikes may require a separate treatment with a knockdown insecticide.
It is recommended that animals with dirty back-ends are dagged prior to application. If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.
Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.
The use of the product has harmful effects on dung flies.
Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is a **serious** risk to aquatic life if this advice is not followed.
Special precautions to be taken by the person administering the veterinary medicinal product to animals
• Operators should wear synthetic rubber gloves, a face mask and PVC trousers when applying the product.
• In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.
• In case of eye contact rinse eyes with clean water for several minutes and contact physician for advice.
• In case of ingestion call for medical advice immediately and show the package leaflet or the label to the physician.

- Always wash hands with soap and water after handling sheep and before eating and drinking or smoking.
- Wash hands and exposed skin after working with recently treated sheep.
- Do not eat, drink or smoke whilst using the product.

<u>Handling sheep in the weeks following treatment:</u>
• Do not shear sheep in the 3 months after treatment.
• Handle sheep as little as possible after treatment as residues remain on the fleece for some weeks. • If you need to handle sheep after treatment, wear coveralls and Wellington boots. If sheep are wet also wear waterproof trousers and coat.

<u>Special precautions for the protection of the environment:</u>
The use of the veterinary medicinal product has harmful effects on dung flies.
Treated sheep must be kept away from watercourses for at least one hour after treatment. There is serious risk to aquatic life if the advice is not followed.

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater.
The product should not enter water courses as dicyclanil may be dangerous for fish and other aquatic organisms.
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.
Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help protect the environment.

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

POM-VPS (To be supplied only on veterinary prescription)

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

1 litre container

1. NAME OF THE VETERINARY MEDICINAL PRODUCT
CLIK 50 mg/ml Pour-On Suspension for Sheep.

2. COMPOSITION
Each ml contains:
Active Substance:
Dicyclanil 50 mg

Excipients:

Methyl Parahydroxybenzoate (E218)	1.50 mg
Proply Parahydroxybenzoate (E216)	3.00 mg
Butylated Hydroxytoluene (E321)	0.50 mg
Ponceau 4R (E124) Sodium	0.05 mg

Pink-colouredpour-on suspension

3. TARGET SPECIES
Sheep

4. INDICATIONS FOR USE
Prevention of blowfly strike due to *Lucilia sericata* on sheep.
Prevention of blowfly strike due to *Wohlfahrtia magnifica* on sheep.

5. CONTRAINDICATIONS
Do not use in sheep producing milk for human consumption.
Do not use in case of hypersensitivity to the active substance or any of the excipients.

6. SPECIAL WARNING(S)
Special warnings:

Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy.
The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infestation based on its epidemiological features, for each flock.

Special precautions for safe use in the target species:

The veterinary medicinal product is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm.
Established strikes may require a separate treatment with a knockdown insecticide.
It is recommended that animals with dirty back-ends are dagged prior to application.
If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.
Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Operators should wear synthetic rubber gloves, a face mask and PVC trousers when applying the veterinary medicinal product.
In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.
In case of eye contact rinse eyes with clean water for several minutes and contact physician for advice.
In case of accidental ingestion seek medical advice immediately and show the package leaflet or the label to the physician.
Always wash hands with soap and water after handling sheep and before eating and drinking or smoking.
Wash hands and exposed skin after working with recently treated sheep. Do not eat, drink or smoke whilst using the veterinary medicinal product.

Handling sheep in the weeks following treatment:

Do not shear sheep in the 3 months after treatment.
Handle sheep as little as possible after treatment as residues remain on the fleece for some weeks.
If you need to handle sheep after treatment, wear coveralls and Wellington boots. If sheep are wet also wear waterproof trousers and coat.

Special precautions for the protection of the environment:

The use of the veterinary medicinal product has harmful effects on dung flies.
Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is **serious** risk to aquatic life if this advice is not followed.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation.
Use only accordingto the benefit-risk assessment by the responsible veterinarian.
Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic, or maternotoxic effects.

Interaction with other medicinal products and other forms of interaction:

None known.

Overdose:

An overdose of up to at least 5 times the recommended dose does not lead to any signs of local or systemic intolerance. An antidote is not known.

For Animal Treatment Only.

Keep out of sight and reach of children.

7. ADVERSE EVENTS

Target species: Sheep.
None known.
Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For external use only.
The veterinary medicinal product is applied according to the following recommendations:

Body weight (kg)	Dose Volume (ml)
10 – 20	20
21 – 30	25
31 – 50	30
>50	35

(Guide dose volumes correspond to 0.6 – 2 ml (30 – 100 mg dicyclanil) per kg bodyweight).

Shake the container well before use.

The veterinary medicinal product must be applied with a manual or automatic dosing gun (e.g. Elanco Pour-on gun), with a spray nozzle, which guarantees the correct spreading of the product on the fleece. Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray using a 4 stroke method as shown in the pictogram along the spine of the animal in a band at least 10 cm wide from the middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.



The veterinary medicinal product is administered once, before predicted Wohlfahrtia fly activity, or before or at the beginning of Lucilia fly activity. The veterinary medicinal

product will protect against blowfly strike caused by Wohlfahrtia or Lucilia flies for 16 weeks. In individual cases, a strike may occur earlier; therefore, it is good practice to check animals regularly for blowfly strike.
Do not shear sheep in the 3 months after treatment.

9. ADVICE ON CORRECT ADMINISTRATION
Please refer to section on “Special Warnings” for further advice on administration.

10. WITHDRAWAL PERIOD(S)
Meat and offal: 40 days
Not authorised for use in animals producing milk for human consumption.

11. SPECIAL STORAGE PRECAUTIONS
Keep out of the sight and reach of children.

Protect from frost.
Store in the original container.
Keep the container tightly closed.
Protect from direct sunlight.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp.. The expiry date refers to the last day of that month.
Shelf life after first opening the container: 1 year.
When the container is opened for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the container should be discarded should be determined. This discard date should be written in the space provided.

12. SPECIAL PRECAUTIONS FOR DISPOSAL
Medicines should not be disposed of via wastewater. The veterinary medicinal product should not enter water courses as dicyclanil may be dangerous for fish and other aquatic organisms.
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.
Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help protect the environment

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS
Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES
The pack is composed of pigmented white opaque polyethylene back pack container with blue polypropylene screw cap, containing 0.8, 2.2 or 5 litres of finished product.
Not all pack sizes may be marketed.
Vm 52127/5093

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder and contact details to report suspected adverse reactions:

Elanco GmbH
Heinz-Lohmann
Strasse 4 Groden 27472
Cuxhaven
Germany
PV.GBR@elancoah.com
Tel: +44 3308221732

Manufacturer responsible for batch release
Argenta Dundee Limited
Kinnoull Road
Dunsinane Industrial Estate
Dundee
DD2 3XR
United Kingdom

Elanco France S.A.S
26 Rue de la Chapelle
68330 Huningue
France

17. OTHER INFORMATION

MINIMUM PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING
Combined label and package leaflet

POLYETHYLENE BACK PACK CONTAINER
2.5 and 5 litre pack sizes

1. NAME OF THE VETERINARY MEDICINAL PRODUCT
CLIK 50 mg/ml Pour-On Suspension for Sheep.

2. COMPOSITION

Each ml contains:

Active substance:

Dicyclanil 50 mg

Excipients:

Methyl Parahydroxybenzoate (E218)	1.50 mg
Proply Parahydroxybenzoate (E216)	3.00 mg
Butylated Hydroxytoluene (E321)	0.50 mg
Ponceau 4R (E124) Sodium	0.05 mg

Pink-coloured pour-on suspension.

3. PACKAGE SIZE
2.2 litres.
5 litres.

4. TARGET SPECIES
Sheep.

5. INDICATIONS FOR USE

Indications for use

Prevention of blowfly strike due to *Lucilia sericata* on sheep.
Prevention of blowfly strike due to *Wohlfahrtia magnifica* on sheep.

6. CONTRAINDICATIONS

Contraindications

Do not use in sheep producing milk for human consumption.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

7. SPECIAL WARNINGS

Special warnings:

Special warnings:

Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy.
The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infestation based on its epidemiological features, for each flock.

Special precautions for safe use in the target species:

The veterinary medicinal product is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm.
Established strikes may require a separate treatment with a knockdown insecticide.
It is recommended that animals with dirty back-ends are dagged prior to application.
If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.
Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Operators should wear synthetic rubber gloves, a face mask and PVC trousers when applying the veterinary medicinal product.
In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.
In case of eye contact rinse eyes with clean water for several minutes and contact physician for advice.
In case of accidental ingestion seek medical advice immediately and show the package leaflet or the label to the physician.
Always wash hands with soap and water after handling sheep and before eating and drinking or smoking.
Wash hands and exposed skin after working with recently treated sheep. Do not eat, drink or smoke whilst using the veterinary medicinal product.

Handling sheep in the weeks following treatment:

Do not shear sheep in the 3 months after treatment.
Handle sheep as little as possible after treatment as residues remain on the fleece for some weeks.
If you need to handle sheep after treatment, wear coveralls and Wellington boots. If sheep are wet also wear waterproof trousers and coat.

Special precautions for the protection of the environment:

The use of the veterinary medicinal product has harmful effects on dung flies.
Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is **serious** risk to aquatic life if this advice is not followed.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation.
Use only accordingto the benefit-risk assessment by the responsible veterinarian.
Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic, or maternotoxic effects.

Interactions with other medicinal products and other forms of interaction:

None known.

Overdose:

An overdose of up to at least 5 times the recommended dose does not lead to any signs of local or systemic intolerance. An antidote is not known.

Special restrictions for use and special conditions for use:

Not applicable.

Major incompatibilities:

Not applicable.

8. ADVERSE REACTIONS

Adverse events

Target species: Sheep.

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system.

9. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Dosage for each species, routes and method of administration

For external use only.
The veterinary medicinal product is applied according to the following recommendations:

Body weight (kg)	Dose Volume (ml)
10 – 20	20
21 – 30	25
31 – 50	30
>50	35

(Guide dose volumes correspond to 0.6 – 2 ml (30 – 100 mg dicyclanil) per kg bodyweight).

Shake the container well before use.

The veterinary medicinal product must be applied with a manual or automatic dosing gun (e.g. Elanco Pour-on gun), with a spray nozzle, which guarantees the correct spreading of the product on the fleece. Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray using a 4 stroke method as shown in the pictogram along the spine of the animal in a band at least 10 cm wide from the middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.



The veterinary medicinal product is administered once, before predicted Wohlfahrtia fly activity, or before or at the beginning of Lucilia fly activity. The veterinary medicinal product will protect against blowfly strike caused by Wohlfahrtia or Lucilia flies for 16 weeks. In individual cases, a strike may occur earlier; therefore, it is good practice to check animals regularly for blowfly strike.
Do not shear sheep in the 3 months after treatment.

10. ADVICE ON CORRECT ADMINISTRATION
Advice on correct administration

Please refer to section on “Special Warnings” for further advice on administration.

11. WITHDRAWAL PERIOD
Withdrawal periods

Meat and offal: 40 days.
Not authorised for use in animals producing milk for human consumption.

12. SPECIAL STORAGE PRECAUTIONS
Special storage precautions
Keep out of the sight and reach of children.

Protect from frost.

Store in the original container.
Keep the container tightly closed.
Protect from direct sunlight.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp.. The expiry date refers to the last day of that month.
Shelf life after first opening the container: 1 year
When the container is opened for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the container should be discarded should be determined. This discard date should be written in the space provided.

13. SPECIAL PRECAUTIONS FOR DISPOSAL

Special precautions for disposal

Medicines should not be disposed of via wastewater.
The veterinary medicinal product should not enter water courses as dicyclanil may be dangerous for fish and other aquatic organisms.
Any unused veterinary medicinal product or waste materials derived from such veterinary medical product should be disposed of in accordance with local requirements.
Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help protect the environment.

14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.
POM-VPS

15. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Pack sizes

The pack is composed of pigmented white opaque polyethylene back pack container with blue polypropylene screw cap, containing 0.8, 2.2 or 5 litres of finished product.
Not all pack sizes may be marketed.
Vm 52127/5093

16. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the ‘Product Information Database’ on www.gov.uk.

17. CONTACT DETAILS

Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Elanco GmbH
Heinz-Lohmann Strasse 4
Groden
27472 Cuxhaven
Germany
PV.GBR@elancoah.com
Tel: +44 3308221732

Manufacturer responsible for batch release:
Argenta Dundee Limited
Kinnoull Road
Dunsinane Industrial Estate
Dundee
DD2 3XR
United Kingdom

Elanco France S.A.S
26 Rue de la Chapelle
68330 Huningue
France

18. OTHER INFORMATION

19. THE WORDS "FOR ANIMAL TREATMENT ONLY"
For animal treatment only

20. EXPIRY DATE
Exp. {mm/yyyy}
Once opened use within 1 year.
Once opened use by

21. BATCH NUMBER
Lot {number}

22. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"
Keep out of the sight and reach of children

Gavin Hall
Approved: 27 April 2026