

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Box**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Endogard Plus Flavour Tablets for dogs  
*Praziquantel, pyrantel embonate, febantel*

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each tablet contains:  
Praziquantel                      50 mg  
Pyrantel embonate                144 mg  
Febantel                              150 mg

**3. PHARMACEUTICAL FORM**

Tablets

The tablets can be divided into 2 or 4 equal parts..

**4. PACKAGE SIZE**

2 tablets  
4 tablets  
10 tablets  
30 tablets  
50 tablets  
100 tablets  
300 tablets

**5. TARGET SPECIES**

For dogs (small and medium size)

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**8. WITHDRAWAL PERIOD**

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP:

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of waste material in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

KRKA, d.d., Novo mesto  
Šmarješka cesta 6  
8501 Novo mesto  
Slovenia

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 01656/4017

**17. MANUFACTURER’S BATCH NUMBER**

Lot:

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blister**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Endogard Plus Flavour Tablets for dogs  
*Praziquantel, pyrantel embonate, febantel*

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

KRKA

**3. EXPIRY DATE**

EXP:

**4. BATCH NUMBER**

Lot:

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## PACKAGE LEAFLET FOR:

### Endogard Plus Flavour Tablets for dogs

#### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

KRKA, d.d., Novo mesto  
Šmarješka cesta 6  
8501 Novo mesto  
Slovenia

Manufacturer responsible for batch release:

KRKA - FARMA d.o.o., V. Holjevca 20/E, 10450 Jastrebarsko, Croatia  
Virbac S.A., 1ere Avenue, 2065M, LID, 06516 Carros Cedex, France  
KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

#### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Dehinel Plus Flavour Tablets for dogs (Czech Republic, Hungary, Estonia, Latvia, Lithuania, Romania, Slovenia, Slovakia)

*Praziquantel, pyrantel embonate, febantel*

Endogard Plus Flavour Tablets for dogs (United Kingdom, Austria, Belgium, Germany, Denmark, Greece, Ireland, Netherlands, Portugal)

*Praziquantel, pyrantel embonate, febantel*

Endogard Sabor Tablets for dogs (Spain)

*Praziquantel, pyrantel embonate, febantel*

Endogard Flavour Tablets for dogs (Italy)

*Praziquantel, pyrantel embonate, febantel*

Anthelmin Plus Flavour Tablets for dogs (Poland)

*Praziquantel, pyrantel embonate, febantel*

#### 3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each tablet contains:

Praziquantel	50 mg
Pyrantel embonate	144 mg
Febantel	150 mg

Yellow coloured, round, biconvex tablets with visible darker spots, cross-scored on one side.

The tablets can be divided into 2 or 4 equal parts..

#### 4. INDICATION(S)

For the treatment of mixed infestations with the following roundworms and tapeworms in adult dogs and puppies:

Nematodes

Ascarids: *Toxocara canis*, *Toxascaris leonina* (late immature forms and mature forms)

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Cestodes

Tapeworms: *Taenia* spp., *Dipylidium caninum*

## 5. CONTRAINDICATIONS

Do not use simultaneously with piperazine compounds.

Do not exceed the stated dosage when treating pregnant bitches.

Do not use in animals with a known hypersensitivity to the active substance or to any of excipients.

Not for use in dogs younger than 2 weeks of age and/or weighing less than 2 kg.

## 6. ADVERSE REACTIONS

In rare cases transient loose faeces, diarrhoea and/or vomiting may occur in some puppies. In adult dogs, very rare cases of vomiting, with or without diarrhoea, may occur.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Dogs (small and medium size)

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral administration.

### Dosage

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg bodyweight.

Tablets may be halved/quartered to allow accuracy of dosing.

The tablet(s) can be given directly to the dog or disguised in food. No restriction of access to food is required either before or after administration of the product.

## 9. ADVICE ON CORRECT ADMINISTRATION

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible.

Puppies may be wormed with this product from 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 monthly intervals. It is advisable to treat the bitch at the same time as the puppies.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every 2 weeks until weaning.

For routine control a single dose is recommended at 3 monthly intervals. In the event of a heavy roundworm infestation, a repeat dose should be given after 14 days.

## 10. WITHDRAWAL PERIOD

Not applicable.

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

## 12. SPECIAL WARNING(S)

Special warnings for each target species:

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Special precautions for use in animals:

Any part-used tablets should be discarded.

Not for use in dogs younger than 2 weeks and/or weighing less than 2 kg.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In the interests of good hygiene, persons administering the tablet directly to a dog or by adding it to the dog's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

Use during pregnancy, lactation or lay:

Consult a veterinary surgeon before treating pregnant animals for roundworms.

The product may be used during lactation.

Do not use in bitches during the first two-thirds of pregnancy.

Interaction with other medicinal products and other forms of interaction:

Do not use simultaneously with piperazine as the anthelmintic effects of pyrantel and piperazine (used in many worming products for dogs) may be antagonized.  
Concurrent use with other cholinergic compounds can lead to toxicity .

Overdose (symptoms, emergency procedures, antidotes), if necessary:

Benzimidazoles possess wide safety margin. Pyrantel is not absorbed systematically to any extent. Praziquantel also has a wide safety margin of up to five times the recommended dose.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

**15. OTHER INFORMATION**


Print and perforated Alu-Alu blister: 2 tablets (1 blister with 2 tablets), in a box.  
Print and perforated Alu-Alu blister: 4 tablets (2 blisters with 2 tablets), in a box.  
Print and perforated Alu-Alu blister: 10 tablets (1 blister with 10 tablets), in a box.  
Print and perforated Alu-Alu blister: 30 tablets (3 blisters with 10 tablets), in a box.  
Print and perforated Alu-Alu blister: 50 tablets (5 blisters with 10 tablets), in a box.  
Print and perforated Alu-Alu blister: 100 tablets (10 blisters with 10 tablets), in a box.  
Print and perforated Alu-Alu blister: 300 tablets (30 blisters with 10 tablets), in a box.  
Not all pack sizes may be marketed.

\*Valid only for packaging of 100 tablets

1 per 10 kg

**PRACTICE STAMP**

Owner's name: .....  
.....  
Pet's name: .....  
.....  
Dosage:.....  
.....



Approved: 08 July 2020