

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Carton box small and big**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Enrox Flavour 15 mg Tablets for dogs and cats  
Enrofloxacin

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each tablet contains 15 mg of Enrofloxacin.

**3. PHARMACEUTICAL FORM**

Tablets

**4. PACKAGE SIZE**

10 tablets  
100 tablets

**5. TARGET SPECIES**

Dogs and cats.

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
The tablet is given orally once daily or as a divided dose twice daily with or without food.

**8. WITHDRAWAL PERIOD**

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.  
User warnings - please read the package leaflet.

**10. EXPIRY DATE**

EXP:

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of waste material in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

KRKA, d.d., Novo mesto  
Šmarješka cesta 6  
8501 Novo mesto  
Slovenia

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 01656/4007

**17. MANUFACTURER'S BATCH NUMBER**

Lot:

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blisters**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Enrox Flavour 15 mg Tablets for dogs and cats  
Enrofloxacin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

KRKA

**3. EXPIRY DATE**

EXP:

**4. BATCH NUMBER**

Lot:

**5. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

**PACKAGE LEAFLET FOR:**

Enrox Flavour 15 mg Tablets for dogs and cats

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

KRKA, d.d., Novo mesto  
Šmarješka cesta 6  
8501 Novo mesto  
Slovenia

Manufacturer responsible for the batch release:

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia  
Virbac S.A., 1ère avenue, 2065 m L.I.D., 06516 Carros Cedex, France  
KRKA - FARMA d.o.o., V. Holjevca 20/E, 10450 Jastrebarsko, Croatia

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Enrox Flavour 15 mg Tablets for dogs and cats

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each tablet contains 15 mg of Enrofloxacin.  
Round slightly biconvex, cream to light brownish tablets with possible visible white or darker spots and bevel-edged.

**4. INDICATION(S)**

The product is for use in dogs and cats in the treatment of bacterial infections of the alimentary, respiratory and urogenital tracts, skin, secondary wound infections and otitis externa where clinical experience indicates enrofloxacin as the drug of choice.

**5. CONTRAINDICATIONS**

Articular cartilage may be affected during the period of rapid growth, therefore do not use in dogs less than 1 year of age or in exceptionally large breeds of dog with a longer growth period less than 18 months of age.  
Do not use in cats less than 8 weeks of age.  
Do not use in case the dog/cat is allergic to the active substance or to any of the excipients.

Do not use in dogs having seizure disorders, since enrofloxacin may cause stimulation of the central nervous system.  
Do not use for prophylaxis.

## **6. ADVERSE REACTIONS**

During the period of rapid growth, enrofloxacin may affect the development of articular cartilage.

In very rare cases (less than 1 animal in 10,000 animals, including isolated reports) vomiting and anorexia are observed.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dogs and cats.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Do not exceed the recommended dosage. Dog or cat should be given 5 mg/kg of the product orally once daily or as a divided dose twice daily for 5 to 10 days with or without food.

The duration of treatment in dogs may be extended depending on the clinical response and the judgement of the responsible veterinary surgeon.

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing.

The daily dose for cats and small dogs is one tablet per 3 kg body weight.

## **9. ADVICE ON CORRECT ADMINISTRATION**

The tablet should be given orally once daily or as divided dose twice daily with or without food.

## **10. WITHDRAWAL PERIOD**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and blister after "EXP". The expiry date refers to the last day of that month.

## 12. SPECIAL WARNING(S)

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly to other classes of antimicrobials.

Whenever possible, use of fluoroquinolones should be based on susceptibility testing. Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross-resistance.

Official and local antimicrobial policies should be taken into account when the product is used.

Do not use in case of resistance to quinolones, as there exists almost complete cross resistance to other quinolones and complete cross resistance to other fluoroquinolones.

Do not exceed the recommended dosage.

Retinotoxic effects including blindness can occur in cats when the recommended dose is exceeded.

Use the product with caution in dogs with severe renal or hepatic impairment.

As enrofloxacin passes into maternal milk, use only according to the benefit/risk assessment by the responsible veterinarian.

Do not combine with other drugs, such as tetracyclines, phenicols or macrolides because there is a potential that these drugs nullify the desired effect.

Concurrent administration of fluoroquinolones may increase the action of oral anticoagulants (the drugs that prevent the clotting of blood).

Do not combine with theophylline (a drug used in medicine as a bronchial dilator) as this could lead to a prolonged elimination of this substance.

Concurrent administration of magnesium or aluminum containing substances may be followed by retarded absorption of enrofloxacin.

In accidental overdose, vomiting, diarrhoea and central nervous system /behavioural changes may occur.

There is no antidote and treatment should be symptomatic. If necessary, administration of aluminium- or magnesium-containing antacids or activated carbon can be used to reduce absorption of enrofloxacin.

### **User warnings**

Wash hands after use.

In case of contact with the eyes, wash with plenty of clean water.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

People with known allergy to (fluoro)quinolones (a group of antimicrobials) should avoid contact with the product.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

February 2021

**15. OTHER INFORMATION**

Polyamide/Aluminium/Polyvinyl chloride film (OPA/Al/PVC), heat sealed with aluminium foil containing 10 tablets / blister. Each cardboard carton contains 100 tablets in 10 blister packs.

Polyamide/Aluminium/Polyvinyl chloride film (OPA/Al/PVC), heat sealed with aluminium foil containing 10 tablets / blister. Each cardboard carton contains 10 tablets in 1 blister pack.

Not all pack sizes may be marketed.

Approved 22 April 2021

A handwritten signature in black ink, appearing to read "A. Hunter.", is positioned below the approval date.