

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE -

Cardboard box (10 ml, 5 x 10 ml, 10 x 10 ml and 50 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Butador 10 mg/ml solution for injection for horses, dogs and cats

Butorphanol

2. STATEMENT OF ACTIVE SUBSTANCES

Butorphanol (as tartrate) 10 mg/ml

3. PHARMACEUTICAL FORM

Solution for injection

4. PACKAGE SIZE

10 ml

5 x 10 ml

10 x 10 ml

50 ml

5. TARGET SPECIES

Horse, dog, cat

6. INDICATION(S)

-

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Horse IV only / Dog IV, SC, IM / Cat IV, SC

Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Withdrawal period:

Horse: zero days

9. SPECIAL WARNING(S), IF NECESSARY

Accidental injection is dangerous.

10. EXPIRY DATE

EXP {month/year}
Once broached, use within 28 days.

11. SPECIAL STORAGE CONDITIONS

Protect from light.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

VetViva Richter GmbH, 4600 Wels, Austria

16. MARKETING AUTHORISATION NUMBER(S)

Vm 57446/4000

17. MANUFACTURER’S BATCH NUMBER

Batch {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

10 ml and 50 ml clear glass vial type I with bromobutyl rubber stopper and alu-caps

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Butador 10 mg/ml solution for injection

Butorphanol

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Butorphanol (as tartrate) 10 mg/ml

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

10 ml

50 ml

4. ROUTE(S) OF ADMINISTRATION

Horse IV / Dog IV, SC, IM / Cat IV, SC

5. WITHDRAWAL PERIOD(S)

Withdrawal period: zero days

6. BATCH NUMBER

Batch {number}

7. EXPIRY DATE

EXP {month/year}

Once broached use by

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

B. PACKAGE LEAFLET

PACKAGE LEAFLET:

Butador 10 mg/ml solution for injection for horses, dogs and cats (FR, IE, UK)

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

VetViva Richter GmbH, Durisolstrasse 14, 4600 Wels, Austria

Manufacturer responsible for batch release:

Richter Pharma AG, Durisolstrasse 14, 4600 Wels, Austria

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Butador 10 mg/ml solution for injection for horses, dogs and cats

Butorphanol

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

1 ml contains:

Active substance:

Butorphanol	10 mg
(as butorphanol tartrate)	14.58 mg)

Excipient:

Benzethonium chloride	0.1 mg
-----------------------	--------

Clear, colourless to almost colourless solution

4. INDICATIONS

HORSE

As an analgesic

For the short term relief of pain such as colic of gastrointestinal tract origin.

As a sedative and pre-anaesthetic

In combination with α_2 -adrenoceptor agonists (detomidine, romifidine, xylazine):
For therapeutic and diagnostic procedures such as minor standing surgery and sedation of intractable patients.

DOG/CAT

As an analgesic

For relief of moderate visceral pain e.g. pre- and post-surgical as well as post-traumatic pain.

As a sedative

In combination with α_2 -adrenoceptor agonists (medetomidine).

As a pre-anaesthetic

Part of anaesthetic regime (medetomidine, ketamine).

5. CONTRAINDICATIONS

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use for treatment of animals with severe dysfunction of the liver and kidneys, in case of cerebral injury or organic brain lesions and in animals with obstructive respiratory diseases, heart dysfunctions or spastic conditions.

For combination use with α_2 -agonists in horses:

Do not use in horses with a pre-existing cardiac dysrhythmia or bradycardia.

The combination will cause a reduction in gastrointestinal motility and consequently should not be used in cases of colic associated with impaction.

Do not use combination during pregnancy.

6. ADVERSE REACTIONS

HORSE

Undesirable effects are generally related to the known activity of opioids. In published trials with butorphanol, transient ataxia, lasting about 3 to 15 minutes, occurred in about 20 % of horses. Mild sedation occurred in about 10 % of horses. Increased motor activity (running movements) is possible. Gastrointestinal motility may be reduced. This effect is mild and transient.

For combination use:

Any reduction of gastrointestinal motility caused by butorphanol may be enhanced by the concomitant use of α_2 -agonists. The respiratory depressive effects of α_2 -agonists may be enhanced by concomitant butorphanol, particularly if respiratory function is already impaired. Other undesirable effects (e.g. cardiovascular) are likely to be related to the α_2 -agonist.

DOG/CAT

Depression of the respiratory and cardiovascular system. Local pain associated with intramuscular administration. Decreased gastrointestinal motility. In rare cases, ataxia, anorexia and diarrhoea. In cats excitation or sedation, anxiety, disorientation, dysphoria and mydriasis are possible.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

7. TARGET SPECIES

Horse, dog, cat

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Horse: Intravenous use

Dog: Intravenous, subcutaneous and intramuscular use

Cat: Intravenous and subcutaneous use

HORSE

As an analgesic

Monotherapy:

0.1 mg/kg (1 ml/100 kg bw) IV.

As a sedative and as a pre-anaesthetic

With detomidine:

Detomidine: 0.012 mg/kg IV, followed within 5 minutes by

Butorphanol: 0.025 mg/kg (0.25 ml /100 kg bw) IV.

With romifidine:

Romifidine: 0.05 mg/kg IV, followed within 5 minutes by

Butorphanol: 0.02 mg/kg (0.2 ml /100 kg bw) IV.

With xylazine:

Xylazine: 0.5 mg/kg IV, followed after 3 - 5 minutes by

Butorphanol: 0.05 – 0.1 mg/kg (0.5 - 1 ml /100 kg bw) IV.

DOG

As an analgesic

Monotherapy:

0.1 - 0.4 mg/kg (0.01 – 0.04 ml/kg bw) slowly IV (in the lower to medium dose range) as well as IM, SC.

For post-operative pain control the injection should be administered 15 minutes before the end of anaesthesia in order to achieve sufficient pain relief during the recovery phase.

As a sedative

With medetomidine:

Butorphanol: 0.1 mg/kg (0.01 ml/kg bw) IV, IM

Medetomidine: 0.01 mg/kg IV, IM.

As a pre-anaesthetic

With medetomidine and ketamine:

Butorphanol: 0.1 mg/kg (0.01 ml/kg bw) IM
Medetomidine: 0.025 mg/kg IM, followed after 15 minutes by
Ketamine: 5 mg/kg IM.

It is only possible to use atipamezole 0.1 mg/kg body weight for medetomidine-antagonisation when ketamine action has ceased.

CAT

As an analgesic

Monotherapy:

15 minutes prior to recovery
either: 0.4 mg/kg (0.04 ml/kg bw) SC
or: 0.1 mg/kg (0.01 ml/kg bw) IV

As a sedative

With medetomidine:

Butorphanol: 0.4 mg/kg (0.04 ml/kg bw) SC
Medetomidine: 0.05 mg/kg SC.

For wound debridement an additional local anaesthesia is recommended.
Medetomidine-antagonisation is possible with atipamezole 0.125 mg/kg body weight.

As a pre-anaesthetic

With medetomidine and ketamine:

Butorphanol: 0.1 mg/kg (0.01 ml/kg bw) IV
Medetomidine: 0.04 mg/kg IV
Ketamine: 1.5 mg/kg IV.

It is only possible to use atipamezole 0.1 mg/kg body weight for medetomidine-antagonisation when ketamine action has ceased.

The stopper must not be punctured more than 25 times.

9. ADVICE ON CORRECT ADMINISTRATION

Butorphanol is intended for use where short (horse and dog) and short to medium (cat) analgesia is required. The dose may be repeated as required. The need for and timing of repeated treatment will be based on clinical response. For information on the duration of analgesia see section "Other information".

Rapid intravenous injection should be avoided.

Do not mix the product with other veterinary medicinal products in one syringe.

10. WITHDRAWAL PERIOD(S)

Horse

Meat and offal: zero days
Milk: zero hours

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature storage conditions.

Keep the vial in the outer carton in order to protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton after "EXP". The expiry date refers to the last day of that month. Shelf-life after first opening the immediate container: 28 days.

When the container is broached (opened) for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the carton should be discarded should be worked out. This discard date should be written in the space provided.

12. SPECIAL WARNINGS

Special warnings for each target species

The precautionary measures required for contact with animals should be followed and stress factors for the animals should be avoided.

In cats, individual response to butorphanol may be variable. In the absence of an adequate analgesic response, an alternative analgesic agent should be used.

Increasing of the dose may not increase the intensity or duration of analgesia.

Special precautions for use in animals

The safety of the product in puppies, kitten and foals has not been established. Use of the product in these groups should be on the basis of a risk-benefit analysis by the responsible veterinarian.

Due to its antitussive properties, butorphanol may lead to an accumulation of mucous in the respiratory tract. Therefore, in animals with respiratory diseases associated with increased mucous production, butorphanol should only be used after a risk-benefit evaluation by the responsible veterinarian. If respiratory depression occurs, naloxone may be used as an antidote.

Sedation may be noted in treated animals. The combination of butorphanol and α_2 -adrenoceptor agonists should be used with caution in animals with cardiovascular disease. The concurrent use of anticholinergic drugs, e.g atropine should be considered.

Administration of butorphanol and romifidine in one syringe should be avoided due to increased bradycardia, heart block and ataxia.

HORSE

The use of the product at the recommended dose may lead to transient ataxia and/or excitement. Therefore, to prevent injuries in patient and people when treating horses, the location for the treatment should be chosen carefully.

CAT

Cats should be weighed to ensure that the correct dose is calculated. An appropriate graduated syringe must be used to allow accurate administration of the required dose

volume (e.g. insulin syringe or 1 ml graduated syringe). If repeated administrations are required, use different injection sites.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Butorphanol has opioid-like activity. Precautions should be taken to avoid accidental injection/self-injection with this potent drug. The most frequent adverse effects of butorphanol in humans are drowsiness, sweating, nausea, dizziness and vertigo and may occur following unintended self-injection. If accidental self-injection occurs, seek medical advice immediately and show the package leaflet or the label to the physician. Do not drive. An opioid antagonist (e.g. naloxone) may be used as an antidote. Wash splashes from skin and eyes immediately.

Pregnancy and lactation

Butorphanol crosses the placental barrier and penetrates into milk. Studies in laboratory species have not produced any evidence of teratogenic effects. The safety of this veterinary medicinal product has not been established in the target species during pregnancy and lactation. The use of butorphanol during pregnancy and lactation is not recommended.

Interaction with other medicinal products and other forms of interaction

The concomitant administration of other drugs which are metabolised in the liver may enhance the effect of butorphanol.

Butorphanol used with concurrently administered anaesthetics, centrally sedative or respiratory depressive drugs produces additive effects. Any use of butorphanol in this context requires acute control and a careful adaptation of the dose.

Administration of butorphanol may remove the analgesic effect in animals, which have already received pure μ -opioid analgesics.

Overdose (symptoms, emergency procedures, antidotes)

HORSE

Increased dosages could result in respiratory depression as a general opioid effect. Intravenous doses of 1.0 mg/kg (10 x the recommended dose), repeated at 4-hourly intervals for 2 days, led to transient adverse effects, including pyrexia, tachypnoea, CNS signs (hyperexcitability, restlessness, mild ataxia leading to somnolence) and gastrointestinal hypomotility, sometimes with abdominal discomfort. An opioid antagonist (e.g. Naloxone) may be used as an antidote.

DOG/CAT

Miosis (dog)/Mydriasis (cat), respiratory depression, hypotension, disorders of the cardiovascular system and in severe cases respiratory inhibition, shock and coma. Depending on the clinical situation counter-measures should be taken under intense medical monitoring. Monitoring is required for a minimum of 24 hours.

Incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

January 2023

15. OTHER INFORMATION

Pharmacodynamic properties

Butorphanol is a centrally acting analgesic from the group of synthetic opioids with an agonistic-antagonistic effect, agonist at the kappa opioid receptor subtype and antagonist at the mu receptor subtype. The kappa receptors control analgesia, sedation without depression of cardiopulmonary system and body temperature, whereas the mu receptors control supraspinal analgesia, sedation and depression of cardiopulmonary system and body temperature.

The agonist component of butorphanol activity is ten times more potent than the antagonist component.

Analgesia generally occurs within 15 minutes following administration in horse, dog and cat. After a single intravenous dose in the horse analgesia usually lasts up to 2 hours. In the dog it lasts up to 30 minutes after a single intravenous administration. In cats with visceral pain analgesic effects have been demonstrated for up to 6 hours. In cats with somatic pain duration of analgesia has been considerably shorter.

Increased doses do not correlate with increased analgesia, a dosage of about 0.4 mg/kg leads to a ceiling effect.

Butorphanol has minimal cardiopulmonary depressant activity in the target species. It does not cause histamine release in horses. In combination with α_2 -agonists it causes additive and synergistic sedation.

Pharmacokinetic particulars

Post parenteral administration absorption of the product is rapid and almost complete with serum peak levels occurring after 0.5 - 1.5 hours. It is highly bound to plasma proteins (up to 80 %). Metabolism is rapid and mainly occurs in the liver.

Two inactive metabolites are produced. The elimination occurs mainly through urine (to a major extent) and faeces.

HORSE: Volume of distribution is large after IV administration (2.1 l/kg) suggesting wide distribution into tissues. Terminal half life is short: about 44 minutes. 97 % of the dose after IV administration in the horse will be eliminated in less than 5 hours.

DOG: Volume of distribution is large after IV administration (4.4 l/kg) suggesting wide distribution into tissues. Terminal half life is short: about 1.7 hours.

CAT: Volume of distribution is large after IV administration (7.4 l/kg) suggesting wide distribution into tissues. Terminal half life is short: about 4.1 hours.

Package sizes

1 x 10 ml, 5 x 10 ml, 10 x 10 ml, 1 x 50 ml.

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.