

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE {0.5, 1, 2.5 & 5 L  
cardboard box}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Cydectin 0.5% w/v Pour-On

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Active Substance:

Moxidectin 5.00 mg

Excipients:

Butylated hydroxyanisole E320 0.10 mg

Tert butyl hydroquinone 0.03 mg

**3. PACKAGE SIZE**

0.5 L

1 L

2.5 L

5 L

**4. TARGET SPECIES**

Cattle

**5. INDICATION(S)**

**6. ROUTES OF ADMINISTRATION**

For pour-on use.

**7. WITHDRAWAL PERIODS**

Meat and offal: 14 days.

Milk: 6 days (144 hours).

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once opened, use within 6 months.

Once opened, use by: ".../.../..."

## **9. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.  
Protect from light.  
If accidentally frozen, shake vigorously before use.

## **10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

## **11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

## **13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited

## **14. MARKETING AUTHORISATION NUMBERS**

Vm 42058/5131

## **15. BATCH NUMBER**

Lot {number}

## **16. SPECIAL WARNING(S), IF NECESSARY**

Environmental risks have been identified for this product and special precautions apply. Read the package leaflet before use.

## **17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

## **18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-VPS

Veterinary medicinal product subject to prescription.

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {0.5, 1, 2.5 & 5 L  
label}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Cydectin 0.5% w/v Pour-On

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Active Substance:

Moxidectin 5.00 mg

Excipients:

Butylated hydroxyanisole E320 0.10 mg

Tert butyl hydroquinone 0.03 mg

0.5 L

1 L

2.5 L

5 L

**3. TARGET SPECIES**

Cattle

**4. ROUTES OF ADMINISTRATION**

For pour-on use.

Read the package leaflet before use.

**5. WITHDRAWAL PERIODS**

Meat and offal: 14 days.

Milk: 6 days (144 hours).

**6. EXPIRY DATE**

Exp. {mm/yyyy}

Once opened, use within 6 months.

Once opened, use by: ".../.../..."

**7. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Protect from light.

If accidentally frozen, shake vigorously before use.

## **8. NAME OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited

## **9. BATCH NUMBER**

Lot {number}

## **10. SPECIAL WARNING(S), IF NECESSARY**

Environmental risks have been identified for this product and special precautions apply. Read the package leaflet before use.

## **11. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

## **12. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

For animal treatment only.

POM-VPS

Veterinary medicinal product subject to prescription.

**PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Cydectin 0.5% w/v Pour-On for Cattle

**2. COMPOSITION**

Each ml contains:

**Active Substance:**

Moxidectin 5.00 mg

**Excipients:**

Butylated hydroxyanisole E320 0.10 mg

Tert butyl hydroquinone 0.03 mg

Pale yellow oily solution.

**3. TARGET SPECIES**

Cattle.

**4. INDICATIONS FOR USE**

Infections of cattle with parasites sensitive to moxidectin.  
For the treatment of infections caused by:

- Adult and larval gastro-intestinal nematodes:
  - *Haemonchus placei*
  - *Ostertagia ostertagi* (including inhibited larvae)
  - *Trichostrongylus axei*
  - *Nematodirus helvetianus*
  - *Cooperia oncophora*
  - *Cooperia punctata* (adults)
  - *Oesophagostomum radiatum* (adults)
  - *Bunostomum phlebotomum* (adults)
  
- Adult respiratory tract nematode
  - *Dictyocaulus viviparus*
  
- Warbles (migrating larvae)
  - *Hypoderma bovis*
  - *Hypoderma lineatum*
  
- Lice
  - *Linognathus vituli*
  - *Haematopinus eurysternus*

- *Solenopotes capillatus*
- *Bovicola bovis* (*Damalinia bovis*)
  
- Mange Mites
  - *Sarcoptes scabiei*
  - *Psoroptes ovis*
  - *Chorioptes bovis*
  
- Horn Flies
  - *Haematobia irritans*
  
- Cydectin 0.5% w/v Pour-On for Cattle has a persistent effect in preventing against reinfection by:
  - *Ostertagia ostertagi* for 5 weeks
  - *Dictyocaulus viviparus* for 6 weeks

## 5. CONTRAINDICATIONS

Not to be used in other species as severe adverse reactions, including fatalities in dogs, may occur.

## 6. SPECIAL WARNINGS

Special precautions for safe use in the target species:  
For topical application only.

To avoid secondary reactions due to the death of *Hypoderma* larvae in the oesophagus or the spine, it is recommended to administer the veterinary medicinal product at the end of the period of fly activity and before the larvae reach their resting sites: consult the veterinarian to know the correct treatment period.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- Do not smoke, eat or drink while handling the veterinary medicinal product.
- Avoid direct contact with skin and eyes.
- Wash hands after use.
- Protective clothes and gloves are recommended when using the veterinary medicinal product.
- If splashed in the eye or on the skin, wash with plenty of clean, running water immediately.

Special precautions for the protection of the environment

Moxidectin fulfils the criteria for a (very) persistent, bioaccumulative and toxic (PBT) substance; therefore, exposure of the environment to moxidectin must be limited to the extent possible. Treatments should be administered only when necessary and

should be based on faecal egg counts or evaluation of the risk of infestation at the animal and/or herd level.

Like other macrocyclic lactones, moxidectin has the potential to adversely affect non-target organisms, in particular aquatic organisms and dung fauna.

- Faeces containing moxidectin excreted onto pasture by treated animals may temporarily reduce the abundance of dung feeding organisms. Following treatment of cattle with the veterinary medicinal product, levels of moxidectin that are potentially toxic to dung fly species may be excreted over a period of more than 2 weeks and may decrease dung fly abundance during that period. It has been established in laboratory tests that moxidectin may temporarily affect dung beetle reproduction; however, field studies indicate no long-term effects. Nevertheless, in case of repeated treatments with moxidectin (as with veterinary medicinal products of the same anthelmintic class) it is advisable not to treat animals every time on the same pasture to allow dung fauna populations to recover.
- Moxidectin is inherently toxic to aquatic organisms including fish. This implies that when allowing moxidectin to enter water bodies, this may have a severe and lasting impact on aquatic life. To mitigate this risk, the veterinary medicinal product should be used only according to the label instructions. Based on the excretion profile of moxidectin when administered as the pour-on formulation, treated animals should not have access to watercourses during the first week after treatment.

Pregnancy, lactation and fertility:

Moxidectin has been shown to be safe for use in pregnant and lactating animals and breeding bulls.

Major incompatibilities:

Not to be mixed with other veterinary medicinal products before administration.

## 7. ADVERSE EVENTS

Cattle

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site reaction Neurological signs (such as ataxia (incoordination) and trembling) Lethargy
---	--

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

## **8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

The veterinary medicinal product should be administered at a dose rate of 500 µg moxidectin/kg body weight (1 ml for 10 kg) as a single topical application.

To be administered along the midline of the back of the animal from the withers to the tailhead.

Apply to clean healthy skin.

## **9. ADVICE ON CORRECT ADMINISTRATION**

None.

## **10. WITHDRAWAL PERIODS**

Meat and offal: 14 days.

Milk: 6 days (144 hours).

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not store above 25 °C

Protect from light

If accidentally frozen, shake vigorously before use.

Do not use after the expiry date stated on the label after "Exp,"

Shelf-life after first opening the container: 6 months.

When the container is opened for the first time, using the in-use shelf-life which is specified on this package insert, the date on which any veterinary medicinal product remaining in the container should be discarded should be worked out. This discard date should be written in the space provided on the label.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste material derived from such veterinary medicinal products should be disposed of in accordance with local requirements. The veterinary medicinal product should not enter water courses as moxidectin may be dangerous for fish and other aquatic organisms.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.



## 14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 42058/5131

The product is packaged in 0.5 L, 1 L, 2.5 L and 5 L HDPE containers.

Not all pack sizes may be marketed.

## 15. **PID LINK (Do not print heading)**

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

## 16. CONTACT DETAILS

### **Marketing authorisation holder and contact details to report suspected adverse reactions:**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP  
Tel: +44 0345 300 8034

### **Manufacturer for the batch release:**

Zoetis Manufacturing & Research Spain, S.L.  
Ctra. Camprodón s/n "la Riba"  
17813 Vall de Bianya  
Girona  
SPAIN

## 17. OTHER INFORMATION

*Gavin Hall*

Approved: 06 July 2024