

## 1. NAME OF VETERINARY MEDICINAL PRODUCT

Soloxine 0.5 mg Tablet

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

### Each tablet contains:

#### Active substance

Levothyroxine Sodium                      0.50 mg

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Small elliptical white tablets.

Scored on the face of each tablet, strength in milligrams to the right and the word SOLOXINE on the reverse.

## 4. CLINICAL PARTICULARS

### 4.1 Target species

Dogs.

### 4.2 Indications for use, specifying the target species

For the long term treatment of thyroid insufficiency in dogs.

### 4.3 Contraindications

Do not use in animals with known hypersensitivity to the active ingredient.

Do not use in animals suffering from thyrotoxicosis or uncorrected adrenal insufficiency.

### 4.4 Special warnings for each target species

None.

### 4.5 Special precautions for use

#### i) Special precautions for use in animals

Appropriate laboratory tests should be conducted to confirm the diagnosis and ensure correct dosage.

Caution should be exercised in the treatment of dogs with clinically significant cardiac disease, hypertension or any disease rendering the animal susceptible to sharply increased metabolic rate. In such cases, consideration should be given to reducing the starting dose, increasing the dose at intervals whilst monitoring all clinical signs. Dogs with concurrent hypoadrenocorticism should be stabilised with appropriate steroid therapy before commencing treatment with levothyroxine sodium.

The effects of thyroxine therapy are slow in being manifested.

**ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals**

In case of accidental ingestion, seek medical advice immediately and show the doctor the label.

Wash hands after use.

**iii) Other precautions**

None.

**4.6 Adverse reactions (frequency and seriousness)**

When administered at an appropriate dose there should not be any adverse effects associated with therapy. Thyrotoxicosis is unusual, but may develop in dogs receiving high doses or in those with impaired metabolism (i.e. renal or hepatic insufficiency).

Clinical signs include panting, nervousness, tachycardia, aggressive behaviour, polyuria, polydipsia, polyphagia and weight loss.

**4.7 Use during pregnancy, lactation or lay**

Can be used during pregnancy and lactation.

The safety of the product has not been tested in special reproduction studies. However, levothyroxine sodium is an endogenous hormone and thyroid hormones are essential for the developing foetus. Hypothyroidism during pregnancy may result in poor foetal and perinatal outcomes. Therefore, hypothyroid bitches intended to be bred should be monitored on a regular basis before, during and after pregnancy as the dose of levothyroxine sodium may need to be adjusted.

**4.8 Interaction with other medicinal products and other forms of interaction**

In diabetic dogs with concurrent hypothyroidism, careful monitoring of diabetic control is recommended once thyroid hormone treatment commences. Dosages of insulin may need to be increased due to thyroid hormone enhancement of glucose absorption, glycogenolysis and gluconeogenesis.

**4.9 Amounts to be administered and administration route**

The commonly prescribed starting dose is 22 µg/kg bodyweight/day. However, due to individual differences in absorption and metabolism this is frequently too low and doses of up to 44 µg/kg bodyweight/day may be required. The dosage should be adjusted for the individual case based on the clinical response of the patient and laboratory investigations and should be administered daily.

For oral administration.

**4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Chronic overdosage will eventually lead to thyrotoxicosis, manifested by: panting, nervousness, tachycardia, aggressive behaviour, polyuria, polydipsia, polyphagia and weight loss.

**4.11 Withdrawal periods**

Not applicable.

**5. PHARMACOLOGICAL PARTICULARS**

Pharmacotherapeutic group: Thyroid Hormones  
ATC Vet Code: QH03AA01

### **5.1 Pharmacodynamic properties**

Levothyroxine is a synthetic homologue of the naturally occurring thyroid hormone, Thyroxine (T<sub>4</sub>).

Levothyroxine is converted to the more biologically active triiodothyronine (T<sub>3</sub>). T<sub>3</sub> binds via specific receptors within the plasma membrane, mitochondria and chromatin resulting in changes in DNA transcription and protein synthesis. Onset of action is therefore slow. Thyroid hormones may act on cellular processes with effects on the basal metabolic rate, cardiac function and blood flow, lipid and carbohydrate metabolism. They are essential for the normal growth and development of the neurological and skeletal systems.

### **5.2 Pharmacokinetic properties**

Time to reach peak serum concentration takes between 2 and 5 hours and the half life of levothyroxine sodium in dogs following oral administration varies from approximately 6 to 20 hours.

Pharmacokinetic properties, particularly absorption and rate of metabolism, vary markedly between individual dogs, with variations in maximum serum concentration of up to 3 times. Therefore it is important to tailor the dose to the individual dog by regular laboratory and clinical monitoring following commencement of treatment.

Pharmacokinetic trials showed that once daily dosing gave higher peak concentrations than dividing the same dose and giving twice daily.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose Monohydrate  
Microcrystalline Cellulose  
Maize Starch Pregelatinised  
Magnesium Stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

### **6.4 Special precautions for storage**

Do not store above 25° C.  
Protect from light.

### **6.5 Nature and contents of immediate packaging**

High-density, brown, polyethylene bottles containing 250 tablets, hermetically sealed and closed with childproof screw cap.

**6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products, if appropriate**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**7.0 MARKETING AUTHORISATION HOLDER**

Virbac S.A.,  
Virbac 1,  
1ère Avenue - 2065 M - LID,  
BP 27, 06511 Carros Cedex,  
France

**8.0 MARKETING AUTHORISATION NUMBERS**

UK: Vm 05653/4139  
Eire: VPA 10988/69/4

**9.0 DATE OF RENEWAL OF THE AUTHORISATION**

UK: 8<sup>th</sup> January 2007  
Eire: 26<sup>th</sup> October 2008

**10.0 DATE OF ANY REVISION OF THE TEXT**

02 July 2010