

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Methoxasol 20/100 mg/ml solution for use in drinking water for pigs and chickens

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each ml contains:

Active substances:

Trimethoprim	20.0 mg
Sulfamethoxazole	100.0 mg

Excipients:

N-methyl pyrrolidone	690.8 mg
----------------------	----------

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for use in drinking water.  
Clear and-yellow solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Pigs and chickens (broilers).

#### **4.2 Indications for use, specifying the target species**

Pigs: Treatment and metaphylaxis of respiratory infections caused by *Actinobacillus pleuropneumoniae* susceptible to trimethoprim and sulfamethoxazole where the disease has been diagnosed in the herd.

Broilers: Treatment and metaphylaxis of respiratory infections caused by *Escherichia coli* susceptible to trimethoprim and sulfamethoxazole where the disease has been diagnosed in the flock.

#### **4.3 Contraindications**

Do not use in animals suffering from severe liver or kidney disease, oliguria or anuria.  
Do not use in animals with impaired haematopoietic systems.  
Do not use in cases of hypersensitivity to sulphonamides or trimethoprim or any of the excipients.

#### 4.4 Special warnings for each target species

Severely diseased animals can have a decreased appetite and water consumption. If necessary the concentration of the veterinary medicinal product in the drinking water may need to be adjusted to make sure that the recommended dosage is being consumed. However if the concentration of the veterinary medicinal product is increased too much, the intake of the medicated drinking water decreases for palatability reasons. Therefore water intake should be monitored regularly, especially in broilers.

In case of insufficient intake of water, pigs should be treated parenterally.

#### 4.5 Special precautions for use

##### Special precautions for use in animals

Due to the likely variability (time, geographical) in susceptibility of bacteria for potentiated sulphonamides, occurrence of resistance of bacteria may differ from country to country and even from farm to farm, and therefore bacteriological sampling and susceptibility testing are recommended. Use of the veterinary medicinal product should be based on culture and sensitivity of micro-organisms from diseased cases on farm or from recent previous experience on the farm. Use of the veterinary medicinal product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to sulfamethoxazole and trimethoprim and may also decrease the effectiveness of combinations of trimethoprim with other sulphonamides due to the potential for cross resistance. Official and local antimicrobial policies should be taken into account when the veterinary medicinal product is used.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

This veterinary medicinal product contains sulfamethoxazole, which can cause allergic reactions in some people.

People with known hypersensitivity (allergy) to sulphonamides or trimethoprim should avoid contact with the veterinary medicinal product.

During preparation and administration of the medicated drinking water, skin contact with the drug should be avoided. Therefore it is recommended to wear impermeable gloves e.g. rubber or latex when applying the veterinary medicinal product. Do not handle this veterinary medicinal product if you know you are allergic to trimethoprim or sulphonamides.

Do not smoke, drink or eat when handling the veterinary medicinal product.

If you develop symptoms following exposure to the veterinary medicinal product such as skin rash, you should seek medical advice and show the present warning to the doctor. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

This veterinary medicinal product can cause eye-irritation.

Always wear protective glasses when mixing the veterinary medicinal product with drinking water.

In the event of eye contact, rinse the eye with copious amounts of clean water and if irritation occurs, seek medical attention. In the event of accidental ingestion, seek medical advice. Wash hands and contaminated skin immediately after handling the veterinary medicinal product.

Laboratory studies in rabbits and rats conducted with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. The veterinary medicinal product should not be administered by pregnant women and women suspected of being pregnant. Personal protective equipment consisting of gloves should be worn when handling the veterinary medicinal product by women of childbearing age.

Special precautions for the protection of the environment

Not applicable.

Other precautions

Not applicable.

#### 4.6 Adverse reactions (frequency and seriousness)

Pigs

Rare (1 to 10 animals / 10,000 animals treated):	Hypersensitivity.
---	-------------------

Chickens

Rare (1 to 10 animals / 10,000 animals treated):	Hypersensitivity Decreased drinking
---	--

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

#### 4.7 Use during pregnancy, lactation or lay

The safety of the veterinary medicinal has not been established in pigs and chickens during pregnancy, lactation, lay or in animals intended for breeding. Laboratory studies in rats conducted with trimethoprim have shown evidence of teratogenicity at higher doses than the recommended therapeutic ones. Laboratory studies in rabbits and rats conducted with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. The use of the veterinary medicinal product is not recommended during pregnancy and lactation.

#### 4.8 Interaction with other medicinal products and other forms of interaction

Do not combine with other veterinary medicinal products.

#### 4.9 Amount(s) to be administered and administration route

Route of administration: in drinking water use.

The veterinary medicinal product can be added directly to the drinking water to prepare a therapeutic solution at the calculated concentration, but can also be used in a concentrated stock solution by adding 200 ml of the veterinary medicinal product

per litre of water and diluting this further if required. Do not use this veterinary medicinal product undiluted or in higher concentrated stock solutions.

Pigs: 25 mg TMPS per kg bodyweight per day for 3-4 days, corresponding to 1 ml of the veterinary medicinal product per 4,8 kg bodyweight per day. This corresponds to approximately 1 litre of the veterinary medicinal product in 500 L drinking water. Based on the recommended dose, daily water consumption, and the number and weight of the pigs to be treated, the exact daily concentration of the veterinary medicinal product should be calculated according to the following formula:

$$\frac{\text{Mean body weight (kg) of pigs to be treated}}{\text{Mean daily water consumption (l) per pig} \times 4.8} = \text{xx ml veterinary medicinal product per l drinking water}$$

Broilers: 33 mg TMPS per kg bodyweight per day for 3-4 days, corresponding to 1 ml of the veterinary medicinal product per 3,64 kg bodyweight per day. This corresponds to approximately 1 litre of the veterinary medicinal product in 750 L drinking water. Based on the recommended dose, daily water consumption, and the number and weight of the birds to be treated, the exact daily concentration of the veterinary medicinal product should be calculated according to the following formula:

$$\frac{\text{Mean body weight (kg) of broilers to be treated}}{\text{Mean daily water consumption (l) per bird} \times 3.64} = \text{xx ml veterinary medicinal product per l drinking water}$$

One litre of the veterinary medicinal product weighs 1079 gram; therefore weight can also be used to measure the veterinary medicinal product quantity to be added in drinking water, according to the following formula:

$$\text{Amount to be added in drinking water (g/L)} = \text{calculated ml/L} \times 1.079.$$

The daily amount is to be added to the drinking water such that all medication will be consumed in 24 hours. Medicated drinking water and stock solutions should be freshly prepared every 24 hours. During the treatment period animals should not have access to other water sources than the medicated water the dilution of which should be calculated to ensure that animals always have sufficient water available. To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing. The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage the concentration of the veterinary medicinal product has to be adjusted accordingly.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In pigs, a 2 ½ fold overdose induces no adverse reactions.

In chickens an acute overdose will likely not occur because the birds will be reluctant to drink the strongly concentrated drinking water (too bitter taste if above 2 litres of the veterinary medicinal product per 1000 litres drinking water). Chronic overdose in

chickens will result in a strongly diminished water- and feed intake and retarded growth.

#### 4.11 Withdrawal period(s)

Meat and offal:

Pigs: 5 days.

Broilers: 6 days.

Eggs:

Not for use in birds producing or intending to produce eggs for human consumption.

## 5. PHARMACOLOGICAL PROPERTIES

**Pharmacotherapeutic group:** Antibacterials for systemic use - Sulfamethoxazole and trimethoprim

**ATCvet code:** QJ01EW11

### 5.1 Pharmacodynamic properties

*In vitro* trimethoprim is generally bacteriostatic and has a broad spectrum of activity against both gram-positive and gram-negative bacteria. A synergistic and bactericidal effect occurs when trimethoprim is combined with sulfamethoxazole, because trimethoprim and sulfamethoxazole inhibit sequential steps in the synthesis of tetrahydrofolic acid, an essential metabolic cofactor in bacterial synthesis of purine and, subsequently, DNA.

### 5.2 Pharmacokinetic particulars

Following oral administration both active ingredients are rapidly absorbed from the gut. The  $C_{max}$  of sulfamethoxazole in pigs is approximately 6.2 µg/g. The  $C_{max}$  of trimethoprim is 0.29 µg/g. The  $C_{max}$  of sulfamethoxazole in chickens is approximately 9.0 µg/g, whereas that of trimethoprim is 0.12 µg/g.

High trimethoprim concentrations are found in the kidneys, the liver and the lungs. With the exception of the kidneys, sulfamethoxazole concentrations in the tissues are significantly lower than in the plasma. Protein binding for trimethoprim and sulfamethoxazole is not very high.

The drugs are primarily excreted through the kidneys (both actively and passively), but elimination also occurs through the faeces. Elimination is relatively fast both in poultry and pigs. Plasma elimination half-life for trimethoprim in poultry is less than 1 hour and that of sulfamethoxazole, approximately 1.5 hours. In pigs, elimination half-life for both substances is approximately 2.5 hours. Within 48 hours after the last medication trimethoprim, sulfamethoxazole and their metabolites are undetectable in urine and faeces.

### 5.3 Environmental properties

Trimethoprim is persistent in soils.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

N-Methylpyrrolidone  
Propylene glycol  
Sodium hydroxide, (for pH adjustment)  
Water, purified

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening the immediate packaging: 1 year.  
Shelf life after dilution or reconstitution according to directions: 24 hours.

### **6.4 Special precautions for storage**

Do not freeze.

### **6.5 Nature and composition of immediate packaging**

1000 ml HDPE bottle closed with a tamper proof HDPE screw cap.  
5000 ml HDPE can closed with tamper proof HDPE screw cap.  
Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Eurovet Animal Health B.V.  
Handelsweg 25  
5531 AE Bladel  
The Netherlands

## **8. MARKETING AUTHORISATION NUMBER**

Vm 16849/5007

**9. DATE OF FIRST AUTHORISATION**

04 September 2012

**10. DATE OF REVISION OF THE TEXT**

December 2023

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

**11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT**

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk).

A handwritten signature in black ink, consisting of several vertical strokes followed by a horizontal line and a long, sweeping flourish that extends to the right.

Approved 15 December 2023