



## **9. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Insecticidal Collar for cats weighing 14 g and 35 cm in length.

Remove collar from sachet and fasten around the cat's neck adjusting until a comfortable fit is achieved. In the fully grown cat, correct fitting will allow one finger to be inserted between the collar and the neck. Any excess length of collar extending beyond the buckle should be cut off and disposed of with the domestic refuse. Replace every 4 months if flea reinfestation is a risk.

## **10. ADVICE ON CORRECT ADMINISTRATION**

Not Applicable.

## **11. WITHDRAWAL PERIOD**

Not Applicable.

## **12. SPECIAL WARNING(S), IF NECESSARY**

Special warnings for each target species:

Do not allow animals to chew the collar.

If signs of disease persist or appear, consult your veterinary surgeon.

Fleas from pets often infest animals basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated with a suitable insecticide and vacuumed regularly. Remove fleas from young kittens and nursing queens by careful use of a flea comb.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not remove collar from the pack until ready to use.

Do not allow children to play with the collar.

Do not allow children to suck or chew the collar.

Do not smoke, eat or drink while handling collar.

Do not sleep with pets wearing flea collars.

Dimpylate / Diazinon is an organophosphorus compound: DO NOT handle if under medical advice not to work with ANTICHOLINESTERASES.

Avoid excessive handling of the collar.

Cut off any excess collar with scissors and dispose of safely.

If you feel unwell after handling the collar, seek medical advice.

Wash hands thoroughly after handling the collar.

Pregnancy and lactation:

Not for use on nursing or lactating queens.

**13. EXPIRY DATE**

<EXP {month/year}>

**14. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Protect from direct sunlight.

Do not remove collar from the pack until ready to use.

**15. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary

medicinal product should be disposed of in accordance with local requirements.

EXTREMELY DANGEROUS TO FISH AND AQUATIC LIFE. Do not contaminate ponds, waterways or ditches with the collar or empty packaging.

**16. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For external use only.

For animal treatment only.

AVM-GSL

UK authorised veterinary medicinal product.

**17. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**18. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder:

VIRBAC SA

1ère Avenue - 2065 m - L.I.D.

06516 CARROS

France

**19. MARKETING AUTHORISATION NUMBER**

Vm 05653/4093

**20. MANUFACTURER'S BATCH NUMBER**

<Batch> {number}

**21. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

March 2016

**22. OTHER INFORMATION**

Distributed by :

Johnson's Veterinary Products Ltd

Sutton Coldfield

West Midlands

B75 7DF

United Kingdom.

Tel: 0121-378-1684

Email: [info@johnsons-vet.com](mailto:info@johnsons-vet.com)

**PACKAGE LEAFLET FOR:**

JOHNSON'S FLEA GUARD COLLAR FOR CATS

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Virbac S.A.

1ère avenue - 2065m – L.I.D.

06516 Carros

France

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

JOHNSON'S FLEA GUARD COLLAR FOR CATS

**3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS**

**Active substance:**

Diazinon 2.1g (15% W/W) per collar

**4. INDICATION(S)**

For the control of fleas on cats for up to 4 months.

**5. CONTRAINDICATIONS**

Do not use on kittens under 6 months of age.

**6. ADVERSE REACTIONS**

A slight irritation on the neck may occasionally occur.

Remove collar immediately should any signs of skin irritation or adverse reaction occur.

REMOVE COLLAR FROM CAT IMMEDIATELY SHOULD ANY SIGNS OF DROWSINESS, LISTLESSNESS OR RESPIRATORY DIFFICULTIES DEVELOP SOON AFTER FIXING OF COLLAR

**7. TARGET SPECIES**

Cats

**8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Insecticidal Collar for cats weighing 14 g and 35 cm in length.

Remove collar from sachet and fasten around the cat's neck adjusting until a comfortable fit is achieved. In the fully grown cat, correct fitting will allow one finger to be inserted between the collar and the neck. Any excess length of collar extending beyond the buckle should be cut off and disposed of with the domestic refuse. Replace every 4 months if flea reinfestation is a risk.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Not Applicable.

## **10. WITHDRAWAL PERIOD(S)**

Not Applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Protect from direct sunlight.

Do not remove collar from the pack until ready to use.

## **12. SPECIAL WARNING(S)**

### **Special warnings for each target species**

Do not allow animals to chew the collar.

If signs of disease persist or appear, consult your veterinary surgeon.

Fleas from pets often infest animals basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated with a suitable insecticide and vacuumed regularly. Remove fleas from young kittens and nursing queens by careful use of a flea comb.

### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Do not remove collar from the pack until ready to use.

Do not allow children to play with the collar.

Do not allow children to suck or chew the collar.

Do not smoke, eat or drink while handling collar.

Do not sleep with pets wearing flea collars.

Dimpylate / Diazinon is an organophosphorus compound: DO NOT handle if under medical advice not to work with ANTICHOLINESTERASES.

Avoid excessive handling of the collar.

Cut off any excess collar with scissors and dispose of safely.

If you feel unwell after handling the collar, seek medical advice.

Wash hands thoroughly after handling the collar.

**Use during pregnancy, lactation**

Not for use on nursing or lactating queens.

**Interaction with other medicinal products and other forms of interaction**

Do not use any other insecticides while the collar is being worn or within 7 days of removal.

For Animal Treatment Only.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. EXTREMELY DANGEROUS TO FISH AND AQUATIC LIFE. Do not contaminate ponds, waterways or ditches with the collar or empty packaging.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

**15. OTHER INFORMATION**

AVM-GSL

Vm 05653/4093

**Approved: 17/08/2017**

