

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

Cardboard box containing 1 bottle

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Zelys 1.25 mg chewable tablets

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains:

Pimobendan            1.25 mg

**3. PACKAGE SIZE**

60 tablets

**4. TARGET SPECIES**

Dogs

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Oral use.

**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once opened use within 2 months.

Once opened, use by:...

**9. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Keep the bottle tightly closed in order to protect from moisture.

Any unused tablet portion should be returned to the bottle and be used for the next administration.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Ceva Sante Animale

**14. MARKETING AUTHORISATION NUMBERS**

Vm 14966/5085

**15. BATCH NUMBER**

Lot {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

Bottle

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Zelys 1.25 mg chewable tablets

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains:

Pimobendan            1.25 mg

**3. TARGET SPECIES**

Dogs

**4. ROUTES OF ADMINISTRATION**

Oral use. Read the package leaflet before use.

**5. WITHDRAWAL PERIODS**

**6. EXPIRY DATE**

Exp. {mm/yyyy}

Once opened use within 2 months.

**7. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Keep the bottle tightly closed in order to protect from moisture.

Any unused tablet portion should be returned to the bottle and be used for the next administration.

**8. NAME OF THE MARKETING AUTHORISATION HOLDER**

Ceva Sante Animale

**9. BATCH NUMBER**

Lot {number}

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

Cardboard box containing 36 or 96 tablets

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Zelys 1.25 mg chewable tablets

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains:

Pimobendan 1.25 mg

**3. PACKAGE SIZE**

36 tablets

96 tablets

**4. TARGET SPECIES**

Dogs

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Oral use.

**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

Exp. {mm/yyyy}

**9. SPECIAL STORAGE PRECAUTIONS**

Do not store above 30 °C.

Any unused tablet portion should be returned to the blister and be used for the next administration.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Ceva Sante Animale

**14. MARKETING AUTHORISATION NUMBERS**

Vm 14966/5085

**15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING**

Blister

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Zelys



**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

Pimobendan                      1.25 mg

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

## **PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

### **PACKAGE LEAFLET**

#### **1. Name of the veterinary medicinal product**

Zelys 1.25 mg chewable tablets for dogs

#### **2. Composition**

Each tablet contains:

##### **Active substance**

Pimobendan 1.25 mg  
Round in shape beige to light brown tablet, with single score line on one side.  
The tablets can be divided into two equal parts.

#### **3. Target species**

Dogs

#### **4. Indications for use**

For the treatment of canine congestive heart failure originating from valvular insufficiency (mitral and/or tricuspid valve regurgitation) or dilated cardiomyopathy. (See also section "Administration routes and dosage").

#### **5. Contraindications**

Do not use pimobendan in hypertrophic cardiomyopathies or in diseases in which an improvement in cardiac output cannot be achieved for functional or anatomical reasons (e.g. aortic stenosis).

Do not use in dogs with severe impairment of liver function since pimobendan is metabolised mainly via the liver.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients. (See also section "Pregnancy and lactation").

#### **6. Special warnings**

##### Special warnings:

None.

##### Special precautions for safe use in the target species:

The chewable tablets are flavoured. In order to avoid any accidental ingestion, store tablets out of reach of the animals.

##### *For veterinarians only*

The blood glucose should be tested regularly during treatment in dogs with existing diabetes mellitus.

Monitoring of cardiac function and morphology is recommended in animals treated with pimobendan (See also section "Adverse events")

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental ingestion, especially by a child, may lead to the occurrence of tachycardia, orthostatic hypotension, flushing of the face and headaches.

Unused part-tablets should be returned to the open blister space, or to the bottle and inserted back into the outer packaging. Keep in a safe place out of the sight and reach of children.

Close bottle tightly with cap directly after removal of the required number of tablets or part-tablets.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

Pregnancy and lactation:

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic or foetotoxic effects. Laboratory studies in rats and rabbits have shown evidence of maternotoxic and embryotoxic effects at high doses. Pimobendan is excreted into milk. The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Use only according to the benefit/risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

In pharmacological studies no interaction between the cardiac glycosides strophanthin and pimobendan was observed. The pimobendan-induced increase in cardiac contractility is attenuated by the calcium antagonists verapamil and diltiazem and by the  $\beta$ -antagonist propranolol.

Overdose:

In case of overdose, please contact your veterinarian.

*For veterinarians only*

In the case of overdose, a positive chronotropic effect, vomiting, apathy, ataxia, heart murmurs or hypotension may occur. In this situation, the dosage should be reduced and appropriate symptomatic treatment should be initiated.

In prolonged exposure (6 months) of healthy beagle dogs at 3 and 5 times the recommended dose, mitral valve thickening and left ventricular hypertrophy were observed in some dogs. These changes are of pharmacodynamic origin.

## **7. Adverse events**

### Dogs

Rare(1 to 10 animals / 10,000 animals treated):

Vomiting<sup>1</sup>, Diarrhoea<sup>2</sup>,  
Anorexia<sup>2</sup>, Lethargy<sup>2</sup>,  
Increased heart rate<sup>1</sup>, Heart valve disorder<sup>3</sup>

Very rare(<1 animal / 10,000 animals treated, including isolated reports):

Mucosa petechiae<sup>4,5</sup>, Haemorrhage<sup>4,6</sup>

<sup>1</sup> Dose-dependent and can be avoided by reducing the dose.

<sup>2</sup> Transient

<sup>3</sup> An increase in mitral valve regurgitation has been observed during chronic pimobendan treatment in dogs with mitral valve disease.

<sup>4</sup> Although a relationship with pimobendan has not been clearly established, these signs of effects on primary haemostasis disappear when the treatment is withdrawn.

<sup>5</sup> Pinpoint discolouration of mucosae due to bleeding

<sup>6</sup> Subcutaneous

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: [adverse.events@vmd.gov.uk](mailto:adverse.events@vmd.gov.uk)

## **8. Dosage for each species, routes and method of administration**

Oral use.

Do not exceed the recommended dosage.

To ensure a correct dosage, body weight should be determined as accurately as possible. The tablets should be administered at a dose range of 0.2 mg to 0.6 mg pimobendan/kg body weight per day. The preferable daily dose is 0.5 mg pimobendan/kg body weight. The dose should be divided into two administrations (0.25 mg/kg body weight each), using a suitable combination of whole, or half of tablets. One half of the dose in the morning and the other half approximately 12 hours later.

Each dose should be given approximately one hour before feeding.

This corresponds to:

One 1.25 mg chewable tablet in the morning and one 1.25 mg chewable tablet in the evening for a body weight of 5 kg.

Tablets (1.25, 5 and 10 mg tablet) are divisible in two equal parts. The veterinary medicinal product may be combined with a diuretic treatment such as furosemide.

## **9. Advice on correct administration**

Spontaneous intake by the animal or place the tablet directly in the mouth.

## **10. Withdrawal periods**

Not applicable.

## **11. Special storage precautions**

Keep out of the sight and reach of children.

For blisters: Any unused tablet portion should be returned to the blister and be used for the next administration.

Do not store above 30°C.

For bottle: Shelf life after first opening the immediate packaging: 2 months  
Keep the bottle tightly closed in order to protect from moisture.

Any unused tablet portion should be returned to the bottle and be used for the next administration.

Do not store above 25°C.

Do not use this veterinary medicinal product after the expiry date which is stated on the blister or bottle and carton after “Exp”. The expiry date refers to the last day of that month.

## **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.

## **14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

Vm 14966/5085

Pack sizes:

For blisters: Cardboard box with 3 or 8 blisters of 12 tablets.

For bottle: 35 ml bottle containing 60 tablets

Not all pack sizes may be marketed.

## **15. PID LINK (Do not print heading)**

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the ‘Product Information Database’ on [www.gov.uk](http://www.gov.uk).

## **16. Contact details**

Marketing authorisation holder:

Ceva Sante Animale  
8 rue de Logrono  
33500 Libourne  
France

Contact details to report suspected adverse reactions:

Ceva Animal Health Ltd  
Explorer House  
Mercury Park  
Wycombe Lane  
Wooburn Green  
High Wycombe  
Buckinghamshire  
HP10 0HH  
United Kingdom

Manufacturer responsible for batch release:

Ceva Santé Animale  
Boulevard de la communication  
Zone Autoroutière  
53950 LOUVERNE FRANCE

## **17. Other information**

POM-V

Veterinary Medicinal product subject to prescription  
For animal treatment only

*Gavin Hall*  
Approved: 05 November 2025