

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**HDPE Bottle (100ml)**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fiprotec Spray 2.5 mg/ml Cutaneous Spray, Solution  
Fipronil

**2. STATEMENT OF ACTIVE SUBSTANCES**

Fipronil            2.5 mg/ml

**3. PHARMACEUTICAL FORM**

Cutaneous Spray, Solution

**4. PACKAGE SIZE**

100 ml

**5. TARGET SPECIES**

Cats and Dogs

**6. INDICATION(S)**

Treatment and prevention of flea (*Ctenocephalides* spp.) and tick (*Ixodes* spp., including *Ixodes ricinus*) infestations in cats and dogs.  
Treatment of biting lice infestations in dogs (*Trichodectes canis*) and cats (*Felicola subrostratus*).

Insecticidal efficacy against new infestations with adult fleas persists for up to 2 months in cats and up to 3 months in dogs, depending on environmental challenge. The product has a persistent acaricidal efficacy for up to one month against ticks, depending on the level of environmental challenge.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Apply 3 to 6 ml per kg bodyweight. The 100ml bottle contains:

- Approximately 3 to 6 treatments for a dog weighing 5 kg, depending on hair length;
- Approximately 8 treatments for a short haired cat weighing 4 kg;
- Approximately 2 treatments for a long haired dog weighing 8 kg.

**8. WITHDRAWAL PERIOD(S)**

N/A

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP {DD/MM/YY}

Once opened use within 1 year.

**11. SPECIAL STORAGE CONDITIONS**

Highly flammable.

Do not store above 25 °C.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Beaphar B.V.  
Drostenkamp 3  
8101 BX, Raalte  
The Netherlands

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 41941/3002

**17. MANUFACTURER’S BATCH NUMBER**

Batch {number}

## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:**

**Fiprotec Spray 2.5 mg/ml Cutaneous Spray, Solution for Cats and Dogs**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Beaphar B.V.  
Drostenkamp 3  
8101 BX, Raalte  
The Netherlands  
Tel: +31 (0) 572 348 834

Manufacturer responsible for batch release:

Beaphar B.V.  
Oude Linderteseweg 9  
8102 EV Raalte  
The Netherlands

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fiprotec Spray 2.5 mg/ml Cutaneous Spray, Solution for Cats and Dogs  
Fipronil

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each ml of this clear and colourless liquid contains: Fipronil 2.5 mg.

**4. INDICATION(S)**

Treatment and prevention of flea (*Ctenocephalides* spp.) and tick (*Ixodes* spp., including *Ixodes ricinus*) infestations in cats and dogs.

Treatment of biting lice infestations in dogs (*Trichodectes canis*) and cats (*Felicola subrostratus*).

Insecticidal efficacy against new infestations with adult fleas persists for up to 2 months in cats and up to 3 months in dogs, depending on environmental challenge. The product has a persistent acaricidal efficacy for up to one month against ticks, depending on the level of environmental challenge.

**5. CONTRAINDICATIONS**

Do not use on sick or recovering animals

Do not use in rabbits, as adverse reactions and even death could occur.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

## **6. ADVERSE REACTIONS**

If licking occurs, a brief period of excessive salivation may be observed due mainly to the nature of the carrier.

Among the very rare suspected adverse reactions, temporary skin reactions such as reddening, itching or hair loss have been reported after use. Exceptionally, excessive salivation, reversible neurological signs (hyperaesthesia, depression, nervous signs), vomiting or respiratory signs have been observed after use.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated )
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

Alternatively you can report via your national reporting system {national system details}.

## **7. TARGET SPECIES**

Cats and dogs

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Route of administration: spray pump for cutaneous use.

Dosage: It is important to dampen the coat down to the skin. Depending on the length of hair, apply 3 to 6 ml per kg bodyweight, (7.5 to 15mg of active ingredient per kg bodyweight) i.e. 6 to 12 pump applications per kg bodyweight of the 100ml presentation.

The 100ml bottle contains:

- Approximately 3 to 6 treatments for a dog weighing 5 kg, depending on hair length;
- Approximately 8 treatments for a short haired cat weighing 4 kg;
- Approximately 2 treatments for a long haired dog weighing 8 kg.

Method of administration:

Spray the entire body of the animal, and apply from a distance of approximately 10-20cm.

Apply against the lay of the hair and make sure that the entire coat of the animal is dampened. Ruffle the coat, especially in long haired animals, so that the product penetrates down to the skin.

For treatment of the head region, and when treating young or nervous pets, application may be carried out by spraying onto a gloved hand and rubbing the product into the coat. Allow to dry naturally. Do not towel dry.

Properties : The formulation contains a coating agent. Therefore, spraying builds up a film and makes the fur glossy.

In the absence of safety studies, the minimum treatment interval is 4 weeks.

## **9. ADVICE ON CORRECT ADMINISTRATION**

## **10. WITHDRAWAL PERIOD(S)**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Highly flammable.

Do not store above 25 °C.

Do not use this veterinary medicinal product after the expiry date which is stated on the bottle. The expiry date refers to the last day of that month.

Shelf life after first opening the container: 1 year.

The discard date (1 year after opening) should be recorded. Any product remaining after this date, should be discarded.

## **12. SPECIAL WARNING(S)**

Special warnings for each target species:

For optimum efficacy, it is not recommended to bathe or shampoo animals in the two days prior to or following treatment with the product. Bathing or shampooing up to four times in two months has been shown not to significantly reduce the remaining effect of the product. Monthly treatment is recommended when more frequent shampooing is carried out.

The possibility that other animals in the same household can be a source of re-infection with fleas, ticks and biting lice should be considered, and these should be treated as necessary with an appropriate product.

Fleas from pets often infest the animal's basket, bedding and regular resting areas, such as carpets and soft furnishings. In cases of massive infestation and when initiating control measures these areas should also be treated with a suitable insecticide, and vacuumed regularly.

There may be an attachment of single ticks. For this reason a transmission of infectious diseases cannot be completely excluded if conditions are unfavourable.

The product is not suitable for direct treatment of the environment.



Given the volume delivered by the spray pump (0.5ml/pump) the product is very suitable for use in cats and dogs up to 5-10kg bodyweight, depending on the length of hair (see also section Dosage for each species, route(s) and method of administration.) but not adequate for medium and large dogs.”

Special precautions for use in animals:

It is important to make sure that animals do not lick each other following treatment.

Do not exceed the recommended dosage.

Avoid contact with the animal's eyes. In the case of accidental eye contact immediately and thoroughly flush the eyes with water.

Do not spray directly onto areas of broken skin.

Allow treated animals to dry in a well ventilated room (see section Special precautions to be taken by the person administering the veterinary medicinal product to animals).

Do not confine animals in an enclosed space or pet carrier until the coat is totally dry.

In the absence of specific tolerance and efficacy data, the product is not recommended for the treatment of species other than cats and dogs.

Puppies and kittens from 2 days of age may be safely treated.

Keep treated animals away from fires or other sources of heat, and surfaces likely to be affected by the alcohol spray, for at least 30 minutes following spraying and until the fur is totally dry. Do not spray on a naked flame or any incandescent material.

For cutaneous use only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This product can cause mucous membrane and eye irritation. Therefore, contact of the product with mouth and eyes should be avoided.

People with asthma or a known hypersensitivity (allergy) to insecticides or alcohol should avoid contact with the veterinary medicinal product. Avoid contents coming into contact with the fingers. If this occurs, wash hands with soap and water. If irritation occurs, seek medical advice.

After accidental eye contact rinse the eye carefully with plain water.

Treated animals should not be handled until the fur is dry, and children should not be allowed to play with treated animals until the fur is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals are not allowed to sleep with owners, especially children.

Spray animals in the open air or a well ventilated room.

Do not breathe spray. Do not smoke, drink or eat during application.

Wear PVC or nitrile gloves during treatment of animals. It is recommended to wear a waterproof apron for the protection of clothing.

If clothing becomes heavily wetted with the product, it should be removed and washed before re-use.

Dispose of gloves after use and then wash hands with soap and water. Wash splashes from skin with soap and water immediately. If irritation occurs, seek medical advice.

Treatment of more than one animal: Good ventilation is particularly important where several animals are to be treated. Treat the animals outside, or reduce the build-up of vapour by removing the animals from the treatment room while the alcohol is evaporating and ensure that the treatment room is well ventilated between individual treatments. In addition, ensure that the drying room is well ventilated and avoid housing several recently treated animals within the same air space.

Other precautions:

Fipronil may adversely affect aquatic organisms.

Dogs should not be allowed to swim in watercourses for 2 days after application.

Pregnancy and Lactation :

The spray is safe to use on lactating dogs.

Do not use on pregnant cats and dogs or on nursing cats.

Interaction with other medicinal products and other forms of interaction:

Do not use simultaneously with other flea products which are applied directly onto the animal.

Overdose (symptoms, emergency procedures, antidotes):

The risk of experiencing adverse effects may increase when overdosing, so animals should always be treated with the correct dose according to bodyweight.

In case of overdosing, appropriate symptomatic treatment should be started by a veterinarian.

Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Fipronil may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty containers.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

September 2022

**15. OTHER INFORMATION**

100ml bottle

Approved 20 October 2022

