

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

{Carton for 20 ml, 50 ml and 100ml vial}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Rheumocam 5 mg/ml solution for injection for cattle and pigs  
Meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam: 5 mg/ml  
Ethanol (96%): 159.8 mg/ml

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE**

20 ml  
50 ml  
100 ml

**5. TARGET SPECIES**

Cattle (calves and young cattle) and pigs

**6. INDICATIONS**

**Cattle:**

Acute respiratory infection.  
Diarrhoea in calves of over one week of age and young, non-lactating cattle.

**Pigs:**

Non-infectious locomotor disorders.  
Post operative pain associated with minor soft tissue surgery such as castration.

## 7. METHOD AND ROUTES OF ADMINISTRATION

### **Cattle:**

Single subcutaneous or intravenous injection.

### **Pigs:**

Single intramuscular injection. If required, a second administration can be given after 24 hours.

Single intramuscular injection before surgery.

Take care of accurate dosing, use of appropriate dosing device and estimation of body weight.

Read the package leaflet before use.

## 8. WITHDRAWAL PERIODS

Withdrawal period

**Cattle:** meat and offal: 15 days.

**Pigs:** meat and offal: 5 days

## 9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

## 10. EXPIRY DATE

EXP {month/year}

Shelf-life of broached vial: 28 days.

Once broached, use by.....

## 11. SPECIAL STORAGE CONDITIONS

## 12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Read the package leaflet before use.

## 13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Chanelle Pharmaceuticals Manufacturing Ltd.,  
Loughrea,  
Co. Galway,  
Ireland.

**16. MARKETING AUTHORISATION NUMBER**

Vm 08749/5025

**17. MANUFACTURER'S BATCH NUMBER**

BN {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

{Label for 100 ml}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Rheumocam 5 mg/ml solution for injection for cattle and pigs  
Meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam 5 mg/ml

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE**

100 ml,

**5. TARGET SPECIES**

Cattle (calves and young cattle) and pigs

**6. INDICATIONS**

Read package leaflet before use.

**7. METHOD AND ROUTES OF ADMINISTRATION**

**Cattle:** SC or IV injection.

**Pigs:** IM injection.

Read the package leaflet before use

**8. WITHDRAWAL PERIODS**

Withdrawal period

**Cattle:** meat and offal: 15 days

**Pigs:** meat and offal: 5 days.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use

**10. EXPIRY DATE**

EXP {month/year}

Once broached, use by...

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF USUED PRODUCTS OR WASTE MATERIALS, IF ANY**

Read the package leaflet before use

**13. THE WORDS 'FOR ANIMAL TREATMENT ONLY' AND CONDITIONS OF RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only-to be supplied only on veterinary prescription.

**14. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Chanelle Pharmaceuticals Manufacturing Ltd  
Loughrea  
Co. Galway  
Ireland

**16. MARKETING AUTHORISATION NUMBER**

Vm 08749/5025 100 ml

**17. MANUFACTURER'S BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

{Label for 20 ml and 50 ml bottles}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Rheumocam 5 mg/ml Solution for injection for Cattle and Pigs  
Meloxicam

**2. QUANTITY OF THE ACTIVE SUBSTANCE**

Meloxicam 5 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

20 ml  
50 ml

**4. ROUTES OF ADMINISTRATION**

**Cattle:** SC or IV.  
**Pigs:** IM.

**5. WITHDRAWAL PERIOD(S)**

Withdrawal period  
**Cattle:** meat and offal: 15 days  
**Pigs:** meat and offal: 5 days.

**6. BATCH NUMBER**

Lot {number}

**7. EXPIRY DATE**

EXP {month/year}  
Once broached use by ...

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only

**PACKAGE LEAFLET:  
Rheumocam 5 mg/ml solution for injection for cattle and pigs**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Chanelle Pharmaceuticals Manufacturing Ltd., Loughrea, Co. Galway, Ireland

Manufacturers responsible for the batch release:

Chanelle Pharmaceuticals Manufacturing Ltd., Loughrea, Co. Galway, Ireland

and

Eurovet Animal Health B.V., Handelsweg 25, 5531 AE Bladel, The Netherlands

and

Labiana Life Sciences, S.A., C/ Venus, 26, Pol. Ind. Can Parellada, Tarrasa, 08228  
Barcelona

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Rheumocam 5 mg/ml solution for injection for cattle and pigs  
Meloxicam

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS**

One ml contains:

Meloxicam	5 mg
Ethanol (96%)	159.8 mg

Clear, yellow solution.

**4. INDICATIONS**

**Cattle:**

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

## **Pigs:**

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For the relief of post operative pain associated with minor soft tissue such as castration.

## **5. CONTRAINDICATIONS**

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.

Do not use in pigs less than 2 days old.

## **6. ADVERSE REACTIONS**

Subcutaneous, intramuscular as well as intravenous administration is well tolerated; only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In very rare cases anaphylactoid reactions which may be serious (including fatal) may occur and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Cattle (calves and young cattle) and pigs.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

### **Cattle:**

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 10 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

### **Pigs:**

#### Locomotor disorders:

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2.0 ml/25 kg body weight). If required, a second administration of meloxicam can be given after 24 hours.

#### Reduction of post-operative pain:

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 0.4 ml/5 kg body weight) before surgery.

Particular care should be taken with regard to the accuracy of dosing including the use of an appropriate dosing device and careful estimation of bodyweight.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Avoid introduction of contamination during use.

## **10. WITHDRAWAL PERIODS**

**Cattle:** meat and offal: 15 days.

**Pigs:** meat and offal: 5 days.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use after the expiry date (EXP) stated on the carton and vial.

Shelf life after first opening the container: 28 days.

## **12. SPECIAL WARNINGS**

Treatment of piglets with Rheumocam before castration reduces post-operative pain. To obtain pain relief during surgery co-medication with an appropriate anaesthetic/sedative is needed.

To obtain the best possible relieving effect post surgery Rheumocam should be administered 30minutes before surgical intervention.

Special precautions for use in animals:

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental self-injection may give rise to pain. People with known hypersensitivity to Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show this package leaflet or the label to the physician.

Use during pregnancy and lactation:

Cattle: Can be used during pregnancy.

Pigs: Can be used during pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

Overdose (symptoms, emergency procedures, antidotes):

In the case of overdose symptomatic treatment should be initiated.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Waste materials should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**


26/10/2020

**15. OTHER INFORMATION**

Cardboard box containing 1 colourless glass injection vial of 20 ml, 50 ml or 100 ml. Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**United Kingdom**  
Chanelle Vet UK Ltd  
Freemans House  
127 High Street  
Hungerford  
RG17 0DL  
UK

A handwritten signature in black ink, consisting of several loops and a final flourish.

Approved: 11 June 2021