

PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Carton}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Librela 30 mg Solution for injection

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each 1 ml contains 30 mg bedinvetmab.

3. PACKAGE SIZE

1 x 1 ml

2 x 1 ml

6 x 1 ml

4. TARGET SPECIES

Dogs.

5. INDICATION(S)

6. ROUTES OF ADMINISTRATION

Subcutaneous use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

Once broached use immediately.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Do not freeze.

Store in the original package.

Protect from light.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Zoetis UK Limited
1st Floor, Birchwood Building
Springfield Drive
Leatherhead
Surrey
KT22 7LP

14. MARKETING AUTHORISATION NUMBER

Vm 42058/5033

15. BATCH NUMBER

Lot {number}

16. SPECIAL WARNING(S), IF NECESSARY

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

POM-V
To be supplied only on veterinary prescription.

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS {Vial label}**

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Librela



**2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE
SUBSTANCE(S)**

bedinvetmab 30 mg/ml

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Once broached use immediately.

5. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

6. ROUTE(S) OF ADMINISTRATION

SC

7. WITHDRAWAL PERIOD

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Librela 30 mg solution for injection for dogs

2. COMPOSITION

Active substance:

Each vial of 1 ml contains 5 mg, 10 mg, 15 mg, 20 mg or 30 mg bedinvetmab*.

* Bedinvetmab is a canine monoclonal antibody expressed through recombinant techniques in Chinese hamster ovary (CHO) cells.

The product should appear clear to slightly opalescent without any visible particles.

3. TARGET SPECIES

Dogs.

4. INDICATIONS FOR USE

For the alleviation of pain associated with osteoarthritis in dogs.

5. CONTRAINDICATIONS

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in dogs under 12 months.

Do not use in animals intended for breeding.

Do not use in pregnant or lactating animals.

6. SPECIAL WARNING(S)

Special warnings:

This veterinary medicinal product may induce transient or persistent anti-drug antibodies. The induction of such antibodies is uncommon and may have no effect or may result in a decrease in efficacy in animals that responded to treatment previously.

If no or limited response is observed within one month after initial dosing, an improvement in response may be observed after administration of a second dose one month later. However, if the animal does not show a better response after the second dose, the veterinary surgeon should consider alternative treatments.

Special precautions for safe use in the target species:

Where a dog has not been able to properly exercise prior to treatment due to its clinical condition, it is recommended that the dog is gradually (over a few weeks) allowed to increase the amount of exercise they take (to prevent overexercise by some dogs).

Caution should be used when treating patients with the following pre-existing conditions: immune-mediated haemolytic anaemia, immune-mediated polyarthritis, immune-mediated thrombocytopenia.

Caution should be used when treating patients with pre-existing seizure disorder.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Hypersensitivity reactions, including anaphylaxis, could potentially occur in the case of accidental self-injection. Repeated self-administration may increase the risk of hypersensitivity reactions.

The importance of nerve growth factor (NGF) in ensuring normal foetal nervous system development is well-established and laboratory studies conducted on non-human primates with human anti-NGF antibodies have shown evidence of reproductive and developmental toxicity. Pregnant women, women trying to conceive and breastfeeding women should take extreme care to avoid accidental self-injection.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation or in breeding dogs. Laboratory studies with human anti-NGF antibodies in cynomolgus monkeys have shown evidence of teratogenic and foetotoxic effects.

Do not use in pregnant or lactating animals.

Fertility:

Do not use in breeding animals.

Interaction with other medicinal products and other forms of interaction:

In a laboratory study over a 2-week period in young, healthy dogs without osteoarthritis, this veterinary medicinal product had no adverse effect when concomitantly administered with a non-steroidal anti-inflammatory product (carprofen).

There are no safety data on the concurrent long-term use of NSAIDs and bedinvetmab in dogs. In clinical trials in humans, rapidly progressive osteoarthritis has been reported in patients receiving humanised anti-NGF monoclonal antibody therapy. The incidence of these events increased with high doses and in those human patients that received long-term (more than 90 days) non-steroidal anti-inflammatory drugs (NSAIDs) concomitantly with an anti-NGF monoclonal antibody. Dogs have no reported equivalent of human rapidly progressive osteoarthritis.

No other laboratory studies on the safety of concomitant administration of this veterinary medicinal product with other veterinary medicinal products have been conducted. No interactions were observed in field studies where this veterinary medicinal product was administered concomitantly with veterinary medicinal products containing parasiticides, antimicrobials, topical antiseptics with or without corticosteroids, antihistamines and vaccines.

If a vaccine(s) is to be administered at the same time as treatment with this veterinary medicinal product, the vaccine(s) should be administered at a different site to that of the veterinary medicinal products administration, to reduce any potential impact on immunogenicity of the vaccine.

Overdose:

No adverse reactions, except mild reactions at the injection site, were observed in a laboratory overdose study when the veterinary medicinal product was administered for 7 consecutive monthly doses at 10 times the maximum recommended dose.

In case of adverse clinical signs after an overdose the dog should be treated symptomatically.

Major incompatibilities:

Do not mix with any other veterinary medicinal product.

7. ADVERSE EVENTS

Dogs:

Uncommon (1 to 10 animals / 1,000 animals treated):	Injection site reaction (e.g. injection site swelling, injection site warmth) ¹ .
Rare (1 to 10 animals / 10,000 animals treated):	Diarrhoea, Emesis. Ataxia ² . Polyuria, Urinary incontinence. Anorexia ³ , Lethargy, Polydipsia.
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Hypersensitivity reaction (anaphylaxis, facial swelling, pruritus) ⁴ Low amounts of red blood cells (immune-mediated haemolytic anaemia), Joint pain, Lameness and Swelling in multiple joints (immune-mediated polyarthritis), Low amounts of thrombocytes (immune-mediated thrombocytopenia). Paresis, Paralysis, Seizure.

¹ Mild

² Including proprioceptive ataxia.

³ Often related to a transient reduced appetite.

⁴ In case of such reactions, appropriate symptomatic treatment should be administered.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Subcutaneous use.

Dosage and treatment schedule:

The recommended dose is 0.5-1.0 mg/kg bodyweight, once a month.

Dogs weighing <5.0 kg:

Aseptically withdraw 0.1 ml/kg from a single 5 mg/ml vial and administer subcutaneously.

For dogs between 5 and 60 kg administer the entire content of the vial (1 ml) according to the table below:

Bodyweight (kg) of dog	Librela strength (mg) to be administered				
	5	10	15	20	30
5.0-10.0	1 vial				
10.1-20.0		1 vial			
20.1-30.0			1 vial		
30.1-40.0				1 vial	
40.1-60.0					1 vial
60.1-80.0				2 vials	
80.1-100.0				1 vial	1 vial
100.1-120.00					2 vials

For dogs above 60 kg, the contents of more than one vial are required to administer a single dose. In those cases, withdraw the content from each required vial into the same syringe and administer as a single subcutaneous injection (2 ml).

9. ADVICE ON CORRECT ADMINISTRATION

None.

10. WITHDRAWAL PERIOD(S)

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Store in the original package.

Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: use immediately.

12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBER AND PACK SIZES

Vm 42058/5033

Clear glass Type I vials with fluorobutyl rubber stopper.

Pack sizes:

Cardboard box with 1, 2 or 6 vials of 1 ml

Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder and contact details to report suspected adverse reactions:

United Kingdom (Great Britain)

Zoetis UK Limited
1st Floor, Birchwood Building
Springfield Drive
Leatherhead Surrey
KT22 7LP
Tel: +44 (0) 345 300 8034

Marketing authorisation holder and manufacturer responsible for batch release:

Zoetis Belgium
Rue Laid Burniat 1
1348 Louvain-La-Neuve
Belgium

Local representatives and contact details to report suspected adverse reactions:

België/Belgique/Belgien

Zoetis Belgium
Mercuriusstraat 20
BE-1930 Zaventem
Tél/Tel: +32 (0) 800 99 189

Lietuva

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belgija
Tel: +370 610 05088

Република България

Zoetis Belgium
Rue Laid Burniat 1
1348 Louvain-La-Neuve
Белгия
Тел: +359 888 51 30 30

Luxembourg/Luxemburg

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belsch
Tél/Tel: +32 (2) 746 80 11

Česká republika

Zoetis Česká republika, s.r.o.
náměstí 14. října 642/17
CZ 150 00 Praha
Tel: +420 257 101 111

Magyarország

Zoetis Hungary Kft.
Csörsz u. 41.
HU-1124 Budapest
Tel.: +36 1 224 5200

Danmark

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 København
Tlf: +45 70 20 73 05
adr.scandinavia@zoetis.com

Malta

Agrimed Limited
Mdina Road, Zebbug ZBG 9016,
MT
Tel: +356 21 465 797

Deutschland

Zoetis Deutschland GmbH
Schellingstr. 1
DE-10785 Berlin
Tel: +49 30 2020 0049
tierarzneimittelsicherheit@zoetis.com

Nederland

Zoetis B.V.
Rivium Westlaan 74
NL-2909 LD Capelle aan den IJssel
Tel: +31 (0)10 714 0900

Eesti

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belgia
Tel: +370 610 05088

Norge

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 København
Danmark
Tlf: +47 23 29 86 80
adr.scandinavia@zoetis.com

Ελλάδα

Zoetis Hellas S.A.
Φραγκοκκλησιάς 7, Μαρούσι
EL-15125 Αττική
Τηλ: +30 210 6791900

España

Zoetis Spain, S.L.
Parque Empresarial Vía Norte Edificio
nº1,
c/ Quintanavides nº13
ES-28050 Madrid
Tel: +34 91 4191900

France

Zoetis France
10 rue Raymond David
FR-92240 Malakoff
Tél: +33 (0)800 73 00 65

Hrvatska

Zoetis B.V.
Podružnica Zagreb za promidžbu
Petra Hektorovića 2
HR-10000 Zagreb
Tel: +385 1 6441 462

Ireland

Zoetis Belgium S.A. (Irish Branch)
2nd Floor, Building 10,
Cherrywood Business Park,
Loughlinstown,
Co. Dublin,
IE – Dublin D18 T3Y1
Tel: +353 (0) 1 256 9800

Ísland

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 København
Danmörku
Sími: +45 70 20 73 05
adr.scandinavia@zoetis.com

Österreich

Zoetis Österreich GmbH
Floridsdorfer Hauptstr. 1
AT-1210 Wien
Tel: +43 (0)1 2701100 100

Polska

Zoetis Polska Sp. z o.o.
ul. Postępu 17B
PL - 02-676 Warszawa
Tel.: +48 22 2234800

Portugal

Zoetis Portugal Lda.
Lagoas Park, Edifício 10
PT-2740-271 Porto Salvo
Tel: +351 21 042 72 00

România

Zoetis România S.R.L.
Expo Business Park, 54A Aviator
Popișteanu, Clădirea 2, Etaj 1-3, Sector
1,
București, 012095 - RO
Tel: +40785019479

Slovenija

Zoetis B.V.
Podružnica Zagreb za promidžbu
Petra Hektorovića 2,
10000 Zagreb,
Hrvaška
Tel: +385 1 6441 462

Slovenská republika

Zoetis Česká republika, s.r.o.
náměstí 14. října 642/17
150 00 Praha
Česká republika
Tel: +420 257 101 111

Italia

Zoetis Italia S.r.l.
Via Andrea Doria 41M,
IT-00192 Roma
Tel: +39 06 3366 8111

Suomi/Finland

Zoetis Finland Oy
Bulevardi 21 / SPACES
FI-00180 Helsinki/Helsingfors
Suomi/Finland
Puh/Tel: +358 10 336 7000
laaketurva@zoetis.com

Κύπρος

Zoetis Hellas S.A.
Φραγκοκκλησιάς 7, Μαρούσι
15125, Αττική
Ελλάδα
Τηλ: +30 210 6791900

Sverige

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 Köpenhamn
Danmark
Tel: +46 (0) 76 760 0677
adr.scandinavia@zoetis.com

Latvija

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belgija
Tel: +370 610 05088

United Kingdom (Northern Ireland)

Zoetis Belgium S.A. (Irish Branch)
2nd Floor, Building 10,
Cherrywood Business Park,
Loughlinstown,
Co. Dublin,
IE – Dublin D18 T3Y1
Tel: +353 (0) 1 256 9800

United Kingdom (Great Britain)

Zoetis UK Limited
1st Floor, Birchwood Building
Springfield Drive
Leatherhead
Surrey, KT22 7LP
UK
Tel: +44 (0) 345 300 8034

17. OTHER INFORMATION

Gavin Hall
Approved: 14 January 2026