

**INSERT TEXT**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE  
FOR BATCH RELEASE, IF DIFFERENT**

**(EU)**

Norbrook Laboratories (Ireland) Ltd.  
Rossmore Industrial Estate  
Monaghan  
Ireland

**(UK)**

Norbrook Laboratories Limited  
Station Works  
Camlough Road  
Newry  
Co. Down  
BT35 6JP  
Northern Ireland

**MANUFACTURER RESPONSIBLE FOR BATCH RELEASE:**

Norbrook Manufacturing Ltd.  
Rossmore Industrial Estate  
Monaghan  
Ireland

Norbrook Laboratories Limited  
Station Works  
Camlough Road  
Newry  
Co. Down  
BT35 6JP  
Northern Ireland

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Synoclav 50 mg Tablets for Dogs and Cats (UK)  
Clavobay 50 mg Tablets for Dogs and Cats (AT, BE, FR, IS, IT, NL, NO, PT)  
Clavobay Vet 40 mg/10 mg Tablets for Dogs and Cats (SE)  
Clavubay 50 mg Tablets for Dogs and Cats (ES)  
Clavobay Vet 50 mg Tablets for Dogs and Cats (DK)

### 3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

<b>Active Ingredients:</b>	mg per tablet
Amoxicillin	40
(as amoxicillin trihydrate)	45.9
Clavulanic acid	10
(as Potassium clavulanate)	11.9
<b>Excipients:</b>	
Carmoisine Lake (E122)	0.245

### 4. INDICATION(S)

Synuclav 50 mg Tablets are indicated for treatment of the following infections caused by  $\beta$ -lactamase producing strains of bacteria sensitive to amoxicillin in combination with clavulanic acid:

- Skin infections (including superficial and deep pyodermas) caused by susceptible Staphylococci.
- Urinary tract infections caused by susceptible Staphylococci or *Escherichia coli*.
- Respiratory infections caused by susceptible Staphylococci.
- Enteritis caused by susceptible *Escherichia coli*.

It is recommended to carry out suitable tests for sensitivity testing when initiating the treatment. The treatment should only proceed if sensitivity is proven to the combination.

### 5. CONTRAINDICATIONS

Do not use in animals with known cases of hypersensitivity to penicillin or other substances of the beta-lactam group.

Do not use in rabbits, guinea pigs, hamsters or gerbils.

Do not use in animals with serious dysfunction of the kidneys accompanied by anuria and oliguria.

Do not use where resistance to this combination is known to occur.

## 6. ADVERSE REACTIONS

Hypersensitivity reactions unrelated to dose can occur with these agents.

Gastrointestinal symptoms (diarrhoea, vomiting) may occur after administration of the product.

Allergic reactions (e.g. skin reactions, anaphylaxia) may occasionally occur.

In case of occurrence of allergic reaction, the treatment should be withdrawn.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Dogs & Cats.

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

12.5 mg combined actives/kg bodyweight twice daily. The recommended dose of 12.5 mg per kg bodyweight is equivalent to one 50 mg tablet per 4 kg bodyweight.

**Dosage frequency:** The following table is intended as a guide to dispensing Synuclav 50mg Tablets at the standard dose rate of 12.5 mg/kg twice daily.

Bodyweight (kg)	Number of tablets per dose twice daily
	50 mg
1-2	●
3-4	●
5-6	● ●
7-8	● ● ●
9-10	● ● ● ●
11-12	● ● ● ● ●
13-14	● ● ● ● ● ●
15-16	● ● ● ● ● ● ●
17-18	● ● ● ● ● ● ● ●

Duration of therapy: Routine cases involving all indications: The majority of cases respond to between 5 and 7 days therapy.

Chronic or refractory cases: In these cases where there is considerable tissue damage, a longer course of therapy may be required in that it allows sufficient time for damaged tissue to repair.

## **9. ADVICE ON CORRECT ADMINISTRATION**

By the oral route. The tablets may be crushed and added to a little food.

## **10. WITHDRAWAL PERIOD**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Store in the original package in order to protect from moisture.

Do not use after the expiry date stated on the blister or tubs; Keep out of the reach and sight of children; Keep the container in the outer carton

## **12. SPECIAL WARNINGS**

Inappropriate use of the product may increase the prevalence of bacteria resistance to amoxicillin/ clavulanic acid.

In animals with hepatic and renal failure, the dosing regimen should be carefully evaluated.

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies. Narrow spectrum antibacterial therapy should be used for first line treatment where susceptibility testing suggests likely efficacy of this approach.

Dogs and cats diagnosed with *Pseudomonas* infections should not be treated with this antibiotic combination.

Studies in laboratory animals have not produced any evidence of teratogenic effects.

Use only according to the benefit/risk assessment by the responsible veterinarian.

The potential for allergic cross-reactivity with other penicillins should be considered.

Caution is advised in the use in small herbivores other than those reported in contradictions.

Chloramphenicol, macrolides, sulfonamides and tetracyclines may inhibit the antibacterial effect of penicillins because of the rapid onset of bacteriostatic action.

Penicillins may increase the effect of aminoglycosides.

### **Operator Warnings:**

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Handle this product with great care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Wash hands after use.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

### **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

November 2019

### **15. OTHER INFORMATION**

Resistance to many antibiotics is caused by beta-lactamase enzymes which destroy the antibiotic before it can act on the bacteria themselves. The clavulanate in Synuclav Tablets counteracts this defence mechanism by inactivating the beta-lactamases, thus rendering the organisms sensitive to amoxicillin's rapid bactericidal effect, at concentrations readily attainable in the body.

*In vitro* potentiated amoxicillin is active against a wide range of clinically important aerobic and anaerobic bacteria including:

#### Gram-positive:

Staphylococci (including  $\beta$ -lactamase producing strains)  
Clostridia  
Streptococci

#### Gram-negative:

*Escherichia coli* (including most  $\beta$ -lactamase producing strains)  
*Campylobacter* spp  
*Pasteurellae*  
*Proteus* spp

The product is supplied in high-density polyethylene tubs with a polypropylene screw cap lid containing 100 tablets and in high-density polyethylene tubs with a polyethylene

screw cap lid containing 500 tablets. A sachet of desiccant is included in each container. The product is also presented in packs containing 2, 10 and 50 blister strips (aluminium-aluminium) each containing 10 tablets per strip.

Not all pack sizes may be marketed.

Vm: 02000/4224

ManA: 2000

**DISTRIBUTED BY:**

MiGroup, CVS House  
Owen Road  
Diss  
Norfolk  
IP22 4ER  
United Kingdom

## LABEL TEXT

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Synuclav 50 mg Tablets for Dogs and Cats (UK)  
Clavobay 50 mg Tablets for Dogs and Cats (AT, BE, FR, IS, IT, NL, NO, PT)  
Clavobay Vet 40 mg/10 mg Tablets for Dogs and Cats (SE)  
Clavobay 50 mg Tablets for Dogs and Cats (ES)  
Clavobay Vet 50 mg Tablets for Dogs and Cats (DK)

### 2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each tablet contains:

Amoxicillin (as amoxicillin trihydrate)	40 mg,
Clavulanic acid (as Potassium clavulanate)	10 mg
Excipient:	
Carmoisine Lake (E122)	0.245 mg.

### 3. PHARMACEUTICAL FORM

Tablet.

### 4. PACKAGE SIZE

100/500 TABLETS

### 5. TARGET SPECIES

Dogs & Cats.

### 6. INDICATION(S)

See package leaflet for indications

### 7. METHOD AND ROUTE(S) OF ADMINISTRATION

To be given orally. The tablet may be crushed and added to a little food.

#### **8. WITHDRAWAL PERIOD**

Not applicable.

#### **9. SPECIAL WARNING(S), IF NECESSARY**

Penicillins and cephalosporins may occasionally cause severe allergic reactions.  
See package leaflet for user warnings.

#### **10. EXPIRY DATE**

EXP: {month/year}

#### **11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.  
Store in the original package in order to protect from moisture.

#### **12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

#### **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

##### **FOR ANIMAL TREATMENT ONLY**

POM-V

To be supplied only on veterinary prescription

#### **14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”**

**KEEP OUT OF THE REACH AND SIGHT OF CHILDREN**



**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Norbrook Laboratories Limited  
Station Works  
Camlough Road  
Newry  
Co. Down  
BT35 6JP  
Northern Ireland

**16. MARKETING AUTHORISATION NUMBER**

ManA 2000

Vm 02000/4224

**17. MANUFACTURER'S BATCH NUMBER**

BN:

A handwritten signature in black ink, consisting of several loops and a long horizontal stroke at the end.

Approved 30 January 2020