

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS**

Label

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Label

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Animeloxan, 5 mg/ml, solution for injection for dogs and cats
Meloxicam

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Meloxicam 5 mg/ml

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

10 ml, 20 ml, 25 ml

4. ROUTE(S) OF ADMINISTRATION

Dogs: IV or SC
Cats: SC

5. WITHDRAWAL PERIOD

6. BATCH NUMBER

Lot

7. EXPIRY DATE

EXP:
Once broached, use by 28 days.

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

ADDITIONAL

aniMedica GmbH, Im Südfeld 9, D-48308 Senden-Bösensell

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box

<PARTICULARS TO APPEAR ON THE OUTER PACKAGE>

Cardboard box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Animeloxan, 5 mg/ml, solution for injection for dogs and cats
Meloxicam

2. STATEMENT OF ACTIVE SUBSTANCES

1 ml contains

Active substance:

Meloxicam 5 mg

3. PHARMACEUTICAL FORM

Solution for injection

4. PACKAGE SIZE

10 ml, 20 ml, 25 ml, 5 x 10 ml, 5 x 20 ml, 5 x 25 ml

5. TARGET SPECIES

Dogs, cats

6. INDICATION(S)

Read the package leaflet before use.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

For subcutaneous (dogs, cats) or intravenous (dogs) injection.

Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Not applicable.

9. SPECIAL WARNING(S), IF NECESSARY

Pregnant women should take extra care when handling this product. See full user warnings for details.

10. EXPIRY DATE

EXP {month/year}
Once broached, use by 28 days.

11. SPECIAL STORAGE CONDITIONS

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: See package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only.
To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

aniMedica GmbH
Im Südfeld 9
48308 Senden-Bösensell
Germany

Distributor
FORTE Healthcare Ltd.
Cougar Lane
Naul, Co. Dublin
Ireland

16. MARKETING AUTHORISATION NUMBER(S)

Vm 24745/4014

17. MANUFACTURER’S BATCH NUMBER

<Batch> <Lot> <BN> {number}

PACKAGE LEAFLET
Animeloxan, 5 mg/ml, solution for injection for dogs and cats

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER
AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE
FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

aniMedica GmbH
Im Südfeld 9
48308 Senden-Bösensell
Germany

Manufacturer responsible for batch release:

aniMedica GmbH
Im Südfeld 9
48308 Senden-Bösensell
Germany

or

Industrial Veterinaria, S.A.
Esmeralda 19
Esplugues de Llobregat
08950 Barcelona
Spain

Distributor

FORTE Healthcare Ltd.
Cougar Lane
Naul, Co. Dublin
Ireland

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Animeloxan 5 mg/ml, solution for injection for dogs and cats
Meloxicam

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER
INGREDIENT(S)**

1 ml contains

Active substance:

Meloxicam 5 mg

Excipients:

N-Methylpyrrolidone 554.04 mg
Ethanol, anhydrous 110.60 mg

Clear, yellow solution

4. INDICATION(S)

Dogs:

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopaedic and soft tissue surgery.

Cats:

Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery.

5. CONTRAINDICATIONS

Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals less than 6 weeks of age nor in cats of less than 2 kg.

6. ADVERSE REACTIONS

Typical adverse drug reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood, apathy and renal failure have occasionally been reported. In very rare cases elevated liver enzymes have been reported.

In very rare cases, haemorrhagic diarrhoea, haematemesis and gastrointestinal ulceration have been reported. These side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

In very rare cases anaphylactoid reactions may occur and should be treated symptomatically.

Injection site pain.

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

7. TARGET SPECIES

Dogs, cats

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Solution for injection.

Dosage for each species

Dogs: single administration of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg).

Cats: single administration of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg).

Method and routes of administration

Dogs:

Musculo-skeletal disorders: single subcutaneous injection.

Animeloxan 1.5 mg/ml oral suspension for dogs may be used for continuation of treatment at a dosage of 0.1 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Reduction of post-operative pain (over a period of 24 hours): single intravenous or subcutaneous injection before surgery, for example at the time of induction of anaesthesia.

Cats:

Reduction of post-operative pain: single subcutaneous injection before surgery, for example at the time of induction of anaesthesia.

9. ADVICE ON CORRECT ADMINISTRATION

Particular care should be taken with regard to the accuracy of dosing.
Avoid introduction of contamination during use.

10. WITHDRAWAL PERIOD

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature storage conditions.

Shelf-life after first opening the container: 28 days

Do not use this veterinary medicinal product after the expiry date which is stated on the bottle after EXP. The expiry date refers to the last date of that month.

12. SPECIAL WARNING(S)

Special precautions for use in animals

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of renal toxicity.

During anaesthesia, monitoring and fluid therapy should be considered as standard practice.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

If accidental skin contact occurs, wash the affected area thoroughly. Wash hands after use.

NSAIDs and other prostaglandin inhibitors are known to have adverse effects on pregnancy and/or embryofoetal development.

Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects.

Women of childbearing age, pregnant women or women suspected of being pregnant should use the veterinary medicinal product with serious caution to avoid accidental self-injection.

Use during pregnancy, lactation or lay

Do not use in pregnant or lactating animals.

The safety of the veterinary medicinal product has not been established in dogs and cats during pregnancy, lactation or in animals intended for breeding. Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Animesoxan must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic drugs should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such veterinary medicinal products should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacological properties of the products used previously.

Overdose (symptoms, emergency procedures, antidotes), if necessary

In the case of over dosage symptomatic treatment should be initiated.

Incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

October 2023

OTHER INFORMATION

Clear glass (type I) bottles of 10 ml, 20 ml and 25 ml, each closed with bromobutyl rubber stoppers and fixed with Aluminium caps or Aluminium/PP flip caps.

Available in cardboard cartons containing:

1 x 10 ml or 5 x 10 ml


1 x 20 ml or 5 x 20 ml

1 x 25 ml or 5 x 25 ml.

Not all pack sizes may be marketed.

MA-Nr.: Vm 24745/4014

For animal treatment only.



Approved: 13 October 2023