

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Cardboard Box}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Advantage 40 mg spot-on solution

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each 0.4 ml pipette contains: 40 mg imidacloprid.

**3. PACKAGE SIZE**

2 pipettes

3 pipettes

4 pipettes

6 pipettes

**4. TARGET SPECIES**



< 4 kg



< 4 kg

**5. INDICATIONS**

Flea (*Ctenocephalides felis*) prevention and treatment for cats.

Flea treatment for pet rabbits.

One treatment prevents further flea infestation for three to four weeks on cats and up to one week on pet rabbits.

The veterinary medicinal product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) in cats, where this has been previously diagnosed by a veterinary surgeon.

[Pictogram]



Flea

## **6. ROUTES OF ADMINISTRATION**

Spot-on use.

## **7. WITHDRAWAL PERIODS**

Do not use on rabbits intended for human consumption.

## **8. EXPIRY DATE**

Exp. {mm/yyyy}

## **9. SPECIAL STORAGE PRECAUTIONS**

Keep the blister in the outer carton.

## **10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.


## **11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

## **13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Elanco 

## **14. MARKETING AUTHORISATION NUMBER**

Vm 00879/4099

## **15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING  
UNITS {Pipette label}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Advantage



< 4kg



< 4kg

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

0.4 ml

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING**  
**UNITS {Blister}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Advantage



< 4kg



< 4kg

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

40 mg imidacloprid

0.4 ml

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

**PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

**PACKAGE LEAFLET**

**1. Name of the veterinary medicinal product**

Advantage 40 mg feline and bunny spot-on solution (< 4 kg)

Advantage 80 mg feline and bunny spot-on solution (≥ 4 kg)

**2. Composition**

Each unit dose (pipette) contains:

|  | <b>Unit Dose</b> | <b>Imidacloprid</b> |
|--|------------------|---------------------|
| Advantage 40 mg feline and bunny spot-on solution (< 4 kg) | 0.4 ml           | 40 mg               |
| Advantage 80 mg feline and bunny spot-on solution (≥ 4 kg) | 0.8 ml           | 80 mg               |

**Excipients:**

832.0 mg/ml benzyl alcohol (E1519), 1.0 mg/ml butylhydroxytoluene (E321)

Clear yellow to slightly brownish solution

**3. Target species**

Cats



and pet rabbits



**4. Indications for use**

For the prevention and treatment of flea (*Ctenocephalides felis*) infestations on cats and for the treatment of flea infestations on pet rabbits.

Fleas are killed within one day following treatment. One treatment prevents further flea infestation for three to four weeks on cats and up to one week on pet rabbits.

The veterinary medicinal product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) in cats, where this has been previously diagnosed by a veterinary surgeon.

## **5. Contraindications**

Do not treat unweaned kittens of less than 8 weeks of age.

Do not use on pet rabbits of less than 10 weeks of age.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

## **6. Special warnings**

### Special warnings:

Unnecessary use of antiparasitics or use deviating from the instructions given in the package leaflet may increase the resistance selection pressure and lead to reduced efficacy. The decision to use the veterinary medicinal product should be based on confirmation of the parasitic species and burden, or of the risk of infestation based on its epidemiological features, for each individual animal.

The possibility that other animals in the same household can be a source of re-infection with fleas should be considered, and these should be treated as necessary with an appropriate product.

Re-infestation from emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. More than one treatment may therefore be required, depending on the level of fleas in the environment. To aid reduction in environmental challenge, the additional use of a suitable environmental treatment against adult fleas and their developing stages is recommended, especially of the animal's basket, bedding and regular resting areas such as carpets and soft furnishings. In order to reduce further the environmental challenge, it is recommended that all cats and rabbits in the household are treated. Treatment of nursing queens controls flea infestations on both dam and offspring.

The veterinary medicinal product remains effective if the animal becomes wet, for example after exposure to heavy rain. However, re-treatment may become necessary, depending on the presence of fleas in the environment. In these cases do not treat more frequently than once weekly.

### Special precautions for safe use in the target species:

This veterinary medicinal product is for topical use and should not be administered orally.

Care should be taken to avoid the contents of the pipette coming into contact with the eyes or mouth of the recipient animal.

Do not allow recently treated animals to groom each other.

Any collar should be removed prior to application of the veterinary medicinal product. Prior to re-fitting the collar, the treated area should be visually assessed to ensure it is dry.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product contains benzyl alcohol and may cause skin sensitisation or transient skin reactions in rare cases (for example, irritation, tingling). People with known hypersensitivity to imidacloprid and benzyl alcohol should avoid contact with the veterinary medicinal product.

Avoid contact between the veterinary medicinal product and skin, eyes or mouth.

Do not massage the application site.

Do not eat, drink or smoke during application.

Wash off any skin contamination with soap and water.

If the veterinary medicinal product gets into eyes accidentally, the eyes should be thoroughly flushed with water.

If skin or eye irritation persists, or the veterinary medicinal product is accidentally swallowed, seek medical advice immediately and show the package leaflet or the label to the physician.

After application, do not stroke or groom animals until application site is dry.

Wash hands thoroughly after use.

Special precautions for the protection of the environment:

Imidacloprid is toxic to aquatic organisms. See also section 'Special precautions for disposal'.

Other precautions:

The solvent in this veterinary medicinal product may stain certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Pregnancy and lactation:

No reproductive toxic effects have been observed in rats and no primary embryotoxic or

teratogenic toxic effects have been observed during the studies on rats and rabbits.

Studies on pregnant and lactating queens together with their offspring are limited.

Evidence so far suggests that no adverse effects are to be expected in these animals.

Interaction with other medicinal products and other forms of interaction:

No incompatibility has been observed between this veterinary medicinal product at twice the recommended dose and the following commonly used veterinary medicinal products: lufenuron, pyrantel and praziquantel (cats). The compatibility of the veterinary medicinal product was also demonstrated with a wide range of routine treatments under field conditions including vaccination.

## Overdose:

In cats, no adverse clinical signs were produced using doses of five times the therapeutic level weekly for eight consecutive weeks.

In rabbits, no adverse clinical signs were seen using doses of up to 45 mg/kg body weight (4 times the therapeutic level) weekly for 4 consecutive weeks.

In rare cases of overdose or licking of treated fur, nervous system disorders (such as twitching, tremors, ataxia, mydriasis, miosis, lethargy) can occur in cats.

Poisoning following inadvertent oral uptake in animals is unlikely. In this event, treatment should be symptomatic under veterinary medical attention. There is no known specific antidote but administration of activated charcoal may be beneficial.

## **7. Adverse events**

Cats.

Very rare (<1 animal / 10,000 animals treated, including isolated reports):

|   |
|---|
| Agitation (hyperactivity)<br>Diarrhoea <sup>1</sup> , Hypersalivation (drooling) <sup>2</sup> , Vomiting <sup>1</sup><br>Neurological signs (e.g. Depression (lethargy), Incoordination, Tremor)<br>Application site reaction (e.g. Hair loss, Itching, Reddening of the skin, Skin lesion) |
|---|

<sup>1</sup>May occur after oral ingestion.

<sup>2</sup>May occur if the cat licks the application site immediately after treatment due to the bitter taste. This is not a sign of intoxication and disappears within some minutes without treatment.

Rabbits:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):

|  |
|--|
| Agitation (hyperactivity)<br>Diarrhoea <sup>1</sup> , Hypersalivation (drooling) <sup>2</sup><br>Application site reaction (e.g. Hair loss, Itching, Reddening of the skin, Skin lesion) |
|--|

<sup>1</sup>May occur after oral ingestion.

<sup>2</sup>May occur if the rabbit licks the application site immediately after treatment due to the bitter taste. This is not a sign of intoxication and disappears within some minutes without treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder <or its local representative > using the contact details at the end of this leaflet, or via your national reporting system:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: [adverse.events@vmd.gov.uk](mailto:adverse.events@vmd.gov.uk)

## 8. Dosage for each species, routes and method of administration

Spot-on use. For external use only.

Underdosing could result in ineffective use and may favour resistance development.

To ensure a correct dosage, body weight should be determined as accurately as possible.

Any instructions given by a veterinary surgeon for the use of this veterinary medicinal product should be followed.

### *Dosage and treatment schedule*

The recommended minimum dose is 10 mg imidacloprid per kg body weight (bw), equivalent to 0.1 ml/kg bw of the veterinary medicinal product.

| Cat/Pet Rabbit weight (kg) | Pipette size to be used                           | Pipette volume (ml) | Imidacloprid (mg/kg bw) |
|----------------------------|---|---------------------|-------------------------|
| < 4 kg                     | Advantage 40 mg feline and bunny spot-on solution | 0.4                 | minimum of 10           |
| ≥ 4 kg                     | Advantage 80 mg feline and bunny spot-on solution | 0.8                 | minimum of 10           |

Select the appropriate pipette size for the weight of the animal to be treated.

For treatment or prevention of infestations with fleas the need for and frequency of re-treatment(s) should be based on professional advice and should take into account the local epidemiological situation and the animal's lifestyle.

## 9. Advice on correct administration

Remove one pipette from the package. Hold pipette in an upright position, twist and pull off cap. Use reversed cap to twist and remove seal from pipette as shown in figure 1.

### Administration to the cat

Part the hair on the cat's neck at the base of the skull until the skin is visible. Place the tip of the pipette on the skin and squeeze firmly several times to empty the contents directly onto the skin as shown in figure 2. Application at the base of the skull will minimise the opportunity for the cat to lick the veterinary medicinal product. Apply only to undamaged skin.

## Administration to the rabbit

Part the hair on the rabbit's neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze firmly several times to empty the contents directly onto the skin as shown in figure 3. Application at the base of the skull will minimise the opportunity for the rabbit to lick the veterinary medicinal product. Apply only to undamaged skin.

### For monolingual packaging only:

Figures are shown below.

### For multilingual packaging only:

Figures are shown at the end of the leaflet.

*In order to facilitate multilingual package leaflets, figures were numbered, and this text added so that graphics can be printed only once at the end of the leaflet for multilingual leaflets.*




Figure 1




Figure 2




Figure 3

## **10. Withdrawal periods**

Do not use on rabbits intended for human consumption.

## **11. Special storage precautions**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Keep the blister in the outer carton.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and pipette after Exp. The expiry date refers to the last day of that month.

## **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

This veterinary medicinal product should not enter water courses as imidacloprid may be dangerous for fish and other aquatic organisms.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product not subject to prescription.

## **14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

| <b>Product Name</b>                               | <b>Vm Number</b> |
|---|------------------|
| Advantage 40 mg feline and bunny spot-on solution | 00879/4099       |
| Advantage 80 mg feline and bunny spot-on solution | 00879/4104       |

Cardboard box containing a total of 2, 3, 4 or 6 unit dose pipettes in a blister sheet. Each unit dose pipette contains 0.4 or 0.8 ml of solution. Not all pack sizes may be marketed.

## **15. PID LINK (Do not print heading)**

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

## **16. Contact details**

### Marketing authorisation holder and contact details to report suspected adverse events:

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

+44 3308221732  
PV.GBR@elancoah.com

### Manufacturer responsible for batch release:

KVP Pharma + Veterinär Produkte GmbH  
Projensdorfer Str. 324, 24106 Kiel  
Germany

### Local representatives <and contact details to report suspected adverse events>:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

## **17. Other information**

In further studies, in addition to the adulticide flea efficacy of imidacloprid, a larvicidal flea efficacy in the surroundings of the treated pet has been demonstrated. Larval stages in the pet's surroundings are killed following contact with a treated animal.

|             |
|-------------|
| NFA-<br>VPS |
|-------------|

*Gavin Hall*

Approved: 13 June 2025