

**<PARTICULARS TO APPEAR ON THE OUTER PACKAGE>**

Box with vial(s) of 100 ml/250 ml

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

DINALGEN 150 mg/ml solution for injection for cattle, pigs and horses

Ketoprofen

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Ketoprofen	150 mg
Benzyl alcohol (E1519)	10 mg

**3. PHARMACEUTICAL FORM**

Solution for injection.

**4. PACKAGE SIZE**

1 x 100 ml  
5 x 100 ml  
10 x 100 ml  
1 x 250 ml  
5 x 250 ml

**5. TARGET SPECIES**

Cattle, pigs and horses

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

**Intramuscular or intravenous use.**  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

Cattle:  
Meat and offal: 2 days  
Milk: zero hours

Horses:  
Meat and offal: 1 day

Milk: Not authorised for use in mares producing milk for human consumption

Pigs:  
and offal: 3 days

#### **9. SPECIAL WARNING(S), IF NECESSARY**

Avoid accidental self injection. **Read the package leaflet before use.**

#### **10. EXPIRY DATE**

<EXP {month/year}>

Shelf-life after opening the container for the first time: 28 days.

Once broached, use by: ...

#### **11. SPECIAL STORAGE CONDITIONS**

Keep the vial in the outer carton.

#### **12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

#### **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription

#### **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children

#### **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Ecuphar Veterinaria S.L.U.  
C/Cerdanya, 10-12 Planta 6º  
08173 Sant Cugat del Vallés  
Barcelona (Spain)

Distributed by:  
Duggan Veterinary Supplies Ltd,  
Holycross, Thurles,  
County Tipperary, E41 A093,  
Ireland  
Tel: +353 504 43169  
Email: sales@dugganvet.ie

**16. MARKETING AUTHORISATION NUMBER**

Vm 46037/4003

**17. MANUFACTURER'S BATCH NUMBER**

<Batch> <Lot> <BN> {number}

**<PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE>**

Vials containing 100 ml/250 ml

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

DINALGEN 150 mg/ml solution for injection for cattle, pigs and horses  
Ketoprofen

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:  
Ketoprofen 150 mg  
Benzyl alcohol (E1519) 10 mg

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE**

100 ml  
250 ml

**5. TARGET SPECIES**

Cattle, pigs and horses

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intramuscular or intravenous use.  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

Cattle:  
Meat and offal: 2 days  
Milk: zero hours

Horses:  
Meat and offal: 1 day  
Milk: Not authorised for use in mares producing milk for human consumption

Pigs:  
Meat and offal: 3 days

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

<EXP {month/year}>

Shelf-life after first opening the immediate packaging: 28 days

Once broached, use by .....

**11. SPECIAL STORAGE CONDITIONS**

Keep the vial in the outer carton.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Ecuphar Veterinaria S.L.U.  
C/Cerdanya, 10-12 Planta 6º  
08173 Sant Cugat del Vallés  
Barcelona (Spain)

Distributed by:

Duggan Veterinary Supplies Ltd,  
Holycross, Thurles,  
County Tipperary, E41 A093,  
Ireland  
Tel: +353 504 43169  
Email: [sales@dugganvet.ie](mailto:sales@dugganvet.ie)

**16. MARKETING AUTHORISATION NUMBER**

Vm 46037/4003

**17. MANUFACTURER’S BATCH NUMBER**

<Batch> <Lot> <BN> {number}

## PACKAGE LEAFLET

DINALGEN 150 mg/ml solution for injection for cattle, pigs and horses

### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Ecuphar Veterinaria S.L.U.  
C/Cerdanya, 10-12 Planta 6º  
08173 Sant Cugat del Vallés  
Barcelona (Spain)

Manufacturer for the batch release:

Zoetis Manufacturing & Research Spain, S.L.  
Crt. Camprodón s/n, 17813 Vall de Bianya (Girona)  
Spain

Distributed by:

Duggan Veterinary Supplies Ltd,  
Holycross, Thurles,  
County Tipperary, E41 A093,  
Ireland  
Tel: +353 504 43169  
Email: sales@dugganvet.ie

### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

DINALGEN 150 mg/ml solution for injection for cattle, pigs and horses

Ketoprofen

### 3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each ml contains:

<b>Active substance:</b>	
Ketoprofen	150 mg
<b>Excipients:</b>	
Benzyl alcohol (E1519)	10 mg

A clear colourless to yellowish solution.

### 4. INDICATIONS

Cattle:

- Reduction of inflammation and pain associated with post-partum, musculoskeletal disorders and lameness
- Reduction of fever associated with bovine respiratory disease

- Reduction of inflammation, fever and pain in acute clinical mastitis in combination with antimicrobial therapy where appropriate.

Pigs:

- Reduction of pyrexia in cases of respiratory disease and Postpartum

Dysgalactia

Syndrome-PDS- (Metritis Mastitis Agalactia syndrome) in sows, in combination with antimicrobial therapy, where appropriate.

Horses:

- Reduction of inflammation and pain associated with osteoarticular and musculoskeletal disorders (lameness, laminitis, osteoarthritis, synovitis, tendinitis, etc.).
- Reduction of postoperative pain and inflammation.
- Reduction of visceral pain associated with colic.

## **5. CONTRAINDICATIONS**

Do not use in animals where there is the possibility of gastro-intestinal ulceration or bleeding, in order not to aggravate their situation.

Do not use in animals suffering from cardiac, hepatic, or renal disease.

Do not use in cases of known hypersensitivity to ketoprofen or acetyl-salicylic acid or to any of the excipients.

Do not use in animals with evidence of blood dyscrasia or coagulopathy.

Do not administer other non-steroidal anti-inflammatory drugs (NSAIDs) concurrently or within 24 hours of each other.

## **6. ADVERSE REACTIONS**

Intramuscular injection of ketoprofen can cause mild, transient, necrotic subclinical muscular lesions that gradually resolve in the days after completion of treatment. Administration in the neck region minimizes the extension and severity of these lesions.

In horses, transient local reactions, which disappeared after 5 days, were observed after one administration of the product at the recommended volume by extravascular route.

Due to the mechanism of action of ketoprofen, after repeated administrations, erosive and ulcerative lesions of gastrointestinal tract may occur.

In common with all NSAIDs due to their action of inhibition of prostaglandins' synthesis, there can be the possibility in certain individuals of gastric or renal intolerance.

If side effects occur treatment must be stopped and the advice of a veterinarian should be sought.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Cattle, pigs and horse

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Intramuscular or intravenous use.

Cattle:

3 mg ketoprofen/kg body weight, i.e. 1ml of product per 50 kg body weight/ day, administered via the intravenous or intramuscular route, preferably in the neck region.

The duration of treatment is 1-3 days, and should be established according to the severity and duration of symptoms.

Pigs:

3 mg of ketoprofen/kg body weight i.e. 1 ml of the product per 50 kg body weight/ day, administered via the intramuscular route on a single occasion.

Depending on the response observed and based on the benefit-risk analysis by the responsible veterinarian treatment may be repeated at intervals of 24 hours for a maximum of three treatments. Each injection should be given at a different site.

Horses:

2.2 mg of ketoprofen/kg body weight, i.e. 0.75 ml of the product. per 50 kg body weight/ day, administered via the intravenous route.

The duration of treatment is 1-5 days, and should be established according to the severity and duration of symptoms.

In the case of colic one injection is normally sufficient. A second administration of ketoprofen requires a clinical re-examination.

## **9. ADVICE ON CORRECT ADMINISTRATION**

## **10. WITHDRAWAL PERIOD**

Cattle:

Meat and offal: 2 days

Milk: zero hours

Horses:

Meat and offal: 1 day

Milk: Not authorised for use in mares producing milk for human consumption

Pigs:

Meat and offal: 3 days

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Keep the vial in the outer carton.

Shelf-life after opening the container for the first time: 28 days.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of the month.

## 12. SPECIAL WARNING(S)

### Special precautions for use in animals:

Do not exceed the recommended dose. Do not exceed the recommended treatment period.

The use of ketoprofen is not recommended in foals less than one month of age. When administering to animals of less than 6 weeks of age, ponies or in aged animals it is necessary to adjust the dose accurately as well as to perform a close clinical follow-up.

Avoid intra-arterial injection.

Avoid use in any dehydrated, hypovolaemic or hypotensive animals as there is a potential risk of increased renal toxicity.

Since gastric ulceration is a common finding in PMWS (Post-weaning Multisystemic Wasting Syndrome), the use of ketoprofen in pigs affected by this pathology is not recommended, in order not to aggravate their situation.

In horses, avoid extravascular administration.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Avoid contact with the skin, eyes and mucous membranes.

In case of accidental skin, eye or mucous membrane contact, wash the affected area thoroughly with clean running water immediately. Seek medical advice if irritation persists.

Avoid accidental self injection. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

Hypersensitivity reactions (skin rash, urticaria) could occur. People with known hypersensitivity to the active substance or to any of the excipients should avoid contact with the veterinary medicinal product.

### Use during pregnancy and lactation:

Studies in laboratory animals (rats, mice, rabbits) and cattle have not produced any evidence of adverse effects. Can be used in pregnant cows.

The safety of the veterinary medicinal product has not been established during pregnancy in sows and mares. Use only accordingly to the benefit-risk assessment by the responsible veterinarian.

Can be used in lactating cows and sows.

The use is not recommended in lactating mares.

### Interactions with other medicinal products and other forms of interaction:

- Concurrent administration of diuretics or potentially nephrotoxic drugs should be avoided since there is an increase of renal disturbances, including renal

failure. This is secondary to the diminished blood flow caused by the inhibition of prostaglandins synthesis.

- Do not administer with other non steroidal anti-inflammatory drugs (NSAIDs), corticosteroids, anticoagulants or diuretics concurrently or within 24 hours of administration of the product since the risk of gastrointestinal ulceration and other adverse reactions may be exacerbated.

- The treatment free period should however take into account the pharmacological properties of the products used previously.

- Ketoprofen is highly bound to plasma proteins and may compete with other highly bound drugs which can lead to toxic effects.

Overdose (symptoms, emergency procedures, antidotes):

Overdose with non-steroidal anti-inflammatory drugs can lead to gastrointestinal ulceration, loss of proteins, hepatic and renal impairment.

In tolerance studies performed in pigs, up to 25% of the animals treated at three times the maximum recommended dose (9 mg/kg bw) for three days or at the recommended dose (3 mg/kg bw) for triple the maximum recommended time (9 days) showed erosive and/or ulcerative lesions in both the aglandular (pars oesophagica) and glandular parts of the stomach. Early signs of toxicity include loss of appetite and pasty faeces or diarrhoea.

The intramuscular administration of the product to cattle, at up to 3 times the recommended dose or for 3 times the recommended duration of the treatment (9 days) did not result in clinical signs of intolerance. However, inflammation as well as necrotic subclinical lesions were detected at the injection site of the treated animals as well as an increase in CPK levels. The histopathological examination showed erosive or ulcerative abomasal lesions related to both dosage regimes.

Horses have been found to tolerate intravenous dosages of ketoprofen up to 5 times the recommended dose for three times the recommended duration (15 days) with no evidence of toxic effects.

If clinical signs of overdose are observed, there is no specific antidote, therefore symptomatic treatment should be initiated.

Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

### **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

April 2022

**15. OTHER INFORMATION**

- . Box with 1, 5 or 10 vials of 100 ml.
- . Box with 1 or 5 vials of 250 ml.

Not all pack sizes may be marketed.

For animal treatment only. To be supplied only on veterinary prescription.

Approved 05 April 2022

A handwritten signature in black ink, appearing to read "Hunter.", is written below the approval date. The signature is stylized and includes a vertical line to the left of the name.