

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Bravoxin suspension for injection for cattle and sheep

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each 1 ml contains:

#### **Active substances:**

<i>C. perfringens</i> type A ( $\alpha$ ) toxoid	$\geq 0.5$ IU <sup>#</sup>
<i>C. perfringens</i> type B & C ( $\beta$ ) toxoid	$\geq 18.2$ IU*
<i>C. perfringens</i> type D ( $\epsilon$ ) toxoid	$\geq 5.3$ IU*
<i>C. chauvoei</i> whole culture, inactivated	$\geq 90\%$ protection**
<i>C. novyi</i> toxoid	$\geq 3.8$ IU*
<i>C. septicum</i> toxoid	$\geq 4.6$ IU*
<i>C. tetani</i> toxoid	$\geq 4.9$ IU*
<i>C. sordellii</i> toxoid	$\geq 4.4$ U <sup>1</sup>
<i>C. haemolyticum</i> toxoid	$\geq 17.4$ U <sup>#</sup>

\* ELISA According to Ph.Eur.

<sup>1</sup> In house ELISA

\*\* Guinea pig challenge test according to Ph.Eur.

# *In vitro* toxin neutralisation test based on haemolysis of sheep erythrocytes.

#### **Adjuvant:**

Aluminium<sup>1</sup> 3.026 – 4.094 mg

<sup>1</sup> from aluminium potassium sulphate (alum)

#### **Excipients:**

Thiomersal 0.05 – 0.18 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Light brown aqueous suspension that settles on storage.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cattle and sheep.

## 4.2 Indications for use, specifying the target species

For the active immunisation of sheep and cattle against disease associated with infections caused by *Clostridium perfringens* type A, *C. perfringens* type B, *C. perfringens* type C, *C. perfringens* type D, *Clostridium chauvoei*, *Clostridium novyi* type B, *Clostridium septicum*, *Clostridium sordellii* and *Clostridium haemolyticum* and against tetanus caused by *Clostridium tetani*.

For the passive immunisation of lambs and calves against infections caused by the above mentioned clostridial species (except *C. haemolyticum* in sheep).

### Onset of immunity:

Sheep and Cattle: 2 weeks after the basic vaccination course (as demonstrated by serology only).

### Duration of active immunity:

As demonstrated by serology only:

Sheep: 1 year against *C. perfringens* type A, B, C and D, *C. novyi* type B, *C. sordellii*, *C. tetani*;

< 6 months against *C. septicum*, *C. haemolyticum*, *C. chauvoei*;

Cattle: 1 year against *C. tetani* and *C. perfringens* type D;

< 1 year against *C. perfringens* type A, B and C;

< 6 months against *C. novyi* type B, *C. septicum*, *C. sordellii*, *C. haemolyticum*, *C. chauvoei*.

An anamnestic humoral immune response (immunological memory) to all components was demonstrated 1 year following the basic course of vaccination.

### Duration of passive immunity:

As demonstrated by serology only:

Lambs: At least 2 weeks for *C. septicum* and *C. chauvoei*;

At least 8 weeks for *C. perfringens* type B and *C. perfringens* type C;

At least 12 weeks for *C. perfringens* type A, *C. perfringens* type D, *C. novyi* type B, *C. tetani* and *C. sordellii*;

No passive immunity was observed for *C. haemolyticum*.

Calves: At least 2 weeks for *C. sordellii* and *C. haemolyticum*;

At least 8 weeks for *C. septicum* and *C. chauvoei*;

At least 12 weeks for *C. perfringens* type A, *C. perfringens* type B, *C. perfringens* type C, *C. perfringens* type D, *C. novyi* type B, and *C. tetani*.

## 4.3 Contraindications

Do not use in sick or immunodeficient animals.

## 4.4 Special warnings for each target species

Vaccinate healthy animals only.

The effectiveness of the vaccine in providing passive immunity to young lambs and calves depends on these animals ingesting adequate amounts of colostrum on the first day of life.

Clinical trials have demonstrated that the presence of maternal derived antibodies (MDA), particularly against *C. tetani*, *C. novyi* type B, *C. perfringens* type A (calves only), *C. chauvoei* (lambs only) and *C. perfringens* type D may reduce the antibody response to vaccination in young lambs and calves. Therefore, to ensure an optimal response in young animals with high levels of MDA, the basic vaccination should be delayed until the levels wane (which is after about 8-12 weeks of age, see section 4.2).

#### 4.5 Special precautions for use

##### Special precautions for use in animals

It is good practice to observe animals regularly for adverse reactions at the injection site following vaccination. It is recommended to seek medical advice from a veterinarian in case of a severe injection site reaction.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

##### Special precautions for the protection of the environment

Not applicable.

##### Other precautions

Not applicable.

#### 4.6 Adverse reactions (frequency and seriousness)

Cattle and sheep:

Very common (>1 animal / 10 animals treated):	Injection site swelling <sup>1</sup> .
Common (1 to 10 animals / 100 animals treated):	Injection site abscess, injection site skin discolouration <sup>2</sup> . Hyperthermia <sup>3</sup> .
Uncommon (1 to 10 animals / 1,000 animals treated):	Injection site pain <sup>4</sup> .
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Anaphylactic type reaction <sup>5</sup> .

<sup>1</sup> Up to a mean value of 6 cm in sheep and 15 cm diameter in cattle; sometimes reactions of up to 25 cm diameter may be seen in cattle. Most local reactions resolve within 3-6 weeks in sheep and in less than 10 weeks in cattle. In a minority of animals they may persist longer.

<sup>2</sup> Returns to normal as the local reaction resolves.

<sup>3</sup> Mild.

<sup>4</sup> For 1-2 days post first vaccination.

<sup>5</sup> In such cases appropriate treatment such as adrenaline should be administered without delay.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

#### **4.7 Use during pregnancy, lactation or lay**

##### Pregnancy:

No side effects other than those described under section 4.6 were seen when the vaccine was used in sheep and cattle between 8 and 2 weeks prior to parturition. In the absence of specific data, the use of the vaccine is not recommended during the first or second third of pregnancy.

Avoid stress in pregnant ewes and cows.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

#### **4.9 Amount(s) to be administered and administration route**

Subcutaneous use.

##### *Dose:*

- Sheep: 1 ml – from 2 weeks of age
- Cattle: 2 ml – from 2 weeks of age

##### *Administration:*

By subcutaneous injection preferably in the loose skin on the side of the neck, observing aseptic precautions.

Shake the bottle thoroughly before use.

Syringes and needles should be sterile before use and the injection should be made through an area of clean, dry skin taking precautions against contamination.

*Basic vaccination:* Two doses should be administered, 4-6 weeks apart (see section 4.2 and 4.4).

*Re-vaccination:* A single dose should be administered at 6 to 12 month intervals after the basic vaccination (see section 4.2.)

##### *Use in pregnancy:*

To provide passive protection of the offspring, via the colostrum, a single re-vaccination should be administered between 8 and 2 weeks before parturition, provided that animals have received a full basic vaccination course before pregnancy.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In calves and lambs, local reactions may increase slightly if twice the recommended dose is administered (see section 4.6).

#### **4.11 Withdrawal period(s)**

Zero days.

### **5. IMMUNOLOGICAL PROPERTIES**

**Pharmacotherapeutic group:** Immunologicals for *Bovidae* and *Ovidae*, inactivated bacterial vaccines (including mycoplasma, toxoid and chlamydia) for cattle and sheep, clostridium.

**ATCvet code:** QI02AB01, QI04AB01.

Inactivated clostridium vaccine. To stimulate active immunity in sheep and cattle against *C. chauvoei* and the toxins of *Clostridium perfringens* type A, *C. perfringens* type B, *C. perfringens* type C, *C. perfringens* type D, *C. novyi*, *C. septicum*, *C. tetani*, *C. sordellii*, and *C. haemolyticum* contained in the vaccine.

To provide passive immunity via the colostrum against the above mentioned clostridial infections in young lambs and calves.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Aluminium potassium sulphate (alum)  
Thiomersal  
Sodium chloride  
Water for injections  
Formaldehyde

#### **6.2 Major incompatibilities**

Do not mix with any other veterinary medicinal product.

#### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 30 months.  
Shelf life after first opening the immediate packaging: 8 hours.

#### **6.4 Special precautions for storage**

Store and transport refrigerated (2 °C – 8 °C). Do not freeze.  
Protect from light.

## **6.5 Nature and composition of immediate packaging**

Flexible low density polyethylene (LDPE) bottle with 50 ml or 100 ml, closed with a bromobutyl rubber stopper and held in place with an aluminium cap.

### Pack sizes:

Cardboard box with one bottle of 50 ml (50 doses of 1 ml or 25 doses of 2 ml).

Cardboard box with one bottle of 100 ml (100 doses of 1 ml or 50 doses of 2 ml).

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Limited  
Walton Manor  
Walton  
Milton Keynes  
MK7 7AJ

## **8. MARKETING AUTHORISATION NUMBER**

Vm 01708/5094

## **9. DATE OF FIRST AUTHORISATION**

05 March 2021

## **10. DATE OF REVISION OF THE TEXT**

April 2024


## **PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

## 11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk) in this section.

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 08 May 2024