

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Chloromed 150 mg/g oral powder for calves.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Active substance:

Each g contains:
chlortetracycline hydrochloride 150 mg

Excipients:

Qualitative composition of excipients and other constituents
Medium Chain Triglycerides.
Glucose Monohydrate.
Colloidal anhydrous silica.

Yellow uniform powder.

3. CLINICAL INFORMATION

3.1 Target species

Cattle (Calves less than 6 months of age).

3.2 Indications for use for each target species

The veterinary medicinal product is indicated in the treatment of respiratory disease in calves caused by *Pasteurella* spp., sensitive to chlortetracycline.

3.3 Contraindications

Do not use in calves over 6 months of age and dairy cows.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals with severe liver and renal disorders.

3.4 Special warnings

The uptake of oral medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed, animals should be treated parenterally.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Use of the veterinary medicinal product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to chlortetracycline and may decrease the effectiveness of treatment with related substances, due to the potential for cross-resistance.

Long term use of this veterinary medicinal product is not recommended as it may lead to the development of bacterial resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to tetracyclines should avoid contact with the veterinary medicinal product.

Do not eat, drink or smoke while handling the veterinary medicinal product or medicated feed.

During preparation and administration of the medicated feed, skin contact with the veterinary medicinal product and inhalation of dust particles should be avoided.

Personal protective equipment consisting of protective overall, glasses, impermeable gloves (e.g. rubber or latex) and an appropriate dust mask (e.g. disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143) should be worn when handling the veterinary medicinal product.

Wash hands immediately after handling the veterinary medicinal product or medicated feed.

In the event of skin or eye contact, rinse immediately the affected area with large amounts of clean water. If irritation occurs, seek medical attention.

If you develop symptoms following exposure to the veterinary medicinal product such as skin rash, you should seek medical advice and show the package leaflet or the label to the physician. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

None.

3.6 Adverse events

Cattle (Calves less than 6 months of age):

Rare (1 to 10 animals / 10 000 animals treated):	Allergic reactions;
	Liver disorders;
	Renal disorders;
	Gastrointestinal disorders;
	Photosensitivity.

Due to the possible incorporation of chlortetracycline, treatment of pregnant and newborn animals can lead to dysfunctional development of the skeleton and teeth in foetal and growing animals.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the immediate packaging for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

Not applicable.

3.8 Interaction with other medicinal products and other forms of interaction

This veterinary medicinal product is not recommended for concurrent administration with any other oral medication.

Do not incorporate the veterinary medicinal product in feed overloaded with polyvalent cations such as Ca^{2+} and Fe^{3+} because the formation of chlortetracycline complexes with these cations is possible.

Do not administer together with antacids, kaolin and iron preparations and in conjunction with bactericidal antibiotics like beta-lactams.

The veterinary medicinal product should not be used in case of known resistance to other tetracyclines.

3.9 Administration routes and dosage

Oral use.

The recommended therapeutic dose is 20 mg chlortetracycline per kg bodyweight (equivalent to 20 grams per 150 kg bodyweight) per day administered for seven days. This should be given in a divided daily dose i.e. 10 g morning and 10 g evening.

The veterinary medicinal product should be administered to small quantities of feed for immediate consumption by individual animals. Larger groups should be treated with medicated feeding stuff.

The veterinary medicinal product should be mixed thoroughly into a part of the daily feed ration and should be administered prior to the feeding. It should be ensured that the calculated dose is completely taken up by the animals.

If animals don't recover within 3 days after oral medication, diagnosis should be reconsidered and treatment should be changed, if necessary.

To ensure a correct dosage, bodyweight should be determined as accurately as possible.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Do not exceed the stated dose.

Chlortetracycline toxicity is low. If digestive disturbances do occur, treatment should be discontinued.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Cattle (Calves):

Meat and offal: 10 days.

Milk: Not authorised for use in animals producing milk for human consumption.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code:

QJ01AA03

4.2 Pharmacodynamics

Chlortetracycline hydrochloride is a predominantly bacteriostatic antibiotic, interfering with bacterial protein synthesis of the rapidly growing and reproducing bacterial cell. Chlortetracycline has a broad spectrum of activity, including Gram-positive aerobes, Gram-negative anaerobes and Mycoplasmas. Resistance is known to occur in respiratory pathogens of cattle and cross-resistance occurs between chlortetracycline and other tetracyclines.

The Clinical and Laboratories Standards Institute (CLSI) breakpoints established for tetracyclines are as follows:

Organisms other than streptococci: S: $\leq 4\mu\text{g/ml}$, I: $8\mu\text{g/ml}$; R: $\geq 16\mu\text{g/ml}$.

4.3 Pharmacokinetics

Following oral administration of the recommended dose, maximum blood levels of approximately 1 – 2 µg/ml are achieved within 2 – 8 hours and approximately 37% of the oral dose is systemically available. Steady state plasma concentrations of chlortetracycline are maintained throughout the twice-daily seven day treatment period. Chlortetracycline accumulates in the lung tissue resulting in higher concentrations at the site of activity. Chlortetracycline undergoes little metabolism and is excreted through both the urinary and biliary systems.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.
Shelf-life after first opening the immediate packaging: 28 days.

5.3 Special precautions for storage

Do not store above 25°C.
Store in a dry place.
Store in the original container.
Protect from light.

5.4 Nature and composition of immediate packaging

1 kg, clear low density polyethylene bag laminated with metallised polyester.

5.5 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Univet Ltd

7. MARKETING AUTHORISATION NUMBER

Vm 05150/4001

8. DATE OF FIRST AUTHORISATION

29 October 2009

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

December 2025

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

Gavin Hall

Approved: 27 February 2026