

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

X-SPECTRA Flavoured tablets for medium and small dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains:

#### **Active substances:**

Febantel 150 mg  
Pyrantel (as embonate) 50 mg  
Praziquantel 50 mg

#### **Excipients**

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablet.

Yellow brown, oval tablet with a score line on both sides.

The tablets can be divided into equal halves.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs.

#### **4.2 Indications for use, specifying the target species**

Treatment of mixed infections by cestodes and nematodes of the following species:

##### Nematodes:

Ascarids: *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms) .

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

Whipworms: *Trichuris vulpis* (adults).

##### Cestodes:

Tapeworms: *Echinococcus* species, (*E. Granulosus*, *E. Multilocularis*), *Taenia* species (*T. hydatigena*, *T. pisiformis*, *T. taeniformis*), *Dipylidium caninum* (adult and immature forms).

#### **4.3 Contraindications**

Do not use in animals with a known sensitivity to the active ingredients or to any of the excipients.

See section 4.7 below.

#### **4.4 Special warnings for each target species**

Dogs kept together or in kennels should be treated at the same time.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation may reoccur unless control of intermediate hosts such as fleas, mice etc is undertaken.

#### **4.5 Special precautions for use**

##### **Special precautions for use in animals**

The product is not recommended for use in dogs of less than 3 kg bodyweight.

In debilitated or heavily infested animals, the product should be used only according to a benefit/risk assessment by the responsible veterinarian.

To minimise the risk of reinfestation and new infestation, excreta should be collected and properly disposed of for 24 hours following treatment.

##### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Wash hands after administration to the animal.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

People with known hypersensitivity to any of the ingredients should avoid contact with the veterinary medicinal product.

##### **Other precautions**

Since it contains praziquantel, the product is effective against *Echinococcus* spp. which do not occur in all EU member states but are becoming more common in some. Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

#### **4.6 Adverse reactions (frequency and seriousness)**

Gastro-intestinal signs (vomiting, diarrhoea), possibly associated with lethargy, have been observed very rarely in spontaneous reports.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

#### **4.7 Use during pregnancy, lactation or lay**

##### Pregnancy:

Do not use in pregnant bitches during the first 4 weeks of pregnancy.

##### Lactation:

The product may be used during lactation (see Section 4.9 below).

#### 4.8 Interaction with other medicinal products and other forms of interaction

Do not use simultaneously with piperazine, Levamisole as the anthelmintic effects of pyrantel may be antagonized.

Plasma concentrations of praziquantel may be decreased by concomitant administration with drugs that increase the activity of cytochrome P-450 enzymes (e.g. dexamethasone, phenobarbital).

Concurrent use with other cholinergic compounds can lead to toxicity.

#### 4.9 Amounts to be administered and administration route

Oral use.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 5 mg/kg pyrantel (as embonate) and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg bodyweight, in a single administration.

Dosages are as follows:

Body weight (kg)	Tablet quantity
3-5	1/2
>5-10	1
>10-15	1 1/2
>15-20	2

The tablets are flavoured and consequently taken by most dogs voluntarily.

The tablets can be given to the dog with or without food. No starvation is needed before or after treatment.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

If there is a risk for re-infestation, the advice of a veterinarian should be sought regarding the need for and the frequency of repeat administration. In case of confirmed single infestation by cestode or by nematode, a monovalent product containing a cestocide or a nematocide alone should be used.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

In safety studies, single doses of 5 times the recommended dose or greater gave rise to occasional vomiting.

#### 4.11 Withdrawal period(s)

Not applicable.

### 5. PHARMACOLOGICAL PROPERTIES

ATCvet code: QP52AA51.

Pharmacotherapeutic group: anthelmintics.

## 5.1 Pharmacodynamic properties

In this fixed combination pyrantel and febantel act against all relevant nematodes (ascarids, hookworms, and whipworms) in dogs. In particular the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*. This combination shows synergistic activity in the case of hookworms and febantel is effective against *T. vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular *Taenia* spp, *Dipylidium caninum*, *Echinococcus granulosus* and *Echinococcus multilocularis*. Praziquantel acts against all adult and immature forms of these parasites.

Praziquantel is very rapidly absorbed through the parasite's surface and distributed throughout the parasite. Both *in vitro* and *in vivo* studies have shown that praziquantel causes severe damage to the parasite integument, resulting in the contraction and paralysis of the parasites. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis of the nematodes and thereby allow removal from the gastro- intestinal (GI) system by peristalsis.

Within the mammalian system febantel undergoes ring closure forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption of structures vital to the normal functioning of the helminth.

Glucose uptake, in particular is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2-3 days later.

## 5.2 Pharmacokinetic particulars

After oral administration to dogs, praziquantel is extensively and quickly absorbed from the gastro-intestinal tract. Maximum plasma concentration of 752 µg/L is obtained in less than 2 hours. It is rapidly and extensively metabolised in the liver into hydroxylated derivatives of the parent compound, then rapidly eliminated, mainly in urine.

After oral administration to dogs, febantel is moderately absorbed from the gastro-intestinal tract. Febantel is rapidly metabolised in the liver into fenbendazole and its hydroxy and oxidative derivatives like oxfendazole. Maximum plasma concentration of fenbendazole (173 µg/L) is obtained after about 5 hours. Maximum plasma concentration of oxfendazole (147 µg/L) is obtained after about 7 hours. The excretion occurs mainly in the faeces.

After oral administration to dogs, pyrantel embonate is poorly absorbed. Maximum plasma concentration of 79 µg/L is obtained after about 2 hours. It is rapidly and extensively metabolised in the liver, then rapidly excreted, mainly in the faeces (the unchanged form) and in urine (the metabolites).

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Liver powder flavour  
Tablet grade inactive yeast  
Sodium laurilsulfate  
Croscarmellose sodium  
Povidone K30  
Anhydrous colloidal silica  
Cellulose microcrystalline  
Magnesium stearate  
Maize starch

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.  
Half-tablets should be discarded.

### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

Nature of immediate packaging:  
Polyamide-aluminium-PVC / aluminium blister packs.

Pack sizes:

Box containing 1 blister of 2 tablets  
Box containing 2 blisters of 2 tablets  
Box containing 1 blister of 8 tablets  
Box containing 13 blisters of 8 tablets  
Box containing 52 blisters of 2 tablets

Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Ceva Animal Health Ltd  
Explorer House  
Mercury Park  
Wycombe Lane  
Wooburn Green  
High Wycombe  
Buckinghamshire  
HP10 0HH  
United Kingdom

**8. MARKETING AUTHORISATION NUMBER**

Vm 15052/4055

**9. DATE OF FIRST AUTHORISATION**

26 November 2010

**10. DATE OF REVISION OF THE TEXT**

October 2022

Approved 20 October 2022

