

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

FLIMABO 100 mg/g suspension for use in drinking water for chickens and pigs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each g contains:

Active substance:

Flubendazole 100 mg

Excipients:

Methyl parahydroxybenzoate (E218) 2.0 mg

Sodium benzoate (E211) 5.0 mg

Disodium edetate 0.1 mg

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Suspension for use in drinking water.
White to brownish white suspension.

4. CLINICAL PARTICULARS

4.1 Target species

Pigs (piglets, pigs for fattening, pregnant and lactating sows) and chickens (layer hens, chickens for reproduction, pullets, broilers).

4.2 Indications for use, specifying the target species

In hens/chickens:

Treatment of helminthiasis caused by *Ascaridia galli* (adult stages), *Heterakis gallinarum* (adult stages), *Capillaria* spp. (adult stages).

In pigs:

Treatment of helminthiasis caused by *Ascaris suum* (adult and intestinal larval stages) in piglets, pigs for fattening, pregnant and lactating sows.

4.3 Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings for each target species

In chickens, optimal results can only be achieved if strict rules of hygiene are respected in the maintenance of the cages.

In both species:

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of body weight, misadministration of the veterinary medicinal product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

4.5 Special precautions for use

Special precautions for use in animals

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Direct contact with veterinary medicinal product should be avoided. Personal protective equipment consisting of protective gloves should be worn when handling the veterinary medicinal product. Wash hands after use.

People with known hypersensitivity to flubendazole should avoid contact with the veterinary medicinal product. In the event of eye contact, rinse thoroughly with water. In case of appearance and persistence of conjunctival redness, seek medical advice and show the package leaflet to the physician.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

Not applicable.

4.6 Adverse reactions (frequency and seriousness)

Pigs:

None known.

Chickens:

Undetermined frequency (cannot be estimated from the available data):	Development disorders of the feathers
---	---------------------------------------

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

4.7 Use during pregnancy, lactation or lay

Laboratory studies in rabbits and rats have not produced any evidence of embryotoxicity, teratogenicity at therapeutic doses. High dosages gave equivocal results. In laboratory studies in rats, there were no effects on pups during lactation. The safety of the product has been demonstrated in laying hens, pregnant and lactating sows.

Pregnancy and lactation:

Can be used in pregnant and lactating sows.

Laying birds:

Can be used in laying hens.

4.8 Interaction with other medicinal products and other forms of interaction

None known.

4.9 Amounts to be administered and administration route

Oral use.

Hens/chickens:

1.43 mg flubendazole (= 14.3 mg veterinary medicinal product) per kg body weight daily during 7 days i.e. 1 g of the veterinary medicinal product per 70 kg body weight daily for 7 days.

Pigs:

a) Treatment of helminthiasis caused by *Ascaris suum* (adult stages and intestinal larval stages):

1 mg flubendazole (= 10 mg veterinary medicinal product) per kg body weight daily during 5 days, i.e. 1 g of the veterinary medicinal product per 100 kg body weight daily for 5 days;

b) Treatment of helminthiasis caused by *Ascaris suum* (adult stages):

2.5 mg flubendazole (= 25 mg veterinary medicinal product) per kg body weight daily during 2 days, i.e. 2.5 g of the veterinary medicinal product per 100 kg body weight daily for 2 days.

Pigs should be grouped according to their bodyweight and dosed accordingly, in order to prevent under or overdosing.

Based on the recommended dose and the number and weight of animals to be treated, the exact daily concentration of the veterinary medicinal product should be calculated according to the following formula:

$$\frac{\text{mg veterinary medicinal product/kg body weight/day} \times \text{average body weight (kg) of animals to be treated}}{\text{average quantity of drinking water (l/animal) consumed in 4 h}} = \text{mg veterinary medicinal product per litre of drinking water}$$

This will result in a concentration of flubendazole between 20 and 200 mg per litre.

Method of administration:

Administration in drinking water

- 1) The required quantity of the veterinary medicinal product is in function of the estimated body weight of the total group animals (see table below for guidance).

Hens/chickens, 7 days of treatment

Total weight of chickens	Amount of medication to be used (g/ day)	Total amount of medication used (g/ 7 days)
1400 kg	20 g	7 x 20 g
3500 kg	50 g	7 x 50 g
7000 kg	100 g	7 x 100 g
52500 kg	750 g	7 x 750 g

Pigs, 5 days of treatment

Total weight of pigs	Amount of medication to be used (g/ day)	Total amount of medication used (g/ 5 days)
2000 kg	20 g	5 x 20 g
5000 kg	50 g	5 x 50 g
10000 kg	100 g	5 x 100 g
75000 kg	750 g	5 x 750 g

Pigs, 2 days of treatment

Total weight of pigs	Amount of medication to be used (g/ day)	Total amount of medication used (g/ 2 days)
800 kg	20 g	2 x 20 g
2000 kg	50 g	2 x 50 g
4000 kg	100 g	2 x 100 g
30000 kg	750 g	2 x 750 g

- 2) Each day a predilution is prepared containing the daily required dose of the veterinary medicinal product admixed in 10 to 100 times its weight in water depending on the distribution system. For example: for 500 g of the veterinary medicinal product, add 5 litres to 50 litres of water.
- 3) If less than entire package (a sachet or a container) is required, the required dose should be measured by suitably calibrated weighing equipment.

- 4) If the entire sachet is used, squeeze it gently before use and then empty the contents into the predilution recipient.
- 5) Stir the predilution vigorously with a manual mixer (whisk) for 2 minutes to obtain a white milky homogenous mixture.
- 6) This predilution must be distributed via the general water supply system:
Tanks: add the predilution to the quantity of water usually consumed by the animals over a period of up to 4 hours.
Dosing pumps: adjust the flow rate of the pump to distribute the predilution over a period of up to 4 hours.

In order to ensure administration of the correct dose, a substantial water flow must be present in the drinking water system. Administration of the veterinary medicinal product over a period of up to 4 hours on each treatment day, at times when water consumption is likely to be highest prevents precipitation of flubendazole in the water delivery system and allows washing out of the drinking water system within a 24 hour period after the period of drug administration is finished.

- 7) Prior to and after the period of treatment make sure the water distribution system is cleaned.
- 8) Make sure that all animals in the group receive enough drinking water with the veterinary medicinal product. Withhold drinking water for 2 hours before treatment to stimulate thirst.
- 9) The corresponding dose must always be distributed when the water consumption of the animals is highest.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Flubendazole has a low acute oral toxicity.

In hens, no undesirable effects have been observed after administration of up to 15 mg/kg b.w./day flubendazole.

In pigs, no adverse effects have been observed after administration of up to 50 mg/kg b.w./day flubendazole.

In situations where accidental overdose is suspected of having occurred, there is no antidote and treatment should be symptomatic.

4.11 Withdrawal periods

Pigs (meat and offal):

- dose 1 mg/kg body weight for 5 days: 3 days
- dose 2.5 mg/kg body weight for 2 days: 4 days

Chickens (meat and offal): 2 days

Eggs: zero days.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anthelmintics, Benzimidazoles and related substances

ATCvet code: QP52AC12

5.1 Pharmacodynamic properties

Flubendazole is a benzimidazole anthelmintic. It acts by binding to tubulin of the parasite, the dimeric subunit protein of the microtubules. It inhibits micro tubular assembly in absorptive cells: i.e. in intestinal cells of nematodes or the tegumental cells of cestodes. This is shown by disappearance of cytoplasmic microtubules, accumulation of secretory granules in the cytoplasm due to a block in their transport, leading to an impaired coating of the cellular membrane and a decreased digestion and absorption of nutrients. Irreversible lytic degeneration of the cells, due to the accumulation of secretory substances (hydrolytic and proteolytic enzymes) results in the death of the parasite. These changes are relatively fast and are primarily seen in those organelles directly involved in the secretory and absorptive functions of the cells. In contrast the changes are not seen in cells of the host. Another tubulin-related effect is the strong inhibition of egg hatch by inhibition of microtubule-dependended processes in the developing worm egg (cell division).

5.2 Pharmacokinetic particulars

Flubendazole is poorly soluble in aqueous systems, such as the gastrointestinal tract, which results in a low distribution rate and a low absorption. This is reflected by the high faecal excretion of unchanged parent drug. The small fraction absorbed is extensively metabolised by first-pass metabolism in the liver, involving carbamate hydrolysis and ketone reduction. The biotransformation products are conjugated to glucuronides or sulphate conjugates and excreted with the bile and the urine. The excretion with urine is relatively low and consists almost exclusively of metabolites with only small amounts of unchanged compound. In pigs and chickens, the half-life of flubendazole and its metabolites in plasma is 12 hours to 2 days.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium benzoate (E211)
Methyl parahydroxybenzoate (E218)
Disodium edetate
Carmellose sodium
Xanthan gum
Citric acid monohydrate
Carbomers
Propylene glycol
Water, purified

6.2 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

6.3 Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf life after first opening the container: 6 months.

Shelf life after first opening the sachet: Use immediately. Any suspension remaining in the sachet after first opening should be discarded.

Shelf life after dilution according to directions: 24 hours.

6.4 Special precautions for storage

This veterinary medicinal product does not require any special storage conditions.

6.5 Nature and composition of immediate packaging

Box containing 2 sachets (sachet PE/PET/aluminium/PET) of 20 g suspension for use in drinking water.

Box containing 24 sachets (sachet PE/PET/aluminium/PET) of 20 g suspension for use in drinking water.

Box containing 2 sachets (sachet PE/PET/aluminium/PET) of 50 g suspension for use in drinking water.

Box containing 24 sachets (sachet PE/PET/aluminium/PET) of 50 g suspension for use in drinking water.

Box containing 1 sachet (sachet PE/PET/aluminium/PET) of 100 g suspension for use in drinking water.

Box containing 5 sachets (sachet PE/PET/aluminium/PET) of 100 g suspension for use in drinking water.

Box containing 25 sachets (sachet PE/PET/aluminium/PET) of 100 g suspension for use in drinking water.

Box containing 4 containers (PP) with a closure (LDPE) of 750 g suspension for use in drinking water.

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto
Šmarješka cesta 6
8501 Novo mesto
Slovenia

8. MARKETING AUTHORISATION NUMBER

Vm 01656/5063

9. DATE OF FIRST AUTHORISATION

18 July 2013

10. DATE OF REVISION OF THE TEXT

November 2023

PROHIBITION OF SALE, SUPPLY AND/OR USE

Not applicable.

11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' or 'PID' on www.gov.uk.

Approved 07 December 2023

