

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Clinacin 75 mg Tablets

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**Active substance:** Clindamycin 75 mg (as Clindamycin Hydrochloride)

**Excipient(s):** For a full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablets  
Round, white tablet with a breakline on one side

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs

#### **4.2 Indications for use, specifying the target species**

Clinacin 75 mg Tablets are indicated for the treatment of infected wounds, abscesses, superficial pyoderma and oral cavity/dental infections caused by or associated with clindamycin-sensitive staphylococci, streptococci, bacteroidaceae, *Fusobacterium necrophorum*, *Clostridium perfringens* and osteomyelitis caused by *Staphylococcus aureus*. Clinacin 75 mg Tablets can also be used to help provide antimicrobial cover during dental procedures.

#### **4.3 Contraindications**

Do not administer to animals with hypersensitivity to clindamycin and lincomycin preparations.

Do not administer to rabbits, guinea pigs, chinchillas, hamsters, horses or ruminants.

#### **4.4 Special warnings for each target species**

Before the use of Clinacin 75 mg tablets, the identification of causative pathogenic micro-organisms should be carried out and the sensitivity to clindamycin should be established. Clindamycin and lincomycin show parallel resistance. There is a partial cross-resistance to erythromycin and other macrolide antibiotics.

#### **4.5 Special precautions for use**

## **Special precautions for use in animals**

During prolonged therapy of one month or greater, periodic liver and kidney function tests and blood counts should be performed. Patients with severe renal and/or very severe hepatic disturbances accompanied by severe metabolic aberrations should be dosed with caution and should be monitored by serum examination during high dose clindamycin therapy.

## **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

None

### **4.6 Adverse reactions (frequency and seriousness)**

Clindamycin sometimes causes the overgrowth of non sensitive organisms such as resistant clostridia and yeasts. In cases of superinfection, appropriate measures should be taken according to the clinical situation. Vomiting and diarrhoea are observed occasionally

### **4.7 Use during pregnancy, lactation or lay**

While high dose studies in rats suggests that clindamycin is not a teratogen and does not significantly affect the breeding performance of males and females, safety in gestating bitches or breeding male dogs has not been established.

### **4.8 Interaction with other medicinal products and other forms of interaction**

Neuromuscular blocking effects have been observed with clindamycin possibly leading to an increase of efficacy of other neuromuscular blocking agents. The concomitant use of such drugs must be handled with care. Clindamycin should not be used concomitantly with chloramphenicol or macrolides because they may antagonise each other at the site of action.

### **4.9 Amounts to be administered and administration route**

For oral administration.

For treatment of infected wounds, abscesses, oral cavity/dental infections, administer 5.5 mg/kg bodyweight every 12 hours for 7 - 10 days (i.e. 1 tablet per 13.5 kg bodyweight twice daily). Treatment may be extended to a maximum of 28 days based on clinical judgement. If no improvement is seen within 4 days, the sensitivity of the pathogens involved should be redetermined.

For the treatment of superficial pyoderma administer 11 mg/kg every 24 hours (i.e. 2 tablets per 13.5 kg bodyweight once daily). Continue treatment for at least 21 days.

For the treatment of osteomyelitis administer 11 mg/kg every 12 hours (i.e. 2 tablets per 13.5 kg bodyweight twice daily) for at least 28 days. If no improvement is seen within 14 days, the sensitivity of the pathogens involved should be redetermined.

To help provide antimicrobial cover during dental procedures, a 10 day course of 5.5 mg/kg every 12 hours is recommended (i.e. 1 tablet per 13.5 kg twice a day

beginning 5 days before the intended procedure and continuing for 5 days thereafter).

The minimum bodyweight to be treated is 13.5 kg

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Symptoms of overdose include vomiting, inappetency and diarrhoea. In such cases, treatment should be stopped immediately and the dogs treated symptomatically.

#### **4.11 Withdrawal period(s)**

Not applicable

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Antibiotic  
ATCvet code:QJ01FF01

#### **5.1 Pharmacodynamic properties**

Clindamycin, a chlorinated analogue of lincomycin, is an antibiotic with bacteriostatic action. Bactericidal actions have also been reported.

#### **5.2 Pharmacokinetic particulars**

Clindamycin is rapidly absorbed; following oral administration up to 90% of the active ingredient is absorbed from the gastro-intestinal tract.

After a single administration of one tablet to fasting dogs maximum plasma levels ( $C_{max}$ ) of 5  $\mu\text{g/ml}$  are found compared to 3.4  $\mu\text{g/ml}$  in non-fasting dogs.

Bioavailability is greater in fasting dogs than fed dogs.

Clindamycin crosses the placental barrier and can be detected in milk.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Ludipress (Lactose Monohydrate, Povidone and Crospovidone),  
Microcrystalline Cellulose,  
Sodium lauryl sulphate,  
Colloidal silicon dioxide,  
Magnesium stearate.

#### **6.2 Incompatibilities**

None known.

#### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 5 years.

#### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

#### **6.5 Nature and composition of immediate packaging**

Clinacin 75 mg Tablets are presented in white High Density Polyethylene (HDPE) with child resistant tamper evident closure containing 10, 16, 20, 30, 50, 80 or 100 tablets. Not all pack sizes may be marketed.

#### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Chanelle Animal Health Limited  
7 Rodney Street  
Liverpool  
L1 9HZ  
United Kingdom.

### **8. MARKETING AUTHORISATION NUMBER**

Vm 11990/4018

### **9. DATE OF FIRST AUTHORISATION**

26 January 2001

### **10. DATE OF REVISION OF THE TEXT**

March 2019

Approved: 22 March 2019

