



Veterinary  
Medicines  
Directorate

United Kingdom  
Veterinary Medicines Directorate  
Woodham Lane  
New Haw  
Addlestone  
Surrey KT15 3LS

**DECENTRALISED PROCEDURE**

**PUBLICLY AVAILABLE ASSESSMENT REPORT FOR A VETERINARY  
MEDICINAL PRODUCT**

**Strectis 121 mg/60 mg Spot-on Solution for Cats 5-10kg**

**PuAR correct as of 28/03/2018 when RMS was transferred  
to FR. Please contact the RMS for future updates**

**Date Created: June 2015**

## MODULE 1

### PRODUCT SUMMARY

EU Procedure number	UK/V/0547/001/DC
Name, strength and pharmaceutical form	Strectis 121 mg/60 mg spot-on solution for cats 5-10kg
Applicant	Ceva Animal Health Ltd Unit 3, Anglo Office Park White Lion Road Amersham Buckinghamshire HP7 9FB
Active substance(s)	Fipronil (S)-methoprene
ATC Vetcode	QP53AX65
Target species	Cats
Indication for use	<p>Treatment and prevention of flea and/or tick infestations.</p> <p>The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD).</p> <p>Treatment and prevention of flea infestations (<i>Ctenocephalides spp</i>). Immediate insecticidal efficacy against new infestations with adult fleas is seen persisting for 8 weeks. Prevention of the multiplication of fleas by inhibiting the hatching of flea eggs (ovicidal activity) and the development of flea eggs into adult fleas for 6 weeks after application.</p> <p>Treatment and prevention of tick infestation (<i>Rhipicephalus turanicus</i>). The product has immediate and persistent acaricidal efficacy for 5 weeks after application.</p>

## **MODULE 2**

The Summary of Product Characteristics (SPC) for this product is available on the Product Information Database of the Veterinary Medicines Directorate.

[www.gov.uk/check-animal-medicine-licensed](http://www.gov.uk/check-animal-medicine-licensed)

## MODULE 3

### PUBLIC ASSESSMENT REPORT

Legal basis of original application	Full application in accordance with Article 12 (3) of Directive 2001/82/EC as amended.
Date of completion of the original decentralised procedure	16 <sup>th</sup> December 2014
Date product first authorised in the Reference Member State (MRP only)	Not applicable
Concerned Member States for original procedure	Belgium, France, Germany, Italy, Luxembourg and Spain

#### I. SCIENTIFIC OVERVIEW

Strectis 1.21 mg/60 mg Spot-on Solution for Cats 5-10kg is intended for the treatment and prevention of flea and/or tick infestations. There is immediate insecticidal activity against new infestations of adult fleas (*Ctenocephalides* spp) persisting for 8 weeks and prevention of the multiplication of fleas and development of adults from flea eggs for 6 weeks after application. The product also treats and prevents tick infestation (*Rhipicephalus turanicus*) for 5 weeks post-application. Additionally the product can be used to treat Flea Allergy Dermatitis (FAD).

The product is produced and controlled using validated methods and tests which ensure the consistency of the product released onto the market. It has been shown that the product can be safely used in the target species, any reactions observed are indicated in the SPC.<sup>1</sup> The product is safe for the user and for the environment, when used as recommended. Suitable warnings and precautions are indicated in the SPC. The efficacy <sup>2</sup> of the product was demonstrated according to the claims made in the SPC. The overall benefit/risk analysis is in favour of granting a marketing authorisation.

<sup>1</sup> SPC – Summary of product Characteristics.

<sup>2</sup> Efficacy – The production of a desired or intended result.

## **II. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE CONSTITUENTS**

### ***II.A. Composition***

The product contains Fipronil and (S)-methoprene as active substances. The excipients used are butylhydroxyanisole E320, butylhydroxytoluene (E321), ethanol (anhydrous) and Diethylene glycol monoethyl ether.

The container/closure system consists of:

Front foil: Polypropylene / polyethylene terephthalate

Lidding foil: Polyester / aluminium / polyester / polyethylene terephthalate

The pipettes are packed in child resistant blisters and packaged into packs containing 1, 3, 6, 12, 24, 60 or 120 pipettes. The particulars of the containers and controls performed are provided and conform to the regulation.

The choice of the formulation and the presence of preservative are justified. The product is an established pharmaceutical form and its development is adequately described in accordance with the relevant European guidelines.

### ***II.B. Description of the Manufacturing Method***

The product is manufactured fully in accordance with the principles of good manufacturing practice from a licensed manufacturing site. The product is manufactured using conventional manufacturing techniques. The manufacturing process is a mixing and filtering procedure, followed by shipping of bulk product for filling into pipettes. Process validation for full-scale batches will be performed post-authorisation.

### ***II.C. Control of Starting Materials***

The active substances are fipronil and (S)-methoprene that are established active substances not described in the European Pharmacopoeia (Ph. Eur.). They are manufactured in accordance with the principles of good manufacturing practice.

Each active substance specification is considered adequate to control the quality of the material. Batch analytical data demonstrating compliance with this specification have been provided. Relevant ASMF were submitted.

The excipients are manufactured in accordance with the relevant Ph. Eur. monographs and certificates of analysis have been provided.

#### ***II.C.4. Substances of Biological Origin***

There are no substances within the scope of the TSE Guideline present or used in the manufacture of this product.

#### ***II.D. Control Tests Carried Out at Intermediate Stages of the Manufacturing Process***

The tests performed during production are described and the results of 3 consecutive runs, conforming to the specifications, are provided.

#### ***II.E. Control Tests on the Finished Product***

The finished product specification controls the relevant parameters for the pharmaceutical form. The tests in the specification, and their limits, have been justified and are considered appropriate to adequately control the quality of the product. Satisfactory validation data for the analytical methods have been provided. Batch analytical data from the proposed production site have been provided demonstrating compliance with the specification. Tests include those for appearance, identity and content of active substances and excipients, water content and microbiological purity.

#### ***II.F. Stability***

Stability data on the active substances have been provided in accordance with applicable European guidelines, demonstrating the stability of the active substance when stored under the approved conditions.

##### Fipronil

Three production batches were stored in commercial packaging under VICH conditions at 25°C/60% RH and 40°C/75% RH. Data provided demonstrated that the active substance is stable under these conditions. A retest period of 3 years is acceptable.

##### (S)-methoprene

Commercially prepared batches of the active substance were stored under nitrogen in light-resistant containers at 25°C/60% RH and at 40°C/75% RH. Active substance stored at 25°C/60% RH was tested at 0, 3, 6, 12, 18 and 24 months. Active substance stored at 40°C/75% RH was tested at 0, 1, 3 and 6 months. A retest period was established as being 2 years when stored at 2-8°C.

Stability tests on the finished product were provided for product stored in commercial packaging for up to 24 months at 25°C/60% RH and at up to 6 months at 40°C/75% RH. A shelf-life of 3 years was established.

## **G. Other Information**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years

## **III. SAFETY AND RESIDUES DOCUMENTATION (PHARMACOTOXICOLOGICAL)**

The applicant provided bibliographical data in support of the toxicological aspects of the two active substances, and also provided studies on the irritation and dermal sensitisation of the products. A user risk assessment (URA) and environmental risk assessment (ERA) were also provided.

### **III.A Safety Documentation**

#### **Pharmacological Studies**

##### **Pharmacodynamics**

###### **Fipronil**

In the target organisms, fipronil antagonises the gamma aminobutyric acid (GABA) channels resulting in the blocking of the pre-synaptic transfer of chloride ions across the cell membrane. This results in uncontrolled activity of the nervous system, a manifestation of the insecticidal and acaricidal activity of fipronil. Fipronil additionally binds glutamate activated chloride channels not found in mammals.

###### **(S)-methoprene**

This active substance is an analogue of the juvenile hormone of insects. Addition of (S)-methoprene therefore causes abnormalities to occur during the developmental stages of the target organisms, it additionally, has negative effects on the reproductive capacity of adult insects.

##### **Pharmacokinetics**

###### **Fipronil**

Dermal absorption studies were provided for humans, rats and rabbits. At 200 mg/ml, dermal penetration was seen to a greater extent in rabbits and rats, but at 0.2 mg/ml dermal penetration was similar between human, rabbits and rats. In a rat dermal study using <sup>14</sup>C-fipronil, the quantity of active substance absorbed was less than 1% of the dose applied. In a single oral dose study in rats using radio-labelled fipronil, the active substance was found to be widely distributed. Faeces appeared to be the main route of excretion.

Sulfone, amide and reduction products are common metabolites of fipronil metabolism, and fipronil is commonly found within the hair and hair follicles of treated animals.

### (S)-methoprene

A single radio-labelled dose of 25 mg/kg was administered orally to rats in one study. Peak concentration occurred after approximately 6 hours, which was followed by a slow decline with a 48 hour half-life. Distribution in rats was found to be in the liver, kidney and lungs, with peak concentration seen at 6-12 hours after dosing. Less of the active substance was seen in fat and muscle. A further study on the metabolism of (S)-methoprene mammals, showed that much is oxidised. The active substance is in general excreted biphasically, with a rapid first phase, followed by a slower second phase.

Suitable data were submitted on the use of the combination of the active substances.

### **Toxicological Studies**

The applicant has provided bibliographical data for toxicological studies.

- **Single Dose Toxicity**

Fipronil was found to be acutely toxic to rats and mice following oral administration, and was slightly toxic to rabbits via the dermal route. In general, (s)-methoprene is considered less toxic than fipronil.

- **Repeated Dose Toxicity**

### Fipronil

Studies were provided for repeat dose studies. In rats, dietary doses were given at 0, 0.07, 0.33, 1.93, or 19.87 mg/kg/day to males and 0, 0.07, 0.37, 2.28 and 24.03 mg/kg/day to females. In this study, the NOEL<sup>3</sup> was established at 0.33 mg/kg/day for males and 0.37 mg/kg/day for females. An oral study in dogs given doses at 0, 0.5, 2.0 and 10 mg/kg/day provided NOEL of 2 mg/kg/day for males and 0.5 mg/kg/day for females. In a further study, gelatine capsules given to dogs at 0, 0.2, 2 or 5 mg/kg/day product a NOEL of 0.2 mg/kg/day in both males and females. Neurotoxic effects were noted.

### (S)-methoprene

Racemic methoprene was administered to male and female dogs at doses of 6.2, 12 and 120 mg/kg/day for 90 days. No deaths occurred, but an increase in liver weight was noted. A NOAEL<sup>4</sup> of 8.6 mg/kg/day was confirmed.

- **Reproductive Toxicity, including Teratogenicity:**

### Fipronil

In one study, rates were administered fipronil in the diet at 0, 0.25, 2.5 and 26 mg/kg/day (male rats), and 0, 0.27, 2.7 and 28 mg/kg/day (female rats). At

---

<sup>3</sup> NOEL – No observed effect level.

<sup>4</sup> NOAEL – No observed adverse effect level.



amounts greater than 2.5 mg/kg/day, systemic effects were seen in the parental animals. Litters of treated parental animals showed adverse effects when given the fipronil-containing diet, and a reduction in fertility was also noted. The NOAEL for parental toxicity was 0.25 mg/kg/day while for reproductive toxicity, it was observed to be 2.5 mg/kg/day.

A study in rats in which fipronil was administered to the cervical region of female rats at doses of 70, 140 and 280 mg/kg/day showed that progesterone levels were increased and oestrodial levels were reduced. This altered the ovulating cycle of the rats. Fipronil reduced the pregnancy index in the highest dose group during mating studies, although other factors such as body weight, weaning weight, implantation and the number of resorptions were not affected.

In addition to further data provided for embryotoxic studies, it was established that fipronil has an effect on reproduction in laboratory animals. Suitable warnings appear in the SPC for use of the product in dogs and cats, i.e. use of the product during pregnancy and lactation is recommended only after the benefit/risk assessment by the responsible veterinarian. A further study in rabbits, administered fipronil at 0, 0.1, 0.2, 0.5 and 1 mg/kg/day demonstrated no treatment-related effect, and established the NOAEL at 1 mg/kg/day.

#### (S)-methoprene

In one study, rats were fed 0, 25 and 75 mg/kg/day racemic methoprene prior to mating. Pups from ensuing litters were utilised in follow-on mating studies. Evidence from the studies suggested that the NOAEL was 33 mg/kg/day, (29mg/kg/day when corrected for purity). A further study suggested a NOAEL for embryotoxicity of 570 mg/kg/day, (the highest dose tested).

In rabbits, a development study using racemic methoprene showed reduced body weight and increase frequency of abortions and the highest dose used, 2000 mg/kg/day. The NOAEL for both maternal and foetal toxicity was established as being 200 mg/kg/day.

- Mutagenicity

Published data was provided that showed the active substances tested negative for genotoxicity.

- Carcinogenicity

#### Fipronil

Fipronil was added to the diet of mice for 78 weeks at doses 0, 0.1, 0.5, 10, 30 or 60 parts per million. An adverse effect possibly due to an increase in liver weights was seen in the highest dose group only. A NOAEL was established at the equivalent of 0.055 mg/kg/day.

In a further combined carcinogenicity/toxicity study, rats were dosed in the diet with fipronil at 0, 0.019, 0.059, 1.27 or 12.68 mg/kg/day and at 0, 0.025, 0.078, 1.61, or 16.75 mg/kg/day for females. High mortality rates prevented the

conclusion of the study with chemical changes and organ alterations noted at the two highest dose rates. Benign and malignant changes occurred in the thyroid glands of both sexes. The NOEL was established for males at 0.019 mg/kg/day and for females at 0.025 mg/kg/day.

#### (S)-methoprene

Racemic methoprene was given in the diet to mice at concentrations of 38, 150 and 380 mg/kg/day. The notable adverse effect was on changes to the liver which led to the establishment of a NOAEL equivalent to 130 mg/kg/day when corrected for purity.

#### ***Studies of Other Effects***

##### Fipronil

Fipronil was administered orally to rats at doses of 0, 0.5, 5 and 50 mg/kg. Clinical toxicity was only observed at the highest dose, with neurotoxic signs being more prevalent in male mice and convulsions seen in both sexes. A NOAEL of 0.5 mg/kg was established. A further study using a spot-on solution was performed in rats, with the active substance applied at 1, 2 and 4 times recommended dose; equivalent to 70, 140 and 280 mg/kg. Some behavioural abnormalities were observed, related particularly to the highest dose.

In a developmental neurotoxicity study in rats, fipronil was administered at 0.05, 0.90 and 15 mg/kg/day. The maternal NOEL was 0.90 mg/kg/day. The NOEL for developmental toxicity was established as 0.05 mg/kg/day, based on a reduction in pup weights and an increase in preputial separation in males.

Studies on thyroid function, which may be altered by fipronil, indicated that biliary clearance of the hormone was affected. This may cause an increase in thyroid-stimulating hormone.

##### (S)-methoprene

(S)-methoprene was shown to have no androgenic, oestrogenic, glucocorticoid or anabolic activity.

##### Combined product

In a variety of studies, the combined product was shown not to be an irritant or sensitiser to skin. Fipronil desulfinyl, the significant metabolite of fipronil seen after photodegradation, was also not observed to cause irritation or sensitisation problems. An investigation to determine the level of active substances on gloves after petting dogs showed that levels were highest 12 hours after administration for all compounds and declined over the course of the treatment to 0.5% by 7 days.

### ***Observations in Humans***

Published data on the adverse effects of fipronil in humans suggests that acute exposure causes no long-term harm and is self-limiting. Adverse reactions include vomiting, drowsiness headache, vertigo and sweating. There were no data relating to the absorption of (S)-methoprene by humans.

### ***User Safety***

The applicant has provided a user risk assessment in compliance with the relevant guideline, which discussed possible routes of exposure.

Warnings and precautions as listed on the product literature are adequate to ensure safety to users of the product.

- This product can cause eye irritation.
- Avoid contact of the product with skin, eyes or mouth. People with known hypersensitivity to any of the ingredients should not treat their animal with this product.
- Treated animals should not be handled or played with for at least 12-hours after treatment. Animals should be treated in the evening in order to minimise contact with the treated animal. On the day of treatment, treated animals should not be permitted to sleep with their owner, especially children.
- Do not eat, drink or smoke while handling the product.
- Wash hands thoroughly after use.
- In case of accidental spillage on skin, wash off immediately with soap and water.
- If the product accidentally gets into the eyes, they should be thoroughly flushed with water.
- If the product is accidentally swallowed, seek medical advice immediately and show the package leaflet to the physician.

### ***Environmental Safety***

The applicant provided a Phase I environmental risk assessment in compliance with the relevant guideline. The product is not to be used in food-producing animals, and the product is not considered to cause a threat to the environment when used as recommended. Warnings and precautions as listed on the product literature are adequate to ensure safety to the environment when the product is used as directed, with a direction that recently treated dogs should not enter watercourses.

## IV CLINICAL DOCUMENTATION

### *IV.1. Pre-Clinical Studies*

#### **Pharmacology**

Refer to Section III, Safety Testing, Pharmacological Studies. Suitable data were provided.

#### Fipronil - Mode of action

The mode of action has been supported by bibliographical references. The principal drug target for fipronil is the GABA gated chloride channel. Binding results in GABA antagonism leading to neurotoxicity and death in invertebrates. Fipronil can result in neurotoxicity in mammals at high doses however; insects are significantly more sensitive to the toxic effects. Fipronil has been demonstrated to have an adulticidal effect in fleas and ticks.

#### S-methoprene - Mode of action

The mode of action of S-methoprene has been supported by bibliographical references. S-methoprene is an insect growth regulator (IGR) with ovicidal and larvicidal activity. Specifically it is defined as a juvenile hormone analogue and its principal target is the insect neuroendocrine system where it exerts a regulatory effect at the level of gene expression. Studies have demonstrated the inhibitory effects of S-methoprene on insect development, with the key periods of sensitivity being early embryonic development and metamorphosis. S-methoprene has both an ovicidal and larvicidal effect on fleas when used topically to treat cats and dogs.

#### Fixed combination of fipronil/(s)-methoprene

The insecticidal and acaricidal effects of fipronil can be demonstrated alongside the IGR effects of S-methoprene in the same animal after topical application.

### **Tolerance in the Target Species**

The applicant has conducted target animal tolerance studies using multiples of the recommended dose in the target species. A GLP<sup>5</sup>-compliant study was performed in a suitable number of clinically eligible young cats, (32 kittens: 18 male and 14 female; age: 8 weeks  $\pm$  2 days on day 0; weight range: 0.27 kg to 0.97 kg on day 0), using a spot-on product containing 1x, 3x or 5x the maximum recommended dose of the final product, over a period of 7 fortnightly applications. Placebo was used as a negative control. This was a five phase, parallel group, randomised, blinded, controlled study. Appropriate observations and clinical measurements were made at suitable time points. Analysis showed that the product was tolerated at x 5 the highest maximum recommended dose level (1.75 ml/kg), for a 12 week period.

A second study evaluated the oral safety of the combined product in adult cats. A 0.35 ml/kg dose was administered to a suitable number of clinically eligible animals, along with a placebo administered at the same dose to additional

---

<sup>5</sup> GLP – Good Laboratory Practice.

animals. This was a GLP-compliant study. The study design was parallel grouped, randomised, blinded and controlled. Placebo was used as a negative control. Appropriate observations and clinical measurements were made at suitable time points. No adverse reactions were seen.

### **Resistance**

Bibliographical references were provided in relation to the possible resistance of ticks and fleas to fipronil, not thought to be a major issue in Europe at the current time. No data is available for (S)-methoprene. Adequate warnings and precautions appear on the product literature and in the SPC.

## **IV.II. Clinical Documentation**

### **Laboratory Trials**

The applicant conducted dose titration and dose confirmation studies, and field clinical trials to demonstrate the efficacy in cats.

#### Efficacy summary for cats:

IGR efficacy against the further development of flea eggs and larvae:

- During a dose titration study, ovicidal and larvicidal efficacy was demonstrated at 6 mg/kg S-methoprene.
- Persistence of IGR (ovicidal activity and prevention of the development of flea eggs to adult fleas) efficacy was shown during two dose confirmation studies until 47 days and 52 days post-treatment respectively.

Adulticidal efficacy against fleas:

- Minimum effective dose 12 mg/kg of fipronil against *Ctenocephalides felis* was shown in a dose titration study.
- Immediate efficacy above 95% was demonstrated during dose titration and dose confirmation studies
- Persistence of adulticidal efficacy until 44 days and 58 days post-treatment was shown during two dose confirmation studies respectively.
- A field study in representative cats across two geographic regions demonstrated non-inferiority to the reference product Frontline Combo Cat, Merial. These studies were randomised controlled studies over 28 days.

Adulticidal efficacy of fipronil against ticks:

- Minimum effective dose 12 mg/kg of fipronil against *Rhipicephalus turanicus*. was shown in a dose titration study.
- Immediate efficacy against *R. turanicus* was demonstrated during two dose confirmation studies. Persistence of efficacy against *R. turanicus* for 30 days and 37 days post-treatment was demonstrated during two dose confirmation studies respectively.

The following indications in cats are supported for the proposed fixed combination:

"Treatment and prevention of flea and/or tick infestations.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD).

Treatment and prevention of flea infestations (*Ctenocephalides spp*). Immediate insecticidal efficacy against new infestations with adult fleas is seen persisting for 8 weeks. Prevention of the multiplication of fleas by inhibiting the hatching of flea eggs (ovicidal activity) and the development of flea eggs into adult fleas for 6 weeks after application.

Treatment and prevention of tick infestation (*Rhipicephalus turanicus*). The product has immediate and persistent acaricidal efficacy for 5 weeks after application."

### **Dose confirmation studies:**

The applicant provided appropriate information on dose determination studies.

A number of dose confirmation studies were presented:

#### **Study 1**

Study title	Dose confirmation efficacy study of a fipronil/(S)-methoprene spot-on formulation against fleas ( <i>Ctenocephalides felis</i> ) on cats
Objectives	<ul style="list-style-type: none"> <li>To confirm the adulticidal efficacy of a fipronil/S-methoprene spot-on formulation against fleas (<i>Ctenocephalides felis</i>) on cats.</li> <li>To confirm the efficacy of a fipronil/S-methoprene spot-on formulation against the further development of flea (<i>Ctenocephalides felis</i>) eggs.</li> <li>To observe any possible adverse events related to the administration of the investigational veterinary product.</li> </ul>
Test site(s)	Single centre.
Compliance with Regulatory guidelines	Good Clinical Practice (GCP).
Test Product	17% fipronil and 8.5% (S)-methoprene  Dose administered dependent on day 0 weight: <ul style="list-style-type: none"> <li>0.35 ml/cat weighing 1 - 5 kg</li> <li>0.71 ml/cat weighing &gt;5 - 10 kg</li> </ul>
Control product/placebo	Negative control.
Animals	16 healthy cats, 8 animals in each group
Outcomes/endpoints	<p>Primary endpoints: The number of adult fleas, the proportion of hatched eggs, and the proportion of adults emerging at each time point in both control and treatment groups.</p> <p>Safety endpoints: No specific adverse events were defined in the protocol, however during the study the health of animals was monitored closely to identify potential adverse events.</p>
Randomisation	Randomised.
Blinding	Blinded.
Method	This was a parallel-grouped study. After acclimatisation,

	animals were given treatment depending of their respective group. At various time points according to study schedule, animals were infested as appropriate (approximately 100 fleas per cat), flea counts were performed, flea eggs were collected, egg hatch and adult flea emergence were assessed to establish adulticidal, ovicidal and larvicidal efficacy.
Statistical method	Statistical analysis was performed using appropriate software. Level of significance was set at 5% (p<0.05). Comparisons were made by ANOVA.  Efficacy against adult fleas, flea egg and larval development (inhibition of adult emergence) was calculated for the various assessment days using the arithmetic mean. The IVP was regarded effective if the efficacy against egg hatch and adult flea emergence was >90% and the adulticidal efficacy against fleas was >95%.
<b>RESULTS</b>	
Outcomes for endpoints	Adulticidal efficacy of the IVP remained above 95% at all time points and until day +44. Ovicidal efficacy of the IVP remained above 90% at all time points and until day +47. Larvicidal efficacy remained above 90% until day +33; on day +40 the efficacy was 89.3% and on day +47 the efficacy was 97.4%. No treatment related adverse events were observed.
DISCUSSION	A fixed combination of 17% fipronil and 8.5% S-methoprene, administered at the recommended dose, demonstrated immediate adulticidal efficacy against existing flea ( <i>Ctenocephalides felis</i> ) infestations on cats and was persistently effective for 6 weeks. The product was effective in preventing the further development of flea eggs (ovicidal activity) for at least 6 weeks post-treatment (47 days).

## **Study 2**

Study title	Dose confirmation study of a fipronil/(S)-methoprene spot-on combination formulation against the further development of flea ( <i>Ctenocephalides felis</i> ) eggs (ovicidal and larvicidal activity) collected from treated cats.
Objectives	To confirm the efficacy of the target dose of a fipronil / (s)-methoprene spot-on combination formulation against the further development of flea ( <i>Ctenocephalides felis</i> ) eggs (ovicidal and larvicidal activity) collected from treated cats.
Test site(s)	Single centre.
Compliance with Regulatory guidelines	Good Clinical Practice (GCP).
Test Product	17% fipronil and 8.5% (S)-methoprene Dose: 0.071 ml/kg (minimum product dose to be received, based on proposed product posology).
Control product/placebo	Negative control.
Animals	16 healthy cats, 8 animals in each group.
Outcomes/endpoints	Primary endpoint: Inhibition of adult emergence. Proportion of emerged adult fleas per incubated egg.

	<p>Secondary endpoint: Ovicidal product effect. Proportion of hatched larvae per incubated egg.</p> <p>Safety endpoints: None specifically defined, however health of enrolled animals was monitored closely to identify potential adverse events.</p>
Randomisation	Randomised.
Blinding	Blinded.
Method	This was a parallel-grouped study. After acclimatisation, animals were given treatment depending of their respective group. At various time points according to study schedule, animals were infested as appropriate (approximately 100 fleas per cat), flea eggs were collected, egg hatch and adult flea emergence were assessed to establish ovicidal and larvicidal efficacy.
Statistical method	<p>Statistical analysis was performed using appropriate software. Level of significance was set at 5% (p&lt;0.05). Comparisons were made by ANOVA.</p> <p>Inhibition of adult emergence and ovicidal product effect were calculated for the various assessment days using the arithmetic mean.</p> <p>The IVP was regarded effective when inhibition of adult flea emergence was &gt;90% compared to controls.</p>
<b>RESULTS</b>	
Outcomes for endpoints	<p>Reported ovicidal efficacy was &gt;90% at all time points and for up to +45 days post treatment.</p> <p>Adult flea emergence efficacy was &gt;90% at all time points and for up to +52 days post treatment.</p> <p>None of the adverse events observed during the study were in the opinion of the examining veterinarian related to administration of the IVP.</p>
<b>DISCUSSION</b>	
	The IVP was safe in cats and >90% effective in preventing the further development of flea eggs to adult fleas for up seven weeks (52 days) post treatment.

### **Study 3**

Study title	A dose confirmation study to determine the efficacy of a combined fipronil/(S)-methoprene product against fleas ( <i>Ctenocephalides felis</i> ) on cats under laboratory conditions.
Objectives	The objective of this dose confirmation study was to determine the efficacy after a single application, of a combined product (17% fipronil / 8.5% S-methoprene) when compared with an untreated control group against artificially induced infestations of fleas ( <i>Ctenocephalides felis</i> ) on cats.
Test site(s)	Single centre.
Compliance with Regulatory guidelines	Good Clinical Practice (GCP).
Test Product	17% fipronil and 8.5% (S)-methoproene



	Dose: 0.071 ml/kg (minimum product dose to be received, based on proposed product posology).
Control product/placebo	Negative control.
Animals	16 healthy cats, 8 animals each group.
Outcomes/endpoints	Primary endpoint: flea adulticidal activity. Flea count compared to baseline (control) count.  Safety endpoints: None specifically defined, however health of enrolled animals was monitored closely to identify potential adverse events.
Randomisation	Randomised.
Blinding	Blinded.
Method	This was a parallel-grouped study. After acclimatisation, animals were given treatment depending of their respective group. At various time points according to study schedule, animals were infested as appropriate (approximately 100 fleas per cat) and flea counts were performed to establish adulticidal efficacy.
Statistical method	Statistical analysis was performed using appropriate software. Level of significance was set at 5% (p<0.05). Comparisons were made by ANOVA.  Efficacy against adult fleas was calculated for the various assessment days using the arithmetic mean.  The IVP was regarded effective when >95% reduction in flea count for treated animals compared to controls was achieved.
<b>RESULTS</b>	
Outcomes for endpoints	Adulticidal efficacy of the IVP was 100% at all time points and until day +58. There were no adverse reactions.
<b>DISCUSSION</b>	
	The results of this study demonstrate that a single topical application of a flea treatment (containing 17% fipronil and 8.5% (S)-methoprene solution) administered at 0.071 ml/kg was 100% effective (based on arithmetic means) against artificially induced infestations of <i>Ctenocephalides felis</i> fleas on cats when compared to the untreated control cats. There were no adverse reactions.

#### **Study 4**

Study title	Dose confirmation efficacy study of a fipronil/S-methoprene spot-on formulation against ticks ( <i>Rhipicephalus turanicus</i> ) on cats.
Objectives	To confirm the efficacy of the target dose of a fipronil/(S)-methoprene combination formulation against ticks ( <i>Rhipicephalus turanicus</i> ) on cats. To observe any possible adverse events related to the administration of the investigational veterinary product.

Test site(s)	Single centre.
Compliance with Regulatory guidelines	Good Clinical Practice (GCP).
Test Product	17% fipronil and 8.5% (S)-methoprene  Dose administered dependent on day 0 weight: <ul style="list-style-type: none"> <li>• 0.35 ml/cat weighing 1 - 5 kg</li> <li>• 0.71 ml/cat weighing &gt;5 - 10 kg</li> </ul>
Control product/placebo	Negative control.
Animals	14 healthy cats, 7 animals each group.
Outcomes/endpoints	Primary endpoint: Adulticidal efficacy against ticks. Safety endpoints: None specifically defined, however health of enrolled animals was monitored closely to identify potential adverse events.
Randomisation	Randomised.
Blinding	Blinded.
Method	This was a parallel-grouped study. After acclimatisation, animals were given treatment depending of their respective group. At various time points according to study schedule, animals were infested as appropriate (50 ticks per cat), tick counts were performed to establish acaricidal efficacy. (Tick counts were conducted 48 ± 2 hours post infestation or treatment).
Statistical method	Statistical analysis was performed using appropriate software. Level of significance was set at 5% (p<0.05). Comparisons were made by ANOVA.  Efficacy against adult ticks was calculated for the various assessment days using the arithmetic mean. The IVP was regarded as effective when the efficacy was greater than or equal to 90%.
<b>RESULTS</b>	
Outcomes for endpoints	Adulticidal efficacy of the IVP remained above 90% at all time points (until at least day +28). No treatment related adverse events were recorded in any cats.
<b>DISCUSSION</b>	17% fipronil and 8.5% S-methoprene, administered at the target dose had a greater than 90% immediate (day +2) efficacy and persistent efficacies greater than 90% against <i>Rhipicephalus turanicus</i> for four weeks (day +28).

### **Study 5**

Study title	Dose confirmation efficacy study of a fipronil / (S)-methoprene spot-on formulation against ticks ( <i>Rhipicephalus turanicus</i> ) on cats.
Objectives	To confirm the efficacy of the target dose of a fipronil / (S)-methoprene combination formulation against ticks ( <i>Rhipicephalus turanicus</i> ) on cats. To observe any possible

	adverse events related to the administration of the investigational veterinary products.
Test site(s)	Single centre.
Compliance with Regulatory guidelines	Good Clinical Practice (GCP).
Test Product	17% fipronil and 8.5% (S)-methoprene  Dose: 0.071 ml/kg (minimum product dose to be received, based on proposed product posology).
Control product/placebo	Negative control.
Animals	16 healthy cats, 8 animals each group.
Outcomes/endpoints	Primary endpoint: Adulticidal efficacy against ticks.  Safety endpoints: None specifically defined, however health of enrolled animals was monitored closely to identify potential adverse events.
Randomisation	Randomised.
Blinding	Blinded.
Method	This was a parallel-grouped study. After acclimatisation, animals were given treatment depending of their respective group. At various time points according to study schedule, animals were infested as appropriate (50 ticks per cat), tick counts were performed to establish acaricidal efficacy. Elizabethan collars were used to prevent ticks being groomed off. Tick counts were performed 48 hours post-treatment or infestation according to a systematic protocol.
Statistical method	Statistical analysis was performed using appropriate software. Level of significance was set at 5% ( $p < 0.05$ ). Comparisons were made by ANOVA. Efficacy against ticks was calculated for the various assessment days using the arithmetic mean. The IVP was regarded effective when adulticidal efficacy was $>90\%$ compared to untreated controls.
<b>RESULTS</b>	
Outcomes for endpoints	Reported adulticidal efficacy: 96.3% on day +2, and between 99.2% and 100% at all other assessment points up to +35days. For both day +2 and +30 the mean tick count from control animals was below 12.5 ticks (25%) and therefore below the CVMP specified minimum. On day +2 this was considered to be due to the four day period between infestation and assessment; on day +30 the lower count was not considered of impact on the conclusions due to the zero tick recovery from IVP treated cats. Efficacy was therefore $>90\%$ from +2 to +37 days post treatment. Adverse events observed were not due to treatment.
DISCUSSION	The IVP (17% fipronil and 8.5% (S)-methoprene solution) was immediately and persistently effective ( $\geq 90\%$ ) against <i>Rhipicephalus turanicus</i> infestations on cats for up to five weeks post treatment when administered at a dosage of 0.071 ml/kg.

## Field Trials

### **Study 6 (Pivotal Field Study)**

Study title	Field evaluation of the efficacy and safety of a topical formulation of Fipronil + (S)-methoprene product in the treatment and prevention of natural infestation of fleas on cats presented as veterinary patients in Europe.
Objectives	Evaluation of the efficacy and safety (over 28 days) of a spot-on solution for Cats (17% fipronil / 8.5% methoprene) in the treatment and prevention of naturally acquired infestations of fleas on cat. The study evaluated the product at a minimum recommended dose of 12 mg/kg of fipronil and 6 mg/kg of S-methoprene spot-on solution (IVP) administered once topically to the skin. The study also evaluated (over 28 days) the effectiveness of the IVP as part of a treatment strategy for the control of flea allergy dermatitis (FAD).
Test site(s)	Multicentre.
Compliance with Regulatory guidelines	Good Clinical Practice (GCP)/ Good Laboratory Practice (GLP).
Test Product	17% fipronil and 8.5% (S)-methoprene (Final formulation)  Pipette volume / dose: IVP1: Small cats ( $\geq 1.0$ kg - $\leq 5$ kg) – 0.35 ml IVP2: Large cats ( $> 5$ kg) – 0.71 ml
Control product/placebo	Positive controls: The Control Veterinary Product: Frontline Combo Spot-on Cat (fipronil and S-methoprene, Merial)  Pipette volume / dose: CVP1: Cats ( $\geq 1.0$ kg) – 0.5 ml (50 mg fipronil and 60 mg S-methoprene).
Animals	203 cats, 95 female, 108 male, aged 3 months to 16 years.
Outcomes/endpoints	Primary endpoints: 1. Efficacy against fleas: presence or absence of fleas on cats. 2. Safety: clinical observations.  Flea counts and clinical observations were performed by a veterinarian on day 0 (baseline) and day +14 ( $\pm 2$ days) and +28 ( $\pm 2$ days). All treated cats with 2 or more fleas on day 0 were included in the efficacy analysis. All treated cats and dogs were included in the safety analysis.  Secondary endpoints: 1. Efficacy against fleas: analysis of arithmetic and geometric mean flea counts, and the reduction in geometric mean flea counts in each treatment group compared with day 0. 2. Efficacy against flea allergic dermatitis (FAD): assessed based on recorded dermatological findings in cats.
Randomisation	Randomised.
Blinding	Parallel group, blinded.
Statistical method	Statistical protocol was defined a priori.

	<p>The statistical unit was the individual cat. The per protocol population (PP) were defined as all treated cats with two or more fleas on day zero that had met all the eligibility criteria and followed the protocol rigidly. The intention to treat population (ITT) were defined as all cat treated with the product regardless of the presence of fleas on day 0.</p> <p>Data analysis was conducted using appropriate software. A significance level of 5% (P&lt;0.05) was applied to all statistical tests. No adjustment for multiple tests was performed. Application conditions for each treatment group were tested for homogeneity of variance and for normality before applying appropriate tests.</p> <p>Efficacy:</p> <p>Primary efficacy endpoint (presence or absence of fleas on 'per-protocol' (PP) cats on days +14 to +28 was tested by comparing the proportion of cats with no fleas on day +14 to the end of the trial in the IVP and CVP groups. Data was analysed with a one-sided non-inferiority test and delta -15%.</p> <p>Secondary endpoints were also identified and assessed.</p> <p>Flea count data were Log transformed prior to analysis. Percent efficacy for each treatment group and time point was calculated as the percent change from Day 0 in geometric mean flea count.</p> <p>The differences between study days in each treatment group were tested for significance, and the percentage reduction in treatment groups was compared at each post-treatment time point.</p> <p>Clinical signs of FAD (flea allergy dermatitis), examination of PP cats for signs of FAD on Day 0, were summarised. For PP cats with signs of FAD on day 0, a summary of the evolution of clinical signs of FAD over time in each treatment group (T01 and T02) was performed.</p> <p>Safety assessment: Adverse and suspected adverse drug reactions were reported and summarised in frequency tables.</p>
<b>RESULTS</b>	
Outcomes for endpoints	<p>Baseline measurements were appropriately assessed.</p> <p>Efficacy: The result of the analyses showed a non-significant difference between the IVP and CVP groups at both efficacy assessment time points. Statistical analysis for non-inferiority was established for both the primary efficacy criterion (the percentage of cats with no (zero) fleas from Days 14 to 28), and the secondary efficacy criteria (analysis of arithmetic and geometric mean flea counts on Days 14 and 28, and the percentage reductions in geometric mean</p>

	<p>flea counts on Days 14 and 28 compared with Day 0).</p> <p><b>Safety:</b> The field safety of the IVP was demonstrated. The potential risk of local adverse reactions at the site of application should be considered. No serious adverse events occurred in the ITT population and no significant safety concerns were raised. Concomitant treatments were administered and none interfered with the interpretation or integrity of the study and there was no evidence of interactions.</p>
DISCUSSION	<p>The IVP, applied as a single topical administration at a recommended minimum dosage of fipronil 12 mg/kg and S-methoprene 6 mg/kg, was well tolerated and clinically efficacious in cats presented as veterinary patients in Europe. Statistical analysis demonstrated that it was non-inferior to the CVP (an approved product in the treatment and prevention of fleas for 28 (<math>\pm 2</math>) days on cats). Administration to a wide range of cats during the trial and with a variety of concomitant medications under typical field conditions in Europe resulted in no drug interactions and only 3 adverse events occurred during the trial from which all cats completely recovered.</p> <p>Clinical signs of FAD in cats improved throughout the study (for 28 (<math>\pm 2</math>) days), compared with Day 0, in both treatment groups. The alleviation of a range of clinical signs of FAD over time, in cats treated with IVP, indicates that the product may be beneficial as part of a treatment strategy for control of FAD in cats.</p>

All data contributed to the formulation being assessed as acceptable for the indications as specified in the SPC.

## V OVERALL CONCLUSION AND BENEFIT– RISK ASSESSMENT

The data submitted in the dossier demonstrate that when the product is used in accordance with the Summary of Product Characteristics, the benefit/risk profile of the product is favourable.

## **MODULE 4**

### **POST-AUTHORISATION ASSESSMENTS**

The SPC and package leaflet may be updated to include new information on the quality, safety and efficacy of the veterinary medicinal product. The current SPC is available on the Product Information Database of the Veterinary Medicines Directorate website.

([www.gov.uk/check-animal-medicine-licensed](http://www.gov.uk/check-animal-medicine-licensed))

The post-authorisation assessment (PAA) contains information on significant changes which have been made after the original procedure which are important for the quality, safety or efficacy of the product.

The PAA for this product is available on the Product Information Database of the Veterinary Medicines Directorate website.

([www.gov.uk/check-animal-medicine-licensed](http://www.gov.uk/check-animal-medicine-licensed))