

ANNEX

Identification number of the additive	Name of the holder of authorisation	Additive (Trade name)	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content		Maximum content	Other provisions	End of period of authorisation	Provisional maximum residue limits (MRLs) in the relevant foodstuffs of animal origin
						mg of active substance/kg of complete feedstuff with a moisture content of 12 %	mg of active substance/kg of complete feedstuff with a moisture content of 12 %				
Coccidiostats and histomonostats											
51701	Huvepharma NV Belgium	Monensin sodium (Coxidin)	<p><i>Additive composition</i></p> <p>Monensin sodium technical substance equivalent to monensin activity: 25 % Perlite: 15 %-20 % Calcium carbonate q.s.100 %</p> <p><i>Active substance</i></p> <p>$C_{36}H_{61}O_{11}Na$ Sodium salt of polyether monocarboxylic acid produced by <i>Streptomyces cinnamonensis</i> 28682, LMG S-19095 in powder form.</p> <p><i>Factor composition</i></p> <p>Monensin A: not less than 90 % Monensin A + B: not less than 95 % Monensin C: 0,2-0,3 %</p> <p><i>Analytical method (1)</i></p> <p>Method for determination of the active substance: high performance liquid chromatography (HPLC) with post-column derivatization and UV-VIS (EN ISO standard method 14183:2008)</p>	Chickens reared for laying	16 weeks	100	125	<p>1. Use prohibited at least 1 day before slaughter.</p> <p>2. The additive shall be incorporated in compound feedstuffs in the form of a premixture.</p> <p>3. Monensin sodium shall not be mixed with other coccidiostats.</p> <p>4. Indicate in the instructions for use:</p> <p>'Dangerous for equines. This feedstuff contains an ionophore: avoid simultaneous administration with tiamulin and monitor for possible adverse reactions when used concurrently with other medicinal substances'.</p> <p>5. Wear suitable protective clothing, gloves and eye/face protection. In case of insufficient ventilation in the premises, wear suitable respiratory equipment.</p> <p>6. A post-market monitoring programme on the resistance to <i>Eimeria</i> spp. shall be planned and executed by the holder of authorisation.</p>	9 March 2022	<p>25 µg monensin sodium/kg of wet skin + fat</p> <p>8 µg monensin sodium/kg of wet liver, wet kidney and wet muscle</p>	

(1) Details of the analytical methods are available at the following address of the European Union Reference Laboratory for Feed Additives: http://firmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx