

PARTICULARS TO APPEAR ON <THE OUTER PACKAGE – Box label

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin 15 % w/w Flea and Tick Collar for Dogs over 12 weeks old

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Active ingredient

Dimpylate 15 % w/w
(3.6g per collar of 24g)
(6.15g per collar of 41g)

3. PHARMACEUTICAL FORM

Collar.

4. PACKAGE SIZE

Collar weighing 24 g and 60 cm in length, for dogs.

Collar weighing 41 g and 65 cm in length, for large dogs.

5. TARGET SPECIES

Dogs.

6. INDICATION(S)

For the control of fleas and ticks on dogs for up to 5 months.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Insecticidal collar weighing 24 g and 60 cm in length, for dogs.

Insecticidal collar weighing 41 g and 65 cm in length, for large dogs.

Remove the collar from sachet/blister and fasten around the dog's neck. When fitted to the growing dog, ensure the collar is loosened as the animal grows. In the fully-grown dog, a correct fitting will allow two fingers to be inserted between the collar and the neck. Any excess length of collar extending beyond the buckle should be cut and disposed of with the domestic refuse.

The collar does not need to be removed when the animal is being bathed as the insecticidal activity remains unaffected by water. Leave in place until the collar effective life is exhausted. Replace after 5 months if reinfestation with fleas or ticks is a risk.

8. WITHDRAWAL PERIOD

Not applicable.

9. SPECIAL WARNING(S), IF NECESSARY

Do not allow children to play with the collar.

Do not allow children to suck or chew the collar.

Do not smoke, eat or drink while handling collar.

Do not sleep with pets wearing collars.

Dimpylate is an organophosphorus compound : DO NOT handle if under medical advice not to work with ANTICHOLINESTERASES.

If you feel unwell after handling the collar, seek medical advice.

Wash hands thoroughly after handling the collar.

10. EXPIRY DATE

EXP:

11. SPECIAL STORAGE CONDITIONS

Do not remove the collar from the pack until ready-to-use.

Do not store above 25 °C.

Protect from direct sunlight.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

EXTREMELY DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the collar or empty packaging.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]

AVM- GSL

For animal treatment only.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Virbac S.A.

1ère avenue - 2065 m - L.I.D.

06516 Carros Cedex

France

16. MARKETING AUTHORISATION NUMBER(S)

Vm 05653/4091

17. MANUFACTURER'S BATCH NUMBER

Batch:

No Approved Packaging

PACKAGE LEAFLET FOR:

Bob Martin 15 % w/w Flea and Tick Collar for Dogs over 12 weeks old

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Virbac S.A.
1ère avenue - 2065 m - L.I.D.
06516 Carros Cedex
France

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bob Martin 15 % w/w Flea and Tick Collar for Dogs over 12 weeks old

3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS

Active ingredient :

Dimpylate 15 % w/w
(3.6g per collar of 24g)
(6.15g per collar of 41g)

4. INDICATION(S)

For the control of fleas and ticks on dogs for up to 5 months.

5. CONTRAINDICATIONS

Do not use on puppies under 12 weeks of age.

6. ADVERSE REACTIONS

Remove the collar immediately should any signs of skin irritation or adverse reaction occur.

Remove THE collar from dog immediately should any signs of drowsiness, listlessness, TREMOR, HYPERSALIVATION or respiratory difficulties develop soon after fixing of collar.

7. TARGET SPECIES

Dogs.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Insecticidal collar weighing 24 g and 60 cm in length, for dogs.

Insecticidal collar weighing 41 g and 65 cm in length, for large dogs.

Remove the collar from sachet/blister and fasten around the dog's neck. When fitted to the growing dog, ensure the collar is loosened as the animal grows. In the fully-grown dog, a correct fitting will allow two fingers to be inserted between the collar and the neck. Any excess length of collar extending beyond the buckle should be cut and disposed of with the domestic refuse.

The collar does not need to be removed when the animal is being bathed as the insecticidal activity remains unaffected by water. Leave in place until the collar effective life is exhausted. Replace after 5 months if reinfestation with fleas or ticks is a risk.

9. ADVICE ON CORRECT ADMINISTRATION

Not applicable.

10. WITHDRAWAL PERIOD(S)

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Do not remove the collar from the pack until ready-to-use.

Do not store above 25 °C.

Protect from direct sunlight.

12. SPECIAL WARNING(S)

Special warnings for each target species

Do not allow animals to chew the collar.

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated with a suitable insecticide and vacuumed regularly.

Special precautions for use in animals

For external use only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Do not allow children to play with the collar.

Do not allow children to suck or chew the collar.

Do not smoke, eat or drink while handling collar.

Do not sleep with pets wearing collars.

Dimpylate is an organophosphorus compound : DO NOT handle if

under medical advice not to work with ANTICHOLINESTERASES.

If you feel unwell after handling the collar, seek medical advice.

Wash hands thoroughly after handling the collar.

Use during pregnancy, lactation or lay

Not for use on nursing or lactating bitches.

Remove fleas from young puppies and nursing bitches by careful use of a flea comb.

Interaction with other medicinal products and other forms of interaction

Do not use any other insecticide while the collar is being worn or within 7 days of removal.

For Animal Treatment Only

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

EXTREMELY DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the collar or empty packaging.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

15. OTHER INFORMATION>

Vm 05653/4091

AVM- GSL

Approved: 03/10/2017

