

**PARTICULARS TO APPEAR ON <THE OUTER PACKAGE> <AND> <THE IMMEDIATE PACKAGE> {NATURE/TYPE}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ticacyl Spot On Solution 744 mg for Small Dogs up to 15 kg

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

This pack contains 3 x 1ml tubes each containing: permethrin (cis:trans 25:75) 744mg.

**3. PHARMACEUTICAL FORM**

Spot On Solution

**4. PACKAGE SIZE**

3 x 1ml

**5. TARGET SPECIES**

Dogs up to 15kg

**6. INDICATION(S)**

Ticacyl Spot On Solution 744mg for small dogs under 15kg contains permethrin and is formulated to eliminate and control tick on dogs, having a persistent acaricidal efficacy up to 4 weeks. Ticacyl is also effective against fleas for up to 4 weeks

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Spot on solution

Take one tube and snip the end off the nozzle.

Avoid solution coming into direct contact with your fingers during application.

Part dog's coat to expose skin between the shoulder blades and at nape of the neck.

Squeeze the entire contents of 1 tube onto the exposed skin. Avoid applying the solution into your dog's fur and do not rub into your dog's skin as this will reduce efficacy.

**8. WITHDRAWAL PERIOD**

**N/A**

## **9. SPECIAL WARNING(S), IF NECESSARY**

Avoid the drops coming into contact with your fingers. Wash off any splashes immediately with plenty of clean water. Avoid contact with eyes. In case of accidental eye contact, rinse with clean, fresh water. If eye irritation persists, seek medical advice. Wash hands and exposed skin with soap and water after use. Users making multiple applications, such as in kennels, should wear protective gloves (neoprene or nitrile 0.3mm thickness). If you feel unwell as a result of handling this product, you should consult your doctor. It is preferable to treat during the evening. Keep away from food and drink, including animal feeding stuffs. Do not handle the area of application for 6 hours following treatment. Treated dogs should not be allowed to play or sleep with people, particularly children, for 8 hours immediately following treatment.

It is very important that treated dogs are kept away from cats for 72 hours after treatment. If a cat becomes exposed to this Dog Spot On, seek veterinary advice immediately, as this product is very poisonous to cats.

Do not allow your dog near to varnished, polished or leather surface until application site is completely dry. Take particular care around leather upholstery.

For the treatment of fleas and ticks on dogs only.

For external use only.

Extremely dangerous to fish. Extremely dangerous to fish. Do not contaminate ponds, waterways or ditches with the product or packaging. Treated dogs should not be allowed to go swimming for 12 hours after treatment.

A small number of dogs may experience a minor irritation on the skin where the solution has been applied. Should this occur, bathe your dog with a mild shampoo and consult your vet. Do not use insecticidal shampoo. If signs of disease persist or appear consult your veterinary surgeon.

## **10. EXPIRY DATE**

EXP: end

## **11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C. protect from light. Do not refrigerate. Keep out of reach of children. Do not remove the tube from the blister and packaging until required for use.

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of empty packaging and any remaining product in the household refuse.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]**

NFA-VPS

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Bob Martin (UK) Ltd

Wemberham Lane

Yatton, North Somerset, BS49 4BS, UK

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 00715/4054

**17. MANUFACTURER'S BATCH NUMBER**

BN:

Approved: 22/03/2018

