

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felpreva spot-on solution for large cats (> 5.0 – 8.0 kg)

tigolaner/emodepside/praziquantel

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

115.52 mg tigolaner/24.01 mg emodepside/96.05 mg praziquantel

3. PHARMACEUTICAL FORM

Spot-on solution

4. PACKAGE SIZE

1.18 ml

1 applicator

2 applicators

10 applicators

20 applicators

5. TARGET SPECIES

Cats

> 5.0 - 8.0 kg

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Spot-on use.

Read the package leaflet before use.

8. WITHDRAWAL PERIOD(S)

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP

11. SPECIAL STORAGE CONDITIONS

Keep the applicator in the aluminium blister in order to protect from moisture.
Keep the container in the outer carton.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Vetoquinol SA
Magny Vernois
70200 Lure
France

16. MARKETING AUTHORISATION NUMBER

Vm 06462/5018

17. MANUFACTURER’S BATCH NUMBER

Lot

POM-V

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

Blister

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felpreva > 5.0 - 8.0 kg
tigolaner, emodepside, praziquantel (EN and/or Latin)



2. NAME OF THE MARKETING AUTHORISATION HOLDER

Vetoquinol SA

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Spot-on applicator

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felpreva > 5.0 - 8.0 kg
tigolaner, emodepside, praziquantel

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

115.52 mg tigolaner, 24.01 mg emodepside, 96.05 mg praziquantel

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

1.18 ml

4. ROUTE(S) OF ADMINISTRATION



5. WITHDRAWAL PERIOD(S)

6. BATCH NUMBER

Lot

7. EXPIRY DATE

EXP

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

B. PACKAGE LEAFLET

PACKAGE LEAFLET:

Felpreva spot-on solution for small cats (1.0 - 2.5 kg)
Felpreva spot-on solution for medium cats (> 2.5 – 5.0 kg)
Felpreva spot-on solution for large cats (> 5.0 – 8.0 kg)

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder and manufacturer responsible for batch release:

Vetoquinol SA
Magny-Vernois
70200 Lure
France

Manufacturer responsible for batch release:

VETOQUINOL BOWET Sp. z o.o.
Żwirowa 140
66-400 Gorzów Wlkp.
Poland

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felpreva spot-on solution for small cats (1.0 - 2.5 kg)
Felpreva spot-on solution for medium cats (> 2.5 – 5.0 kg)
Felpreva spot-on solution for large cats (> 5.0 – 8.0 kg)
tigolaner/emodepside/praziquantel

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each spot-on applicator delivers:

Felpreva spot-on solution	Volume of unit dose [ml]	Tigolaner [mg]	Emodepside [mg]	Praziquantel [mg]
for small cats (1.0 - 2.5 kg)	0.37	36.22	7.53	30.12
for medium cats (> 2.5 – 5.0 kg)	0.74	72.45	15.06	60.24
for large cats (> 5.0 – 8.0 kg)	1.18	115.52	24.01	96.05

Excipients: 2.63 mg/ml butylhydroxyanisole (E320) and 1.10 mg/ml butylhydroxytoluene (E321) as antioxidants.

Spot-on solution.
Clear yellow to red solution.

Change in colour may occur during storage. This phenomenon does not affect product quality.

4. INDICATION(S)

For cats with, or at risk from, mixed parasitic infections/ infestations. The veterinary medicinal product is exclusively indicated when ectoparasites, cestodes and nematodes are targeted at the same time.

Ectoparasites

- For the treatment of flea (*Ctenocephalides felis*) and tick (*Ixodes ricinus*, *Ixodes holocyclus*) infestations in cats providing immediate and persistent killing activity for 13 weeks.
- The veterinary medicinal product can be used as part of a treatment strategy for the control of flea allergy dermatitis (FAD).
- For the treatment of mild to moderate cases of notoedric mange (caused by *Notoedres cati*).
- For the treatment of ear mite infestations (*Otodectes cynotis*)

Gastrointestinal roundworms (nematodes)

For the treatment of infections with:

- *Toxocara cati* (mature adult, immature adult, L4 and L3)
- *Toxascaris leonina* (mature adult, immature adult and L4)
- *Ancylostoma tubaeforme* (mature adult, immature adult and L4)

Lungworms (nematodes)

For the treatment of infections with:

- *Aelurostrongylus abstrusus* (adult)
- *Troglostrongylus brevior* (adult)

Tapeworms (cestodes)

For the treatment of tapeworm infections:

- *Dipylidium caninum* (mature adult and immature adult)
- *Taenia taeniaeformis* (adult)

5. CONTRAINDICATIONS

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

6. ADVERSE REACTIONS

Mild and transient application site reactions following administration such as scratching, erythema, thinning of hair and/or inflammation may occur in very rare cases. Cosmetic effects such as a temporary spiking of the hair at the application site are commonly observed.

Neurological disorders such as ataxia and tremor may occur in very rare cases. Mild and transient digestive tract disorders such as hypersalivation or vomiting may occur in very rare cases. These effects are thought to occur as a result of the cat licking the application site immediately after treatment. In individual cases these

signs can be accompanied by non-specific signs such as agitation, vocalisation or inappetence.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

7. TARGET SPECIES

Cats.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Spot-on use. For external use only.

Posology

The recommended minimum doses are 14.4 mg tigolaner / kg body weight, 3 mg emodepside / kg body weight, 12 mg praziquantel / kg body weight equivalent to 0.148 ml veterinary medicinal product / kg body weight.

Body Weight of Cat (kg)	Applicator size to be used: Felpreva spot-on solution	Volume of unit dose (ml)	Tigolaner (mg/kg bw)	Emodepside (mg/kg bw)	Praziquantel (mg/kg bw)
1.0 - 2.5	for small cats	0.37	14.5 – 36.2	3.0 - 7.5	12.0 – 30.1
2.6 - 5.0	for medium cats	0.74	14.5 – 27.9	3.0 – 5.8	12.0 – 23.2
5.1 - 8.0	for large cats	1.18	14.4 – 22.7	3.0 - 4.7	12.0 - 18.8
> 8.0	Use an appropriate combination of applicators				

Treatment schedule

Treatment is only indicated when ectoparasites, cestodes and nematodes are targeted at the same time. In the absence of mixed infections or risk of mixed infections, appropriate narrow spectrum antiparasitic products should be used.

Fleas and ticks

The veterinary medicinal product remains active against fleas and ticks for a period of 13 weeks.

If re-treatment is necessary within 13 weeks after administration, an appropriate narrow-spectrum product should be used.

Mites

For the treatment of ear mites (*Otodectes cynotis*) and notoedric mange (*Notoedres cati*) a single dose of the veterinary medicinal product should be administered.

The treatment success and the need for re-treatment with an appropriate narrow-spectrum antiparasitic product should be determined by the treating veterinarian after 4 weeks.

Due to individual cases of single surviving ear mites and, therefore, the risk of a new cycle of otocariosis, the treatment success should be confirmed by the veterinarian 1 month after treatment.

Gastrointestinal nematodes and tapeworms

For the treatment of roundworms and tapeworms a single dose of the veterinary medicinal product should be administered. The need for and frequency of re-treatment should be in accordance with the advice of the prescribing veterinarian and take into account the local epidemiological situation as well as the cat's lifestyle. If a re-treatment is necessary within 3 months after administration, an appropriate narrow-spectrum product should be used.

Lungworms

For treatment against the lungworm *Aelurostrongylus abstrusus* and *Troglostrongylus brevior*, one treatment with the product followed by a second treatment two weeks apart with a spot-on solution for cats containing 21.4 mg/ml emodepside and 85.8 mg/ml praziquantel is recommended as there is no veterinary medicinal product containing only emodepside as active substance.

9. ADVICE ON CORRECT ADMINISTRATION

Use scissors (1) to open the childproof blister. Pull foils apart (2) and remove spot-on applicator from package (3).



Hold applicator in upright position (4), twist and pull off cap (5) and use the opposite end of the cap to break the seal (6).



Part the fur on the cat's neck at the base of the skull until the skin is visible (7). Place the tip of the applicator on the skin and squeeze firmly several times to empty the contents directly onto the skin (7). Application on the base of the skull will minimise the ability of the cat to lick the product off.



10. WITHDRAWAL PERIOD(S)

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature storage conditions.

Keep the applicator in the aluminium blister in order to protect from moisture.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton. The expiry date refers to the last day of that month.

12. SPECIAL WARNING(S)

Special warnings for each target species:

Ectoparasites need to start feeding on the host to become exposed to tigolaner; therefore, the risk of the transmission of vector borne diseases cannot be excluded.

Parasite resistance to any particular class of antiparasitics included in the fixed combination may develop following frequent, repeated use of antiparasitics of those classes under specific circumstances. The use of this veterinary medicinal product should be based on the assessment of each individual case and on local epidemiological information about the current susceptibility of the target species in order to limit the possibility of a future selection for resistance.

Shampooing or immersion of the animal in water directly after treatment may reduce the efficacy of the product. Treated animals, therefore, should not be bathed until the solution has dried.

Special precautions for use in animals:

In the absence of available data, treatment of kittens under 10 weeks of age or weighing less than 1 kg is not recommended.

This veterinary medicinal product is for topical use and should not be administered in any other way e.g. orally.

Apply only onto intact skin. Apply as described under "advice on correct administration" to prevent the animal from licking and ingesting the veterinary

medicinal product. Avoid the treated cat or other cats in the household licking the site of application while it is wet. For signs observed after oral ingestion (e.g. licking) please refer to section “adverse reactions”.

The product may be an eye irritant. If accidental ocular contact occurs, flush the eyes immediately with clean water. If eye irritation occurs, seek veterinary advice.

There is no experience on the use of the veterinary medicinal product in sick and debilitated animals, thus the veterinary medicinal product should only be used based on a benefit-risk assessment for these animals.

Acute signs of pneumonia may occur after treatment as a result of the inflammatory host response against the death of *T. brevior* lungworms especially in young cats.

The veterinary medicinal product should not be administered at intervals shorter than 8 weeks. Due to the product’s activity against fleas and ticks for a period of 3 months from a clinical point of view, the use of the product is not indicated at intervals shorter than three months.

No target animal safety data beyond 4 consecutive treatments are available and accumulation of tigolaner is likely. Repeated treatments should be restricted to limited individual situations according to a benefit-risk evaluation by the responsible veterinarian. Please refer to section “Overdose”.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The veterinary medicinal product may cause neurological symptoms and can transiently elevate blood glucose levels following accidental ingestion.

Do not smoke, eat or drink during application.

Wash hands after use.

Used applicators should be disposed of immediately and should not be left within the sight or reach of children.

In case of accidental contact of the applicator content to skin, wash off immediately with soap and water.

The veterinary medicinal product may be an eye irritant. If the veterinary medicinal product accidentally gets into eyes, they should be thoroughly flushed with plenty of water.

If skin or eye symptoms persist, or in case of accidental ingestion, especially by children, seek medical advice immediately and show the package leaflet or the label to the physician.

Since foetotoxic effects are described in laboratory animals after exposure to tigolaner and emodepside, pregnant women and women intending to conceive should wear gloves to avoid direct contact with the product.

Pregnant women should avoid contact with the site of application during the first 24 hours after application of the product and until the treated area is no longer noticeable. Keep children away from treated animals during the first 24 hours after application of the product . Care should be taken not to allow children to have prolonged intensive contact with treated cats until the treated area is no longer noticeable. It is recommended to treat animals in the evening. On the day of treatment, treated animals

should not be permitted to sleep in the same bed as their owner, especially children and pregnant women.

The veterinary medicinal product may stain or damage certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Pregnancy and lactation:

Foetotoxic effects are described in laboratory animals after exposure to tigolaner and emodepside. The safety of the veterinary medicinal product has not been established in pregnant or lactating cats and, therefore, use in such animals is not recommended.

Interaction with other medicinal products and other forms of interaction:

Emodepside is a substrate for P-glycoprotein. Co-treatment with other substances that are P-glycoprotein substrates/inhibitors (for example, ivermectin and other antiparasitic macrocyclic lactones, erythromycin, prednisolone and cyclosporine) could give rise to pharmacokinetic drug interactions.

Overdose (symptoms, emergency procedures, antidotes):

After administration of 4 consecutive treatments of up to 5 times the maximum recommended dose in kittens starting at 10 weeks of age and adult cats, a decrease in thyroid weight was noted in some male animals. In adult cats, a transient elevation in liver enzymes (AST, ALT), accompanied by multifocal liver congestion in one individual, was noted in the high dose group (5x) and an elevation in cholesterol in all overdose groups (3x, 5x). No systemic clinical signs were observed.

In the high (5x) dose group, cases of local reactions at the application site occurred (alopecia, erythema, hyperplasia of the epidermis and/or inflammatory infiltrates).

There is no antidote known.

For Animal Treatment Only

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

The product should not enter water courses as this may be dangerous for fish and other aquatic organisms.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

14. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

15. OTHER INFORMATION

Vm 06462/5016
Vm 06462/5017
Vm 06462/5018

POM-V

Pack sizes: 1, 2, 10 or 20 applicators. Not all pack sizes may be marketed.

Tigolaner belongs to the chemical class of bispyrazoles. Tigolaner acts as a potent inhibitor of the neurotransmitter gamma-aminobutyric acid (GABA) receptor. Tigolaner exhibits higher functional potency to block insect/acarine receptors compared to mammalian receptors. It is an acaricide and insecticide and is efficacious against ticks (*Ixodes 15eonine*, *I. holocyclus*), fleas (*Ctenocephalides felis*), and mites (*Notoedres cati*, *Otodectes cynotis*) on cats. Fleas already on the animal prior to administration are killed within 12 hours. For newly infecting fleas the onset of efficacy is within 8 hours for 2 months after product administration and within 24 hours afterwards. Fleas and ticks must attach to the host and start feeding in order to be exposed to tigolaner. *Ixodes 15eonine* ticks on the animal prior to administration are killed within 24 hours. *Ixodes 15eonine* ticks newly infested are killed within 48 hours for 13 weeks.

Emodepside is a semi-synthetic compound belonging to the chemical group of depsipeptides. It is active against all stages of roundworms (ascarids and hookworms). In this product, emodepside is responsible for the efficacy against *Toxocara cati*, *Toxascaris 15eonine*, *Ancylostoma tubaeforme*, *Aelurostrongylus abstrusus* and *Troglostrongylus brevior*.

It acts at the neuromuscular junction by stimulating presynaptic receptors belonging to the secretin receptor family which results in paralysis and death of the parasites.

Praziquantel is a pyrazinoisoquinoline derivative effective against the tapeworms *Dipylidium caninum*, and *Taenia taeniaeformis*.

Praziquantel is rapidly adsorbed via the surface of the parasites and acts primarily by changing the Ca⁺⁺ permeability of the parasite membranes. This results in severe damage to the parasite integument, contraction and paralysis, disruption of metabolism and finally leads to the death of the parasite.

[Read me information to be placed on the top of the package leaflet]

Dear cat-owner,
Your cat has been prescribed Felpreva, an approved veterinary medicinal product for cats. This leaflet contains valuable information for application and usage of Felpreva. Please read this leaflet carefully and follow the instructions.

Gavin Hall
Approved: 19 January 2025