PARTICULARS TO APPEAR ON THE OUTER PACKAGE
{BOX}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

RSPCA FLEAaway Combo 50 mg/60 mg Spot-on Solution

2. STATEMENT OF ACTIVE SUBSTANCES

WHAT'S IN THIS PRODUCT?
Each 0.5 ml pipette contains:

Active substances:
50 mg fipronil
60 mg S-methoprene

3. PACKAGE SIZE

1 x 0.5 ml
3 x 0.5 ml
6 x 0.5 ml

4. TARGET SPECIES

Cats

5. INDICATION(S)

- Kills fleas, ticks and lice
- Protects for up to 4 weeks against fleas

WHAT DOES THIS PRODUCT DO?
Kills fleas (*Ctenocephalides spp.*) on treated cats for up to 4 weeks and prevents eggs laid by exposed fleas from developing for up to 6 weeks. Kills ticks (*Ixodes ricinus, Dermacentor variabilis, Rhipicephalus sanguineus*) on treated cats for up to 2 weeks. Treats infestations with biting lice (*Felicola subrostratus*).

6. ROUTE(S) OF ADMINISTRATION

How to apply this product



1 x Treatment
For external use only.

HOW DO I USE THIS PRODUCT?
Before use, read the package leaflet for full instructions and all warnings, including user warnings.

1. Hold the pipette upright. Twist and pull the cap off.
2. Use the tip of the cap to break the pipette's seal.
3. Part the coat on the back of the animal's neck in front of the shoulder blades until the skin is visible.
4. Apply the pipette to the skin and squeeze it several times to empty all of its contents onto the skin in one spot.

Do not reapply more often than at four weekly intervals.
It is important to make sure that the product is applied to an area where the animal cannot lick it off and to make sure that animals do not lick each other following treatment.

7. WITHDRAWAL PERIOD

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

HOW SHOULD I STORE THE PRODUCT
Store in the original package in order to protect from light and moisture.

10. THE WORDS "READ THE PACKAGE LEAFLET BEFORE USE"

Read the package leaflet before use.

IS THIS PRODUCT SUITABLE FOR MY PET?
Only use this product if your cat is at least 8 weeks old and weighs more than 1 kg.
Do not use if your pet is sick or recovering from an illness. Do not use if your pet has a hypersensitivity (allergy) to fipronil or S-methoprene or the other ingredients. Do not apply on wounds or damaged skin.
Only use on cats, do not use in any other animal species.
Do not use on rabbits as adverse drug reactions, including death, may occur.

USER SAFETY- Please read before every use.
This veterinary medicinal product can cause mucous membrane, skin and eye irritation. Therefore, contact of the veterinary medicinal product with mouth, skin and eyes should be avoided. People with a known hypersensitivity (allergy) to fipronil, S-methoprene, or alcohol should avoid contact with the veterinary medicinal product. Avoid contents coming into contact with the fingers. If the veterinary medicinal product comes into contact with fingers, wash hands with soap and water. After accidental eye contact, rinse carefully in clean water. If the veterinary medicinal product is accidentally swallowed, seek medical advice immediately and show the package leaflet or the label to the physician.
Wash hands after use. Do not smoke, drink or eat during application.
Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals are not allowed to sleep with owners, especially children.
Keep pipettes in original packaging until ready to use and dispose of used pipettes immediately.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto

14. MARKETING AUTHORISATION NUMBERS

Vm 01656/3008
Vm 01656/5008

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

{BAG}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

RSPCA FLEAaway Combo

2. QUANTITATIVE PARTICULARS OF THE ACTIVE

Fipronil/ S-methoprene
50 mg/60 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Marketing authorisation holder
KRKA



MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

{PIPETTE}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

RSPCA FLEAaway Combo

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

50 mg/60 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

RSPCA FLEAaway Combo 50 mg/60 mg Spot-on Solution for Cats

2. Composition

Each 0.5 ml pipette contains:

Active substances:

Fipronil	50 mg
S-methoprene	60 mg

Excipients:

Butylhydroxyanisole (E320)	0.1 mg
Butylhydroxytoluene (E321)	0.05 mg

Clear yellow solution.

3. Target species

Cats

4. Indication(s)

To be used against infestations with fleas, alone or in association with ticks and/or biting lice:

- Treatment of flea infestations (*Ctenocephalides* spp.). Insecticidal efficacy against new infestations with adult fleas persists for 4 weeks. Prevention of the multiplication of fleas by inhibiting the development of eggs (ovicidal activity), larvae and pupae (larvicidal activity) originating from eggs laid by adult fleas for six weeks after application.
- Treatment of tick infestations (*Ixodes ricinus, Dermacentor variabilis, Rhipicephalus sanguineus*). The product has a persistent acaricidal efficacy for up to 2 weeks against ticks.
- Treatment of infestations with biting lice (*Felicola subrostratus*).

5. Contraindications

Do not use on kittens less than 8 weeks old and/or weighing less than 1 kg.
Do not use on sick (e.g. systemic diseases, fever) or convalescent animals.

Do not use if your cat has a hypersensitivity (allergy) to fipronil or S-methoprene or the other ingredients.
Only use in cats, do not use in any other animal species.
Do not use on rabbits as adverse drug reactions, including death, may occur.

6. Special warnings

Special warnings:
When treating infestations of parasites, all in-contact animals should be treated with an appropriate product at the same time. Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.
Bathing and/or shampooing should be avoided. No information on the effect of bathing or the use of a shampoo on the effectiveness of the product in cats is available. Based on information available for dogs, bathing/immersion in water within 2 days after application of the product and more frequent bathing than once a week should be avoided.

Special precautions for safe use in the target species:
Avoid contact with the animal’s eyes.
It is important to make sure that the product is applied to an area where the animal cannot lick it off and to make sure that animals do not lick each other following treatment.
There may be an attachment of single ticks. For this reason a transmission of infectious diseases cannot be completely excluded if conditions are unfavourable.
The potential toxicity of the product for kittens of less than 8 weeks of age in contact with a treated queen is not documented. Special care should be taken in this case.

Special precautions to be taken by the person administering the veterinary medicinal product to animals
This veterinary medicinal product can cause mucous membrane, skin and eye irritation. Therefore, contact of the veterinary medicinal product with mouth, skin and eyes should be avoided.
People with a known hypersensitivity (allergy) to fipronil, S-methoprene, or alcohol should avoid contact with the veterinary medicinal product. Avoid contents coming into contact with the fingers. If the veterinary medicinal product comes into contact with fingers, wash hands with soap and water.
After accidental eye contact, rinse carefully with clean water.
If the veterinary medicinal product is accidentally swallowed, seek medical advice immediately and show the package leaflet or the label to the physician.
Wash hands after use.
Do not smoke, drink or eat during application.
Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals are not allowed to sleep with owners, especially children.
Keep pipettes in original packaging until ready to use and dispose of used pipettes immediately.

Pregnancy and lactation:
The product can be used during pregnancy.
The potential toxicity of the product for kittens of less than 8 weeks of age in contact with a treated queen is not documented. Special care should be taken in this case.

Interaction with other medicinal products and other forms of interaction:
Do not use simultaneously with other flea products which are applied directly on to the animal.

Overdose:
No undesirable effects were observed in target animal safety studies in cats and kittens aged 8 weeks and older and weighing about 1 kg treated once a month at five times the recommended dose for six consecutive months.
Itching may occur following treatment.
Overdose application of the product will cause a sticky appearance of hairs at the treatment spot. However, should this occur, it will disappear within 24 hours post application.
The risk of experiencing adverse effects may increase with overdosing (see section 6 Adverse reactions).

7. Adverse events

Do not overdose.

Cat:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site skin scaling, hair loss, itching, reddening* Generalised itching or hair loss
Undetermined frequency (cannot be estimated from the available data)	Increased salivation Vomiting Increased sensitivity to stimulation, depression, other nervous signs

*Transient.

In the case of licking the administration site, a brief period of excessive salivation may be observed due mainly to the nature of the carrier.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medical product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or its local representative using the contact details at the end of this leaflet, or via your national reporting system:
Website: https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine
e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, route(s) and method of administration

For spot-on administration. Apply one pipette of 0.5 ml per cat by topical application to the skin, corresponding to a minimum recommended dose of 5 mg/kg for fipronil and 6 mg/kg for S-methoprene.

Method of administration:

1. Remove the pipette from its packaging. Hold the pipette in an upright position, twist and pull the cap off.
2. Turn the cap around and place the other end of the cap back on the pipette. Push and twist the cap to break the seal, then remove the cap from the pipette.
3. Part the coat on the back of the animal at the base of the neck in front of the shoulder blades until the skin is visible.
4. Place the tip of the pipette onto the skin and squeeze the pipette several times to empty its contents completely and directly onto the skin in one spot.



9. Advice on correct administration

In the absence of safety studies, the minimum treatment interval is 4 weeks.

10. Withdrawal period(s)

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature storage conditions. Store in the original package in order to protect from light and moisture.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after Exp. The expiry date refers to the last day of that month.

12. Special precautions for the disposal

Medicines should not be disposed of via wastewater.
This veterinary medicinal product should not enter water courses as Fipronil and S-methoprene may be dangerous for fish and other aquatic organisms.

Use take-back schemes for the disposal of any unused veterinary medicinal product

waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Do not contaminate ponds, waterways or ditches with the veterinary medicinal product or empty container.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product not subject to prescription.

14. Marketing authorisation numbers and pack sizes

Vm 01656/3008
Vm 01656/5008

Available in cartons containing 1, 3 or 6 pipettes.
Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release
KRKA, d.d., Novo mesto
Šmarješka cesta 6
8501 Novo mesto
Slovenia

Local representatives and contact details to report suspected adverse reactions:
KRKA UK Ltd
United Kingdom
Tel: 02071 646 156
info.uk@krka.biz

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information

AVM-GSL

Gavin Hall

Approved: 16 June 2026