

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE {CARDBOARD BOX}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

IsoFlo 100% w/w Inhalation Vapour, liquid

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Isoflurane 1000 mg/g

**3. PACKAGE SIZE**

250 ml

100 ml

**4. TARGET SPECIES**

Horses, dogs, cats, ornamental birds, reptiles, rats, mice, hamsters, chinchillas, gerbils, guinea pigs and ferrets.

**5. INDICATION(S)**

**6. ROUTES OF ADMINISTRATION**

Inhalation use.

**7. WITHDRAWAL PERIODS**

Withdrawal period: Horses: Meat and offal: 2 days.

Not authorised for use in mares producing milk for human consumption.

**8. EXPIRY DATE**

Exp. {mm/yyyy}

**9. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Store in the original bottle.

Keep the bottle in the outer carton

Keep the bottle tightly closed.

Protect from direct sunlight and heat.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

## **13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

## **14. MARKETING AUTHORISATION NUMBERS**

Vm 42058/5102

## **15. BATCH NUMBER**

Lot {number}

## **16. SPECIAL WARNING(S), IF NECESSARY**

Anaesthetics must be handled correctly. **Read the package leaflet before use.**

## **17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

## **18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

**POM-V**

To be supplied only on veterinary prescription.

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {BOTTLE LABEL}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

IsoFlo 100% w/w Inhalation Vapour, liquid

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Isoflurane 1000 mg/g

**3. TARGET SPECIES**

Horses, dogs, cats, ornamental birds, reptiles, rats, mice, hamsters, chinchillas, gerbils, guinea pigs and ferrets.

**4. ROUTES OF ADMINISTRATION**

Inhalation use.  
Read the package leaflet before use.

**5. WITHDRAWAL PERIODS**

Withdrawal period: Horses: Meat and offal: 2 days  
Not authorised for use in mares producing milk for human consumption.

**6. EXPIRY DATE**

Exp. {mm/yyyy}

**7. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.  
Store in the original bottle.  
Keep the bottle in the outer carton  
Keep the bottle tightly closed.  
Protect from direct sunlight and heat.

**8. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**9. BATCH NUMBER**

Lot {number}

## **10. PACKAGE SIZE**

250 ml

100 ml

## **11. INDICATION(S)**

## **12. SPECIAL WARNING(S), IF NECESSARY**

Anaesthetics must be handled correctly. Read the package leaflet before use.

## **13. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

## **14. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

POM-V

## **15. MARKETING AUTHORISATION NUMBER(S)**

Vm 42058/5102

## **PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

IsoFlo 100% w/w Inhalation Vapour, liquid

### **2. COMPOSITION**

Each g contains:

#### **Active substance:**

Isoflurane 1000 mg

Clear, colourless volatile liquid.

### **3. TARGET SPECIES**

Horse, dog, cat, ornamental bird, reptile, rat, mouse, hamster, chinchilla, gerbil, guinea pig and ferret.

### **4. INDICATIONS FOR USE**

Induction and maintenance of general anaesthesia.

### **5. CONTRAINDICATIONS**

Do not use in cases of known susceptibility to malignant hyperthermia.  
Do not use in cases of hypersensitivity to the active substance.

### **6. SPECIAL WARNING(S)**

For animal treatment only.

#### Special warnings:

The ease and rapidity of alteration of the depth of anaesthesia with isoflurane and its low metabolism, may be considered advantageous for its use in special groups of patients such as the old or young, and those with impaired hepatic, renal or cardiac function.

#### Special precautions for safe use in the target species:

Isoflurane has little or no analgesic properties. Adequate analgesia should always be given before surgery. The analgesic requirements of the patient should be considered before general anaesthesia is ended.

Isoflurane causes depression of the cardiovascular and respiratory systems. It is important to monitor pulse quality and rate in all patients. The use of the product in patients with cardiac disease should only be considered after a benefit risk assessment by the responsible veterinary surgeon. In the case of cardiac arrest, complete cardiopulmonary resuscitation should be performed.

It is important to monitor respiratory rate and quality. It is also important to maintain an open airway and to properly oxygenate tissues during the maintenance of anaesthesia. Respiratory arrest should be treated by assisted ventilation.

The metabolism of isoflurane in birds, and small mammals, can be affected by decreases in body temperature, that may occur secondary to a high surface area to body weight ratio. Furthermore, drug metabolism in reptiles is slow and highly dependent upon environmental temperature. Therefore, body temperature should be monitored and kept stable during treatment.

Reptiles may be difficult to induce with inhalation agents due to breath holding. When using isoflurane to anaesthetise an animal with a head injury, consideration should be given as to whether artificial ventilation is appropriate to help avoid increased cerebral blood flow by maintaining normal CO<sub>2</sub> levels.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not breathe the vapour. Users should consult their National Authority for advice on Occupational Exposure Standards for isoflurane.

Operating rooms and recovery areas should be provided with adequate ventilation or scavenging systems to prevent the accumulation of anaesthetic vapour. All scavenging/ extraction systems must be adequately maintained.

Adverse effects on foetuses and pregnant animals were observed in laboratory animals. Pregnant and breast-feeding women should not have any contact with the product and should avoid operating rooms and animal recovery areas. Avoid using masking procedures for prolonged induction and maintenance of general anaesthesia.

Use cuffed endotracheal intubation when possible for the administration of IsoFlo during maintenance of general anaesthesia.

Care should be taken when dispensing isoflurane, with any spillage removed immediately using an inert and absorbent material e.g. sawdust. Wash any splashes from skin and eyes and avoid contact with the mouth. If severe accidental exposure occurs remove the operator from the source of exposure, seek medical advice immediately and show the package leaflet or the label to the physician.

Halogenated anaesthetic agents may induce liver damage. In the case of isoflurane this is an idiosyncratic response very rarely seen after repeated exposure.

*To the physician:* Ensure a patient's airway and give symptomatic and supportive treatment. Note that adrenaline and catecholamines may cause cardiac dysrhythmias.

Special precautions for the protection of the environment:

Although anaesthetics have a low potential for damage to the atmosphere, it is good practice to use charcoal filters with scavenging equipment, rather than to discharge them into the air.

Pregnancy:

Use only according to the benefit/risk assessment by the responsible veterinarian. Isoflurane has been safely used for anaesthesia during caesarean section in the dog and the cat.

Lactation:

Use only according to the benefit/risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

The action of muscle relaxants in man, especially those of the non-depolarising type such as atracurium, pancuronium or vecuronium is enhanced by isoflurane. Similar potentiation might be expected to occur in the target species, although there is little

direct evidence to this effect. Concurrent inhalation of nitrous oxide enhances the effect of isoflurane in man and similar potentiation might be expected in animals. The concurrent use of sedative or analgesic drugs is likely to reduce the level of isoflurane required to produce and maintain anaesthesia. Isoflurane has a weaker sensitising action on the myocardium, to the effects of circulating dysrhythmogenic catecholamines, than halothane. Isoflurane may be degraded to carbon monoxide by dried carbon dioxide absorbents.

#### Overdose:

Isoflurane overdose may result in profound respiratory depression. Therefore, respiration must be monitored closely and supported when necessary with supplementary oxygen and/ or assisted ventilation.

In cases of severe cardiopulmonary depression, administration of isoflurane should be discontinued, the breathing circuit should be flushed with oxygen, the existence of a patent airway ensured, and assisted or controlled ventilation with pure oxygen initiated. Cardiovascular depression should be treated with plasma expanders, pressor agents, antiarrhythmic agents or other appropriate techniques.

#### Major incompatibilities:

Isoflurane has been reported to interact with dry carbon dioxide absorbents to form carbon monoxide. In order to minimise the risk of formation of carbon monoxide in rebreathing circuits and the possibility of elevated carboxyhaemoglobin levels, carbon dioxide absorbents should not be allowed to dry out.

## **7. ADVERSE EVENTS**

Horse, dog, cat, ornamental bird, reptile, rat, mouse, hamster, chinchilla, gerbil, guinea pig and ferret.

Rare (1 to 10 animals / 10,000 animals treated)	bradycardia <sup>1</sup> , arrhythmias
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	cardiac arrest, hypotension <sup>2</sup> , respiratory arrest, respiratory depression <sup>2</sup> , malignant hyperthermia <sup>3</sup>

<sup>1</sup> Transient.

<sup>2</sup> Dose-related.

<sup>3</sup> Susceptible animals.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or its local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Inhalation use.

The MAC (minimal alveolar concentration in oxygen) or effective dose ED50 and suggested concentrations given below for the target species should be used as a guide or starting point only. The actual concentrations required in practice will depend on many variables including the concomitant use of other drugs and the clinical status of the patient.

Isoflurane may be used in conjunction with other drugs commonly used in veterinary anaesthetic regimes for premedication, induction and analgesia. Some specific examples are given in the individual species information. The use of analgesia for painful procedures is consistent with good veterinary practice.

Recovery from isoflurane anaesthesia is usually smooth and rapid. The analgesic requirements of the patient should be considered before the termination of general anaesthesia.

## **HORSES**

The MAC for isoflurane in the horse is approximately 1.31%.

### **Premedication:**

Isoflurane may be used with other drugs commonly used in veterinary anaesthetic regimes. The following drugs have been found to be compatible with isoflurane: acepromazine, alfentanil, atracurium, butorphanol, detomidine, diazepam, dobutamine, dopamine, [guaiphenesin](#), ketamine, morphine, pentazocine, pethidine, thiamylal, thiopentone and xylazine. Drugs used for premedication should be selected for the individual patient. However, the potential interactions below should be noted.

### **Interactions:**

Detomidine and xylazine have been reported to reduce the MAC for isoflurane in horses.

### **Induction:**

As it is not normally practicable to induce anaesthesia in adult horses using isoflurane, induction should be by the use of a short acting barbiturate such as thiopentone sodium, ketamine or [guaiphenesin](#). Concentrations of 3 to 5% isoflurane may then be used to achieve the desired depth of anaesthesia in 5 to 10 minutes. Isoflurane at a concentration of 3 to 5% in a high flow of oxygen may be used for induction in foals.

### **Maintenance:**

Anaesthesia may be maintained using 1.5% to 2.5% isoflurane.

### **Recovery:**

Recovery is usually smooth and rapid.

## **DOGS**

The MAC for isoflurane in the dog is approximately 1.28%.

### **Premedication:**

Isoflurane may be used with other drugs commonly used in veterinary anaesthetic regimes. The following drugs have been found to be compatible with isoflurane: acepromazine, atropine, butorphanol, buprenorphine, bupivacaine, diazepam,



dobutamine, ephedrine, epinephrine, etomidate, glycopyrrolate, ketamine, medetomidine, midazolam, methoxamine, oxymorphone, propofol, thiamylal, thiopentone and xylazine. Drugs used for premedication should be selected for the individual patient. However, the potential interactions below should be noted.

Interactions:

Morphine, oxymorphone, acepromazine, medetomidine, medetomidine plus midazolam have been reported to reduce the MAC for isoflurane in dogs. The concomitant administration of midazolam/ketamine during isoflurane anaesthesia may result in marked cardiovascular effects, particularly arterial hypotension. The depressant effects of propranolol on myocardial contractility are reduced during isoflurane anaesthesia, indicating a moderate degree of  $\beta$ -receptor activity.

Induction:

Induction is possible by face mask using up to 5% isoflurane, with or without premedication.

Maintenance:

Anaesthesia may be maintained using 1.5% to 2.5% isoflurane.

Recovery:

Recovery is usually smooth and rapid.

**CATS**

The MAC for isoflurane in the cat is approximately 1.63%.

Premedication:

Isoflurane may be used with other drugs commonly used in veterinary anaesthetic regimes. The following drugs have been found to be compatible with isoflurane: acepromazine, atracurium, atropine, diazepam, ketamine, and oxymorphone. Drugs used for premedication should be selected for the individual patient. However, the potential interactions below should be noted.

Interactions:

Intravenous administration of midazolam-butorphanol has been reported to alter several cardio-respiratory parameters in isoflurane-induced cats as has epidural fentanyl and medetomidine.

Isoflurane has been shown to reduce sensitivity of the heart to adrenalin (epinephrine).

Induction:

Induction is possible by face mask using up to 4% isoflurane, with or without premedication.

Maintenance:

Anaesthesia may be maintained using 1.5% to 3% isoflurane.

Recovery:

Recovery is usually smooth and rapid.

## **ORNAMENTAL BIRDS**

Few MAC/ED<sub>50</sub> values have been recorded. Examples are 1.34% for the Sandhill crane, 1.45% for the racing pigeon, reduced to 0.89% by the administration of midazolam, and 1.44% for cockatoos, reduced to 1.08% by the administration of butorphanol analgesic.

The use of isoflurane anaesthesia has been reported for many species, from small birds such as zebra finches, to large birds such as vultures, eagles and swans.

### Drug interactions/compatibilities:

Propofol has been demonstrated in the literature to be compatible with isoflurane anaesthesia in swans.

### Interactions:

Butorphanol has been reported to reduce the MAC for isoflurane in cockatoos. Midazolam has been reported to reduce the MAC for isoflurane in pigeons.

### Induction:

Induction with 3 to 5% isoflurane is normally rapid. Induction of anaesthesia with propofol, followed by isoflurane maintenance, has been reported for swans.

### Maintenance:

The maintenance dose depends on the species and individual. Generally, 2 to 3% is suitable and safe.

Only 0.6 to 1% may be needed for some stork and heron species.

Up to 4 to 5% may be needed for some vultures and eagles.

3.5 to 4% may be needed for some ducks and geese.

Generally, birds respond very rapidly to changes in concentration of isoflurane.

### Recovery:

Recovery is usually smooth and rapid.

## **REPTILES**

Isoflurane is considered by several authors to be the anaesthetic of choice for many species. The literature records its use on a wide variety of reptiles (eg. various species of lizard, tortoise, iguanas, chameleon and snakes).

The ED<sub>50</sub> was determined in the desert iguana to be 3.14% at 35°C and 2.83% at 20°C.

### Drug interactions/ compatibilities:

No specific publications on reptiles have reviewed compatibilities or interactions of other drugs with isoflurane anaesthesia.

### Induction:

Induction is usually rapid at 2 to 4% isoflurane.

### Maintenance:

1 to 3% is a useful concentration.

### Recovery:

Recovery is usually smooth and rapid.

## **RATS, MICE, HAMSTERS, CHINCHILLAS, GERBILS, GUNIEA PIGS AND FERRETS**

Isoflurane has been recommended for anaesthesia of a wide variety of small mammals.

The MAC for mice has been cited as 1.34%, and for the rat as 1.38%, 1.46% and 2.4%.

### Drug interactions/ compatibilities:

No specific publications on small mammals have reviewed compatibilities or interactions of other drugs with isoflurane anaesthesia.

### Induction:

Isoflurane concentration 2 to 3%.

### Maintenance:

Isoflurane concentration 0.25 to 2%.

### Recovery:

Recovery is usually smooth and rapid.

## **SUMMARY TABLE CONTAINING ANAESTHESIA INDUCTION AND MAINTENANCE DATA BY SPECIES**

Species	MAC (%)	Induction (%)	Maintenance (%)	Recovery
Horses	1.31	3.0 – 5.0	1.5 – 2.5	Smooth and rapid
Dogs	1.28	Up to 5.0	1.5 – 2.5	Smooth and rapid
Cats	1.63	Up to 4.0	1.5 – 3.0	Smooth and rapid
Ornamental birds	See posology	3.0 – 5.0	See posology	Smooth and rapid
Reptiles	See posology	2.0 – 4.0	1.0 – 3.0	Smooth and rapid
Rats, mice, hamsters, chinchillas, gerbils, guinea pigs and ferrets	1.34 (mouse) 1.38/1.46/2.40 (rat)	2.0 – 3.0	0.25 – 2.0	Smooth and rapid

## **9. ADVICE ON CORRECT ADMINISTRATION**

Isoflurane should be administered using an accurately calibrated vaporiser in an appropriate anaesthetic circuit since levels of anaesthesia may be altered rapidly and easily. Isoflurane may be administered in oxygen or oxygen/nitrous oxide mixtures.

## **10. WITHDRAWAL PERIOD(S)**

Horses:

Meat and offal: 2 days.

Not authorised for use in mares producing milk for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not store above 25°C.

Store in the original bottle.

Keep the bottle in the outer carton.

Keep the bottle tightly closed.

Protect from direct sunlight and heat.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and the carton after Exp. The expiry date refers to the last day of that month.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater or household waste.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements. These measures should help to protect the environment..

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

## **14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

Vm 42058/5102

### Package sizes:

100 ml bottle in a cardboard box.

250 ml bottle in a cardboard box.

Not all pack sizes may be marketed.

## **15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED**

Detailed information on this veterinary medicinal product is available in the Union Product Database.

## **16. CONTACT DETAILS**

### Marketing authorisation holder:

Zoetis UK Limited

1st Floor, Birchwood Building

Springfield Drive

Leatherhead


Surrey

KT22 7LP

Manufacturer responsible for batch release:

Zoetis Belgium SA  
Rue Laid Burniat 1  
1348 Louvain-la-Neuve  
Belgium

**17. OTHER INFORMATION**

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 26 June 2023