

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {Bottle labels}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Zermasect Sheep Pour-On Alphacypermethrin 12.5 g/l pour-on solution

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Alphacypermethrin 12.5 g/l pour on solution

3. PHARMACEUTICAL FORM

Pour-On Solution

4. PACKAGE SIZE

5 Litres

5. TARGET SPECIES

Sheep

6. INDICATION(S)

For the treatment and control of blowfly strike in sheep and lambs, and for the treatment of lice and ticks on sheep.

Ready to use topical insecticidal solution. Treats a maximum of 124 ewes or a maximum of 200 lambs.

Contains the pyrethroid alphacypermethrin, which has both a repellent and insecticidal action to protect ewes and lambs from blowfly strike. The product has been formulated as a ready to use pour-on. Treated animals are easily identified by the natural green colour of the product which will fade after a few days without staining the fleece.

For the best results, apply before the spring or autumn tick rise and before the first appearance of headfly or blowfly; treat for lice when they are present. To ensure the best results the whole flock should be treated.

Use:

Blowfly – For the control of blowfly a single application will normally provide 8 to 10 weeks protection. Re-treatment may be necessary after this period.

For the treatment of blowfly strike, a single treatment applied to the infected area will ensure blowfly larvae are killed.

Ticks – A single treatment will normally provide control of ticks for 8 to 12 weeks.

Headfly – A single treatment will reduce the incidence for up to 6 weeks.

Lice – A single treatment will normally kill all lice.

7. CONTRAINDICATIONS

Avoid treating animals in hot weather. Do not treat sheep producing milk for human consumption. Do not treat lambs under one week old. Discolouration of the fleece may be observed in some sheep from approximately 6 to 8 weeks after treatment.

8. METHOD AND ROUTE(S) OF ADMINISTRATION

For external use only.

READ THE INSTRUCTIONS CAREFULLY BEFORE USE

| Pest | Dosage, ml | |
|---------|----------------|----------------|
| | Sheep (> 25kg) | Lambs (< 25kg) |
| Blowfly | 40 | 25 |
| Lice | 40 | |
| Ticks | 40 | |
| Headfly | 5 | 5 |

For dosages of 40 ml and 25ml, apply as two half-doses. Apply the first half-dose evenly along the back line from the neck to the saddle, and the second half-dose evenly from the saddle to the tail head and around the rump.

For the control of blowfly it is important that there is good coverage around the tail area. The T-bar must be held against the fleece so that the dose is applied directly into it. For treatment of existing strike, apply the dose directly to the struck area. Most strikes will require the application of 5ml (saucer size) or 10ml (dinner plate size).

For the reduction of headfly strike, 5mls per animal is applied between the horns. Because alphacypermethrin is retained in the wool grease of the fleece, it is recommended that the product be applied to sheep with a minimum wool length of 1.0cm.

9. WITHDRAWAL PERIOD

Meat: 49 days.

Milk: Not for use in sheep producing milk for human consumption.

10. SPECIAL WARNING(S), IF NECESSARY

Protective clothing must be worn. Remove the transit cap from the bottle and pierce the seal. Fit the spouted cap to the bottle. Fit the tube from the applicator gun to the spouted cap and fit the metal spring over the tube on the cap spout to ensure it is fixed securely. Check the gun is fitted with the T-bar and that it is working correctly.

Before use the gun must be primed and set to the required volume. Clean the gun after use and maintain the gun to ensure accurate dosing.

The SimcroTech 25ml Pour-On Applicator gun is recommended as the product may have a detrimental effect on certain parts of conventional dosing guns.

Operator warnings:

Protective clothing, boots, gloves and waterproof apron must be worn when applying the product and when handling recently treated sheep. Wash hands and exposed skin before eating, drinking or smoking, and after work. Wash splashes from skin and eyes immediately. Use in a well-ventilated area. Remove heavily contaminated clothing immediately and wash before re-use.

11. EXPIRY DATE

Expires end:

12. SPECIAL STORAGE CONDITIONS

Store upright in the original container tightly closed in a safe place away from food, drink and animal feeding stuffs.

Do not store above 25°C. Store in a dry place

Protect from direct sunlight. Do not freeze. If contents freeze they must be thawed and thoroughly mixed before use.

13. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the product or empty container.

14. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

POM-VPS

To be supplied only on veterinary prescription.

For animal treatment only

15. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

16. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Zoetis Belgium S.A.
2nd Floor, Building 10
Cherrywood Business Park
Loughlinstown
Dublin 18
D18 T3Y1
Ireland

Distributed by:
Downland Marketing Limited
Main Mill
Warwick Mill Business Centre
Warwick Bridge
Carlisle
Cumbria
CA4 8RR

17. MARKETING AUTHORISATION NUMBER

Vm 60021/3035

18. MANUFACTURER’S BATCH NUMBER

Batch Number:

PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Shipper label}

PLEASE NOTE: Whilst the 4 x 5 Litre outer label has been approved by the VMD in the past, this component is in fact a shipper label so should not be under any regulatory control.

Approved 20 January 2025
Gavin Hall