

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Carton box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Lozenord 5 mg/ml solution for injection

2. STATEMENT OF ACTIVE SUBSTANCES

Meloxicam 5 mg/ml

3. PACKAGE SIZE

10 ml

4. TARGET SPECIES

Dogs and cats

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Dogs: intravenous or subcutaneous use.

Cats: subcutaneous use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

Once broached use within 28 days.

Once broached use by:.....

9. SPECIAL STORAGE PRECAUTIONS

Keep the vial in outer carton in order to protect from light.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Accord Healthcare B.V

14. MARKETING AUTHORISATION NUMBERS

Vm 42153/3000

15. BATCH NUMBER

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE
PACKAGING UNITS**

Vial, 10 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Lozenord 5 mg/ml solution for injection

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Each ml contains:

Active substance:
Meloxicam 5 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}
Once broached use within 28 days.
Once broached use by:.....

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Lozenord 5 mg/ml solution for injection for dogs and cats

2. Composition

Each ml contains:
Meloxicam 5 mg

Qualitative composition of excipients and other constituents	Quantitative composition
Ethanol anhydrous	150 mg

A clear, yellow solution.

3. Target species

Dogs and cats

4. Indications for use

Dogs:

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopaedic and soft tissue surgery.

Cats:

Alleviation of mild to moderate post-operative pain and inflammation following surgical procedures in cats, e.g. orthopaedic and soft tissue surgery

5. Contraindications

Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals less than 6 weeks of age nor in cats of less than 2 kg.

6. Special warnings

Special precautions for safe use in the target species:

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of renal toxicity. During anaesthesia, monitoring and fluid therapy should be considered as standard practice.

For post-operative pain and inflammation following surgical procedures in cats:

In case additional pain relief is required, multimodal pain therapy should be considered.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show this package leaflet or the label to the physician.

This product can cause eye irritation. In case of contact with the eyes, immediately rinse thoroughly with water.

Special precautions for the protection of the environment:

Not applicable.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation. Do not use in pregnant or lactating animals.

Interaction with other medicinal products and other forms of interaction:

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Meloxicam may also antagonise the antihypertensive effects of ACE inhibitors. The veterinary medicinal product must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic drugs should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such veterinary medicinal products should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacological properties of the products used previously.

Overdose:

In case of overdose symptomatic treatment should be initiated.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

7. Adverse events

Dogs and Cats

Very rare (<1 animal / 10,000 animals treated, including isolated reports)	Appetite loss ¹ , Lethargy ¹ Vomiting ¹ , diarrhoea ¹ , blood in faeces ^{1,2} , haemorrhagic diarrhoea ¹ , haematemesis ¹ , gastric ulcer ¹ , small intestine ulcer ¹ Elevated liver enzymes ¹ Renal failure ¹ Anaphylactoid reaction ³
--	--

- 1 These adverse events occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.
- 2 Occult
- 3 Should be treated symptomatically

If adverse reactions occur, treatment should be discontinued, and the advice of a veterinarian should be sought.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

8. Dosage for each species, routes and method of administration

Dosage for each species

Dogs: single administration of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg).

Cats: Single administration of 0.2 mg meloxicam/kg body weight (i.e. 0.04 ml/kg) when administration of meloxicam is to be continued as an oral follow-up therapy.
Single administration of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg) where no oral follow-up treatment is possible, e.g. feral cats.

Method and routes of administration

Dogs: Intravenous or subcutaneous use.

Cats: Subcutaneous use.

Dogs:

Musculo-skeletal disorders: single subcutaneous injection.

A suitable oral meloxicam formulation, e.g. suspension or tablet, administered in accordance with label recommendations, may be used for continuation of treatment 24 hours after administration of the injection.

Reduction of post-operative pain (over a period of 24 hours): single intravenous or subcutaneous injection before surgery, for example at the time of induction of anaesthesia.

Cats:

Reduction of post-operative pain and inflammation when administration of meloxicam is to be continued as an oral follow-up therapy: Single subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.04 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia. A suitable oral meloxicam formulation administered in accordance with label recommendations, may be used for continuation of treatment 24 hours after administration of the injection.

Reduction of post-operative pain and inflammation where no oral follow-up treatment is possible e.g. feral cats:

Single subcutaneous injection at a dosage of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia. In this case do not use oral follow up treatment.”

The rubber stopper should not be punctured more than 24 times.

9. Advice on correct administration

Particular care should be taken with regard to the accuracy of dosing. To ensure a correct dosage, body weight should be determined as accurately as possible. The use of suitably calibrated measuring equipment is recommended. Avoid introduction of contamination during use.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Keep the vial in outer carton in order to protect from light.

Shelf life after first opening of the immediate packaging: 28 days.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and the vial after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local

requirements and with any applicable national collection systems. These measures should help to protect the environment. Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

Colourless Type-I glass injection vial of 10 ml, closed with a grey chlorobutyl fluorotec rubber stopper and sealed with aluminium cap and flip off plastic tamper evident top.

Cardboard box containing a single glass vial.

15. Date on which the package leaflet was last revised

To be completed nationally.

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

16. Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Accord Healthcare B.V
Winthontlaan 200, Utrecht,
3526 KV, Netherlands
Telephone number: +44 (0) 208 901 3383

Manufacturer responsible for batch release:

Laboratori Fundació DAU
Calle Lletra C De La Zona Franca 12-14, Poligono Industrial De La Zona Franca
De Barcelona, Barcelona, 08040, Spain

17. Other information

Gavin Hall
Approved: 04 February 2025