

**<PARTICULARS TO APPEAR ON THE OUTER PACKAGE>**

**Cardboard box**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

FIPROSPOT 134 mg Spot-on solution for medium dogs  
Fipronil

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each pipette contains fipronil 134 mg

Excipients:

Butylhydroxyanisole (E 320) 0.268 mg

Butylhydroxytoluene (E 321) 0.134 mg

**3. PHARMACEUTICAL FORM**

Spot-on solution

**4. PACKAGE SIZE**

3 pipettes  
6 pipettes  
12 pipettes  
24 pipettes  
60 pipettes  
120 pipettes

**5. TARGET SPECIES**

For dogs  
(10 – 20 kg)

**6. INDICATION(S)**

Against fleas and ticks.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

Not applicable

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Store in the original package.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

IDT Biologika GmbH  
Am Pharmapark  
06861 Dessau-Rosslau  
Germany

Distributed by:  
to be determined nationally

**16. MARKETING AUTHORISATION NUMBER(S)**

## 17. MANUFACTURER'S BATCH NUMBER

Batch number



**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blister**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

FIPROSPOT 134 mg Spot-on solutions for medium dogs  
Fipronil

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

LOGO (of IDT)  
IDT Biologika GmbH

or

Distributed by:  
LOGO (of distributor)  
Name of distributor (to be determined nationally)

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

<b>MINIMUM PARTICULARS TO APPEAR ON PIPPETES</b>
--------------------------------------------------

<b>Pipette</b>
----------------

<b>1. NAME OF THE VETERINARY MEDICINAL PRODUCT</b>
----------------------------------------------------

FIPROSPOT  
134 mg Fipronil

<b>2. EXPIRY DATE</b>
-----------------------

EXP

<b>3. BATCH NUMBER</b>
------------------------

Lot

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET FOR

### FIPROSPOT 134 mg Spot-on solution for medium dogs

#### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder

IDT Biologika GmbH  
Am Pharmapark  
D-06861 Dessau-Rosslau  
Germany

Manufacturer responsible for batch release

Klocke Verpackungs-Service GmbH  
Max-Becker-Str. 6  
D-76356 Weingarten  
Germany

Distributed by  
to be determined nationally

#### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

FIPROSPOT 134 mg Spot-on solution for medium dogs  
Fipronil

#### 3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each 1.34 ml pipette contains

**Active substance:**

Fipronil	134.00 mg
----------	-----------

**Excipients**

Butylhydroxyanisole (E 320)	0.268 mg
Butylhydroxytoluene (E 321)	0.134 mg

Clear, colourless to yellowish solution.

#### 4. INDICATIONS

For the treatment of dogs against flea infestations (*Ctenocephalides* spp.)

Insecticidal efficacy against new infestation with fleas persists for up to 6 weeks.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD), where this has been previously diagnosed by a veterinary surgeon.

Although the product does not consistently show an immediate acaricidal efficacy (several ticks may be present after 48 hours), it has a persistent acaricidal efficacy for up to 4 weeks against *Dermacentor variabilis* and up to 3 weeks against *Rhipicephalus sanguineus*.

## **5. CONTRAINDICATIONS**

Do not use on puppies less than 2 months old and/or weighing less than 2 kg in the absence of available data.

Do not use on sick (systemic disease, fever, etc.) or convalescent animals.

Do not use on rabbits, as adverse drug reactions and even death could occur.

This veterinary medicinal product has been developed specifically for dogs. Do not use on cats as this could lead to overdosing.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

## **6. ADVERSE REACTIONS**

If licking occurs, a brief period of hypersalivation may be observed due mainly to the nature of the carrier substance.

Among the very rare suspected adverse reactions, transient cutaneous reactions at the application site (scaling, local alopecia, pruritus, erythema) and general pruritus or alopecia have been reported after use.

Very rare, hypersalivation, reversible neurological signs (hyperaesthesia, depression, nervous signs), vomiting or respiratory signs have been observed after use.

The frequency of adverse reactions is defined using the following convention:

Very common (more than 1 in 10 animals displaying adverse reaction(s) during the course of one treatment)

Common (more than 1 but less than 10 animals in 100 animals)

Uncommon (more than 1 but less than 10 animals in 1,000 animals)

Rare (more than 1 but less than 10 animals in 10,000 animals)

Very rare (less than 1 animal in 10,000 animals, including isolated reports)

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dog (10-20 kg)

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Method of application and posology:

Only by topical application to the skin. External use only.

Animals should be weighed accurately prior to treatment.

1 pipette of 1.34 ml is sufficient for the treatment of a dog with a body weight of 10 kg up to 20 kg corresponding to a recommended minimum dose of 6.7 mg fipronil/kg body weight.

The minimum interval between two treatments should be not less than 4 weeks.

## **9. ADVICE ON CORRECT ADMINISTRATION**



Disconnect one of the blisters from the blister card. This helps to avoid accidental opening of the adjacent blister package in order to protect the still unopened pipettes from exposure to humidity. Open the blister with scissors. To avoid damaging of the pipette cut along the line marked with the scissors icon. Carefully peel back the foil from the cut off end and withdraw the pipette.

Hold the pipette upright. Tap lightly to ensure the entire liquid contents are within the main body of the pipette. Bend the upper border strip backwards. Then the pipette can be set aside, if necessary. To open the pipette snap off the top of the pipette along the scored line.

Part the coat between the shoulder blades until the skin is visible. Place the tip of the pipette on the skin and squeeze the pipette several times to empty its content completely and directly onto the skin in one spot.

Application of the solution near the base of the head minimises the possibility that the animal will lick the solution off. Care should be taken after the application that animals do not mutually lick off the solution.

Care should be taken to avoid excessive wetting of the hair with the product since this will cause a sticky appearance of hairs at the treatment spot.

However, should this occur, it will disappear within 24 hours post application.

## **10. WITHDRAWAL PERIOD**

Not applicable

## **11. SPECIAL STORAGE PRECAUTION**

This veterinary medicinal product does not require any special storage conditions. Store in the original package.

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP.

## **12. SPECIAL WARNINGS**

The veterinary medicinal product does not prevent an infestation of the animal by ticks.

Ticks will usually die within 48 hours of infestation however attached ticks (both live and killed) may be seen at this time. Some of these will have had a blood meal.

Death normally occurs before the ticks are fully engorged so that the risk of transmission of infectious diseases by ticks is minimised, but cannot be completely ruled out. As soon as the ticks are dead they generally fall off the animal; remaining ticks can be removed with a gentle pull.

For the optimal control of flea problems in households with several animals all dogs and cats should be treated with an authorised insecticide.

Fleas from pets often infest animal's baskets, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.

Shampooing with a medicated shampoo, followed by thorough drying, 1 to 2 hours before treatment application and bathing once weekly over a period of 6 weeks, has been shown not to affect the efficacy of this veterinary medicinal product against fleas. Bathing and intensive wetting of the coat should be avoided for the first 2 days following administration of the veterinary medicinal product.

When used as part of a strategy for the treatment of Flea Allergy Dermatitis, monthly applications to the allergic patients and to other animals in the household are recommended.

#### Special precautions for use in animals:

This veterinary medicinal product should not be used on dogs weighing less than 10 kg.

It is important to make sure that the product is applied to an area where the animal cannot lick it off. Do not allow recently treated animals to lick each other.

Avoid contact with the animal's eyes. Should the veterinary medicinal product come into contact with the eyes, rinse thoroughly at once with water.

Do not apply the veterinary medicinal product to wounds or skin lesions.

There may be an attachment of some ticks. For this reason transmission of infectious diseases cannot be completely excluded if conditions are unfavourable.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This product can cause mucous membrane and eye irritation. Therefore, contact between the product and the mouth or eyes should be avoided.

In the case of accidental eye contact immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the physician.

Avoid contents coming into contact with the fingers. If this occurs, wash off immediately with soap and water.

Wash hands after use.

Do not smoke, drink or eat during application.

People with a known hypersensitivity to fipronil or other excipients should avoid contact with the veterinary medicinal product.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Keep pipettes in the original packaging and dispose of used pipettes immediately.

#### Other precautions

Dogs should not be allowed to swim in watercourses for 2 days after application.

The product may have adverse effects on painted, varnished or other household surfaces or furnishings.

#### Use during pregnancy, lactation or lay

The safety of the veterinary medicinal product has not been established in dogs during pregnancy and lactation. Laboratory studies using fipronil have not shown any evidence of teratogenic or foetotoxic effect. Use in pregnancy and lactation only in accordance with the benefit/risk assessment of the responsible veterinarian.”

Overdose (symptoms, emergency procedures, antidotes)

The toxicity of the veterinary medicinal product administered to the skin is very low. The risk of experiencing adverse effects (see section 6.) may however increase when overdosing, so animals should always be treated with the correct pipette size according to body weight.

Specific studies investigating the safety of the product following repeated administration or at overdosage have not been conducted due to the known safety profile of the active substance and excipients.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY**

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

Fipronil may be harmful to aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED:**

**15. OTHER INFORMATION**

The veterinary medicinal product distributes itself within 48 hours over the entire skin of the animal.

Package sizes:

A blister card consists of 3 blisters each containing a single pipette.

Packs containing 3, 6, 12, 24, 60 and 120 pipettes.

Not all pack sizes may be marketed