

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**CARTON**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flunixin 50 mg/ml Solution for Injection for Cattle, Horses and Pigs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Flunixin, (as flunixin meglumine) 50 mg

**3. PHARMACEUTICAL FORM**

Solution for Injection

**4. PACKAGE SIZE**

50 ml

100 ml

250 ml

**5. TARGET SPECIES**

Cattle, Horses and Pigs

**6. INDICATION(S)**

Read the package leaflet before use.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Cattle: Meat & offal: 7 days

Milk: 36 hours

Horses: Meat & offal: 7 days

Pigs: Meat & offal: 22 days

Do not use in mares producing milk for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use

**10. EXPIRY DATE**

EXP:

**11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Keep the vial in the outer carton to protect from light.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Norbrook Laboratories Limited  
Station Works  
Camlough Road  
Newry  
Co. Down  
BT35 6JP  
Northern Ireland

**Distributed By:**

Norbrook Laboratories Limited  
Carnbane Industrial Estate  
Newry  
BT35 6QQ, Co Down  
Northern Ireland

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 02000/4170

**17. MANUFACTURER'S BATCH NUMBER**

B.N.:

LOGO

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**LABEL**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flunixin 50 mg/ml Solution for Injection for Cattle, Horses and Pigs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Flunixin (as flunixin meglumine) 50 mg

**3. PHARMACEUTICAL FORM**

Solution for Injection

**4. PACKAGE SIZE**

50 ml  
100 ml  
250 ml

**5. TARGET SPECIES**

Cattle, Horses and Pigs

**6. INDICATION(S)**

Read the package leaflet before use.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Cattle: Meat & offal: 7 days

Milk: 36 hours

Horses: Meat & offal: 7 days

Pigs: Meat & offal: 22 days

Do not use in mares producing milk for human consumption

**9. SPECIAL WARNING(S), IF NECESSARY**

**10. EXPIRY DATE**

EXP:  
Once broached use by.....

**11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.  
Keep the vial in the outer carton to protect from light.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Norbrook Laboratories Limited  
Station Works  
Camlough Road  
Newry  
Co. Down  
BT35 6JP  
Northern Ireland

**Distributed By:**  
Norbrook Laboratories Limited  
Carnbane Industrial Estate  
Newry  
BT35 6QQ, Co Down  
Northern Ireland

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 02000/4170

**17. MANUFACTURER’S BATCH NUMBER**

B.N.:

**B. PACKAGE LEAFLET**



**PACKAGE LEAFLET FOR:**

Flunixin 50 mg/ml Solution for Injection for Cattle, Horses and

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Norbrook Laboratories Limited  
Station Works  
Camlough Road  
Newry  
Co. Down  
BT35 6JP  
Northern Ireland

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flunixin 50 mg/ml Solution for Injection for Cattle, Horses and Pigs

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each ml contains:

Flunixin (as flunixin meglumine)	50 mg
Phenol (as preservative)	5 mg
Sodium Formaldehyde Sulphoxylate Dihydrate	2.5 mg

A clear colourless solution

**4. INDICATION(S)**

In the horse, Flunixin Injection is indicated for the alleviation of inflammation and pain associated with musculo-skeletal disorders and for the alleviation of visceral pain associated with colic. In cattle, Flunixin Injection is indicated for the control of acute inflammation associated with respiratory disease. It may also be used as adjunctive therapy in the treatment of acute mastitis. In Pigs, Flunixin Injection is indicated as an adjunctive therapy in the treatment of swine respiratory diseases.

**5. CONTRAINDICATIONS**

Do not exceed the recommended dose or the duration of treatment.

Do not administer to pregnant mares.

Do not administer to pregnant sows, gilts at mating and in breeding boars.

Use is contraindicated in animals suffering from cardiac, hepatic or renal disease, where there is the possibility of gastrointestinal ulceration or bleeding, where there is evidence of a blood dyscrasia or hypersensitivity to the product.

Do not use in dehydrated animals suffering from ileus-associated colics.

Do not use the product within 48 hours before expected parturition in cows.

## **6. ADVERSE REACTIONS**

Flunixin meglumine is a non steroidal anti-inflammatory drug (NSAID). Untoward effects include gastrointestinal irritation, ulceration and, in dehydrated or hypovolaemic animals, potential for renal damage.

Rare cases of anaphylactic reaction have been reported. In horses (rare) and cattle (very rare) anaphylaxis type reactions can include neurological signs such as convulsion, loss of consciousness and ataxia. Such reactions may be exacerbated by intra-arterial injection.

In pigs (very rare), transient irritation may occur at the injection site, this resolves spontaneously within 14 days.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

If you notice any serious effects or other effects not mentioned in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Cattle, Horses and Pigs

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Flunixin Injection is indicated for intravenous administration to cattle and horses and intramuscular injection to pigs.

Do not exceed the recommended dose or duration of treatment.

**HORSES:** For use in equine colic, the recommended dose rate is 1.1 mg flunixin/kg bodyweight equivalent to 1 ml per 45 kg bodyweight by intravenous injection. Treatment may be repeated once or twice if colic recurs.

For use in musculo-skeletal disorders, the recommended dose rate is 1.1 mg flunixin/kg bodyweight equivalent to 1 ml per 45 kg bodyweight, injected intravenously once daily for up to 5 days according to clinical response.

For the treatment of endotoxaemia or septic shock associated with gastric torsion and other conditions in which the circulation of blood to the gastro-intestinal tract is compromised: 0.25 mg/kg (1 ml per 200 kg) every 6-8 hours.

**CATTLE:** The recommended dose rate is 2.2 mg flunixin/kg bodyweight equivalent to 2 ml per 45 kg bodyweight injected intravenously and repeated as necessary at 24 hour intervals for up to 5 consecutive days.

**PIGS** For use in pigs, the recommended dose rate is 2 ml per 45 kg bodyweight (equivalent to 2.2 mg flunixin/kg) once by intramuscular injection, in the neck, in conjunction with appropriate antimicrobial therapy. The injection volume should be limited to a maximum of 5 ml per injection site.

## **9. ADVICE ON CORRECT ADMINISTRATION**

An appropriately graduated syringe must be used to allow accurate administration of the required dose volume. This is particularly important when injecting small volumes.

This veterinary medicinal product must not be mixed with other veterinary medicinal products.

Avoid introduction of contamination.

The stopper should not be punctured more than 50 times. A draw off needle should be used to avoid excessive puncturing of the stopper.

## **10. WITHDRAWAL PERIOD(S)**

**Cattle:** Meat & offal: 7 days  
Milk: 36 hours

**Horses:** Meat & offal: 7 days

**Pigs:** Meat & offal: 22 days

Do not use in mares producing milk for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C. Keep the vial in the outer carton to protect from light.

**Keep out of the sight and reach of children**

Do not use after the expiry date stated on the carton and the label

Following withdrawal of the first dose, use the product within 28 days. Discard unused product.

When the container is breached for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the container should be discarded should be worked out. This discard date should be written in the space provided on the label.

## **12. SPECIAL WARNING(S)**

### **Special warnings for each target species:**

The cause of the underlying inflammatory condition or colic should be determined and treated with appropriate concomitant therapy.

Non-steroidal anti-inflammatory drugs are not permitted under the Rules of Racing and under rules covering other competitive events. Horses intended for racing and competition should be prevented from racing or competing when in need of treatment and horses which have been recently treated should be dealt with according to local requirements. Appropriate precautions must be taken to ensure compliance with competition regulations.

Cattle should be treated with flunixin in conjunction with disease-specific therapy and an improvement in housing conditions.

The use of flunixin in conjunction with disease-specific antibiotic therapy may mask antibiotic resistance of the bacteria, due to alleviation of inflammation symptoms.

### **Special precautions for use in animals:**

Avoid intra-arterial injection.

Use in any animal less than 6 weeks of age or in aged animals may involve an additional risk. If such use cannot be avoided animals may require a reduced dosage and careful clinical management.

Do not use in piglets weighing less than 6 kg.

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, except in the case of endotoxaemia or septic shock, as there is a potential risk of increased renal toxicity.

It is preferable that NSAIDs, which inhibit prostaglandin synthesis, are not administered to animals undergoing general anaesthesia until fully recovered.

Due to the excipient propylene glycol, life-threatening shock reactions may occur in rare cases. The solution for injection should therefore be injected slowly and be of approximate body temperature.

NSAIDs are known to have the potential to delay parturition through a tocolytic effect by inhibiting prostaglandins that are important in signalling the initiation of parturition. The use of the product in the immediate post-partum period may interfere with uterine involution and expulsion of fetal membranes resulting in retained placentae.

Flunixin is toxic to avian scavengers. Do not administer to animals susceptible to enter wild fauna food chain. In case of death or sacrifice of treated animals, ensure that they are not made available to wild fauna.

**Special precautions to be taken by the person administering the veterinary medicinal product to animals:**

The veterinary medicinal product can cause skin and eye irritation. Avoid contact with skin and eyes. In case of accidental skin exposure, wash the affected area immediately with plenty of water. In case of accidental eye contact, rinse immediately with plenty of water. If skin and/or eye irritation persists, seek medical advice immediately and show the package leaflet or the label to the physician.

The veterinary medicinal product can cause hypersensitivity (allergy) reactions. People with known hypersensitivity to non-steroidal anti-inflammatory drugs should avoid contact with the veterinary medicinal product. Adverse reactions can be serious. Gloves should be worn during application.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

**Use in Pregnancy, lactation or lay:**

May be used in pregnant and lactating cattle.

Do not administer to pregnant mares. Do not administer to pregnant sows, gilts at mating and in breeding boars. Safety studies in pregnant mares or sows have not been conducted.

The product should not be used in lactating sows.

The product should only be administered within the first 36 hours post-partum following a benefit/risk assessment performed by the responsible veterinarian and treated animals should be monitored for retained placentae.

**Interaction with other medicinal products and other forms of interaction:**

Monitor drug compatibility closely where adjunctive therapy is required.

Do not administer other non-steroidal anti-inflammatory drugs (NSAIDs) concurrently or within 24 hours of each other. Some NSAIDs may be highly bound to plasma proteins and compete with other highly bound drugs which can lead to toxic effects.

Concurrent administration of potentially nephrotoxic drugs should be avoided.

**Overdose:**

Overdose studies in the target species have shown the product to be well tolerated. Overdosage is associated with gastrointestinal toxicity.

**Incompatibilities:**

In the absence of incompatibility studies, this medicinal product must not be mixed with other medicinal products.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

June 2023

Detailed information on this product is available on the website of the European Medicines Agency (<http://www.ema.europa.eu/>).

**15. OTHER INFORMATION**

Packaging quantities:

Multi-dose vials of 50 ml, 100 ml and 250 ml.

The product is also presented in packs of 5, 10 and 12 vials for the 50 ml and 100 ml and a pack of 5 vials for the 250 ml.


Flunixin Injection has also been shown to have some benefit in the treatment of experimental acute pulmonary emphysema (fog fever).

Flunixin meglumine is a relatively potent non-narcotic, non-steroidal analgesic with anti-inflammatory, anti-endotoxic and anti-pyretic properties.

Flunixin is toxic to avian scavengers although foreseen low exposure leads to low risk.

**Distributed By:**

Norbrook Laboratories Limited  
Carnbane Industrial Estate  
Newry  
BT35 6QQ, Co Down  
Northern Ireland



Approved: 02 June 2023