

PARTICULARS TO APPEAR ON THE OUTER PACKAGE - CARTON

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

WORMclear Praziquantel 20 mg Spot-on Solution for Cats

2. STATEMENT OF ACTIVE SUBSTANCES

Each tube contains 0.5 ml of solution equivalent to 20 mg praziquantel/tube.

3. PACKAGE SIZE

2 tubes x 0.5 ml

4 tubes x 0.5 ml

20 mg solution

4. TARGET SPECIES

Cats

5. INDICATIONS

For the treatment of tapeworms of cats. The product is effective against mature and immature forms of *Dipylidium caninum*, *Taenia* species and *Echinococcus multilocularis*.

6. ROUTES OF ADMINISTRATION

The minimum dose rate is 8 mg/kg bodyweight equivalent to 1 tube per 2.5 kg.

Bodyweight (kg)	Tubes
1 – 2.5 kg	1
MORE THAN 2.5 – 5 kg	2
MORE THAN 5 kg	3

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp.{mm/yyyy}

Do not use after the expiry date.

9. SPECIAL STORAGE PRECAUTIONS

Store away from food, drink and animal feeding stuffs.
Keep the container and the leaflet in the outer carton.

The solvent in the product may stain certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

For full user safety warnings, read the package leaflet.

Pregnant women and women suspected of being pregnant should not administer the product.

Do not stroke or groom animals until the area of application is dry (typically around 1 hour after application).

Wash hands thoroughly after use.

After use, replace cap on tube and dispose of empty packaging and any remaining product in the household refuse.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Vetoquinol UK Limited

14. MARKETING AUTHORISATION NUMBERS

Vm 08007/5029

15. BATCH NUMBER

Lot {number}

Further information:

Distributed by:

Call our Helpline on 0333 0066236, email us at info@uk.beaphar.com or write to us at Beaphar UK Ltd, Rook Tree Farm, Withersfield Road, Great Wratting, Suffolk, CB9 7HD.

Do NOT administer by mouth.

Do not use this product on cats weighing less than 1.0 kg.

Do not allow recently treated animals to groom each other.
Use during pregnancy and lactation – read the package leaflet.
The safety of the veterinary medicinal product has not been established in cats during pregnancy, lactation or in animals intended for breeding.

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS – TUBE LABEL**

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

WORMclear Praziquantel 20 mg Spot-on

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

20 mg

0.5 ml

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

WORMclear Praziquantel 20 mg Spot-on Solution for Cats

2. Composition

The active ingredient is praziquantel which kills your pet's tapeworms. Each tube of the veterinary medicinal product contains 0.5 ml of liquid containing 20 mg of praziquantel. The solution also contains inactive ingredients which are needed to allow the drug to be absorbed across the skin and into the cat's body where it can be effective. Each tube contains 497.8 mg of N-methylpyrrolidone.

Clear, colourless to slightly reddish liquid.

3. Target species

Cats.

4. Indications for use

The veterinary medicinal product is used for the treatment of tapeworms. It is effective against adult and immature forms of tapeworms known as *Dipylidium caninum* and tapeworms of the *Taenia* species. The veterinary medicinal product has been specifically designed for administration to cats.

5. Contraindications

Do not use this product in animals weighing less than 1 kg.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

6. Special warnings

For external use only.

Do not allow recently treated animals to groom each other.

Special precautions for safe use in the target species:

Care should be taken to avoid the contents of the tube coming into contact with the eyes or mouth of the recipient animal.

If signs of disease persist or appear, consult a veterinary surgeon.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

User safety – IMPORTANT: ALWAYS READ THIS SECTION CAREFULLY

Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. The veterinary medicinal product should not be administered by pregnant women and women suspected of being pregnant. Personal protective equipment consisting of gloves should be worn when handling the veterinary medicinal product by women of childbearing age.

This veterinary medicinal product can be irritant to the skin and eyes.

Care should be taken to prevent contact of the solution with the skin or eyes.

In case of accidental spillage onto skin, wash off any skin contamination with soap and water immediately.

In case of accidental contact with the eyes, flush the affected eyes thoroughly with clean fresh water.

If skin or eye irritation persists, seek medical advice immediately and show the package leaflet or the label to the doctor.

Do not stroke or groom animals until area of application is dry (typically around 1 hour after application).

Wash hands thoroughly after use.

Do not eat, drink or smoke during application.

Other precautions:

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (WOAH), specific guidelines on the treatment and follow-up and on the safeguard of persons need to be obtained from the relevant competent authority.

The solvent in the product may stain certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

For Animal Treatment Only.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established in cats during pregnancy, lactation or in animals intended for breeding. Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. Use only accordingly to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

None known.

Overdose:

Overdosing can lead to slight skin reactions which disappear without treatment within a few days.

Major incompatibilities:

None known.

7. Adverse events

Cats

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site reaction ¹ (e.g. dandruff ¹ , exudation ^{1,2} , hair loss ¹ , reddening ¹ , itching ¹ , scab ¹) Hypersalivation ³
---	---

¹ Transient

² Mild

³ Due to the bitter taste of the product if the cat licks the application site immediately after treatment. This is not a sign of intoxication and disappears after a short time without treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, routes and method of administration

The veterinary medicinal product should be administered at monthly intervals or when signs of tapeworm infestation appear. The number of tubes to be used on your cat depends on how much it weighs.

For cats weighing from 1 to 2.5 kg: use 1 tube.

For cats weighing over 2.5 kg up to 5 kg: use 2 tubes.

For cats weighing more than 5 kg: use 3 tubes.

9. Advice on correct administration

Warning: Do NOT administer the product by mouth.

Weigh your cat to see how many tubes of the veterinary medicinal product will be required.

Underdosing could result in ineffective use and may favour resistance development.


Take the required number of tubes from the blister pack.

Hold one tube in an upright position with the cap at the top.

Twist and pull off the cap.

Reverse the cap and replace it onto the tube.

Twist reversed cap, this will break the seal across the top of the tube.



On the back of the animal's neck, immediately behind the head, part the coat until the skin is visible.

Place the tip of the opened tube on the skin and squeeze tube firmly several times to empty the entire contents directly onto the skin. When it is necessary to use more than one tube, care must be taken to prevent the solution running off the skin of the animal. In this situation it is advised that the applications should be made slowly to allow time for absorption of the solution. Only begin application from the second or third tubes when the contents of the previous tube have been completely absorbed.

It is particularly important to apply the dose to an area where it cannot be licked off. Application to the back of the neck at the base of the skull will minimise the opportunity for the cat to lick the product.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Store this product away from food, drink and animal feeding stuffs.

Do not administer this product to your pet after the expiry date on the carton. The expiry date refers to the last day of that month.

12. Special precautions for disposal

After you have applied the product replace the cap on the tube. Dispose of empty packaging and any remaining product in the household refuse.

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product not subject to prescription.

14. MARKETING AUTHORISATION NUMBER AND PACK SIZES

Vm 08007/5029

Pack size: Carton with 2 or 4 tubes.

To safeguard the health of yourself and your pet, please keep this leaflet and read it before you apply the product to your pet. It provides a summary of information about this veterinary medicine. For any information about this veterinary medicinal product, please contact the local distributor.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder:

Vetoquinol UK Limited
Steadings Barn
Pury Hill Business Park, Nr. Alderton
Towcester
Northamptonshire
NN12 7LS
United Kingdom

Manufacturer responsible for batch release:

VETOQUINOL BIOWET Sp. z o.o.
Żwirowa 140,
66-400, Gorzów, Wlkp.,
Poland

Local representatives and contact details to report suspected adverse reactions:

Beaphar UK Limited
Rook Tree Farm
Withersfield Road
Great Wratting
Suffolk
CB9 7HD
Tel: 0333 0066236
Email: info@uk.beaphar.com

17. Other information

AVM-GSL

Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas and mice is undertaken. Flea infestations can be controlled by the regular use of effective flea control remedies.

Mice control: if cats roam and hunt, they will come into contact with, and possibly consume, mice. As the mouse is one of the hosts of a tapeworm known as *Taenia taeniaformis*, re-infestation of the cat with this tapeworm is impossible to prevent.

Gavin Hall
Approved 16 May 2026