

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**DUO 100 mg/120 mg spot-on solution for very large cats FIPRALONE DU  
FIPROKIL DUO 100 mg/120 mg spot-on solution for very large cats (FR)**

**Box containing 1 individual pipette placed in overblister  
Box containing 2 individual pipettes placed in 1 overblister  
Box containing 3 individual pipettes placed in 2 overblisters  
Box containing 4 individual pipettes placed in 2 overblisters  
Box containing 6 individual pipettes placed in 3 overblisters**



**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

DUO 100 mg/120 mg spot-on solution for very large cats FIPRALONE  
FIPROKIL DUO 100 mg/120 mg spot-on solution for very large cats (FR)  
Fipronil/Pyriproxyfen

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each 1 ml pipette contains: Fipronil 100 mg – Pyriproxyfen 120 mg

**3. PHARMACEUTICAL FORM**

Spot-on solution

**4. PACKAGE SIZE**



1  
1 ml



2  
1 ml



3  
1 ml



4  
1 ml



6  
1 ml

## 5. TARGET SPECIES

Cats >6-12 kg

## 6. INDICATION(S)

**\*\*\*Only for those countries where the product is available without prescription:\*\*\***

To be used against infestations with fleas alone or in association with ticks.  
Fleas, flea eggs & Ticks

## 7. METHOD AND ROUTE(S) OF ADMINISTRATION

Spot-on use.

**\*\*\*Only for multilingual boxes on which the available space will be sufficient:\*\*\***

 [optional]	Cat	Fipronil	Pyriproxyfen
1 ml	>6-12 kg	100 mg	120 mg

Read the package leaflet before use.



[optional]

## 8. WITHDRAWAL PERIOD

Withdrawal period: Not applicable.

### 9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.



[optional]

### 10. EXPIRY DATE

EXP {month/year}

### 11. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.

Store in a dry place.

Keep the blister pack in the outer carton in order to protect from light.



30°C

[optional]

### 12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Read the package leaflet before use.



[optional]

### 13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.

### 14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.



[optional]

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ALFAMED  
13ème Rue – L.I.D.  
06517 CARROS CEDEX  
FRANCE

**16. MARKETING AUTHORISATION NUMBER(S)**

**17. MANUFACTURER'S BATCH NUMBER**

Lot {number}


**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**DUO 100 mg/120 mg spot-on solution for very large cats FIPRALONE DUO 100 mg/120 mg spot-on solution for very large cats (UK)  
FIPROKIL DUO 100 mg/120 mg spot-on solution for very large cats (FR)**

**Overblister packed in 1-pipette blister or 2-pipette blisters divisible per pipette**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

DUO 100 mg/120 mg spot-on FIPRALONE DUO 100 mg/120 mg spot-on (UK)  
FIPROKIL DUO 100 mg/120 mg spot-on (FR)  
Fipronil/Pyriproxyfen



**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

Fipronil	100 mg
Pyriproxyfen	120 mg

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

Pipette volume 1 ml

**4. ROUTE(S) OF ADMINISTRATION**

Spot-on use.

**5. WITHDRAWAL PERIOD**

Not applicable.

**6. BATCH NUMBER**

Lot {number}

**7. EXPIRY DATE**

EXP {month/year}

**8. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.


**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**DUO 100 mg/120 mg spot-on solution for very large cats FIPRALONE DUO 100 mg/120 mg spot-on solution for very large cats (UK)**  
**FIPROKIL DUO 100 mg/120 mg spot-on solution for very large cats (FR)**

**Individual Pipette**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

DUO 100 mg/120 mg spot-on FIPRALONE DUO 100 mg/120 mg spot-on (UK)  
FIPROKIL DUO 100 mg/120 mg spot-on (FR)  
Fipronil/Pyriproxyfen



**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

Fipronil 100 mg  
Pyriproxyfen 120 mg

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

Pipette volume 1 ml

**4. ROUTE(S) OF ADMINISTRATION**

Spot-on use.

**5. WITHDRAWAL PERIOD**

Not applicable.

**6. BATCH NUMBER**

Lot {number}

**7. EXPIRY DATE**

EXP {month/year}

**8. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**PACKAGE LEAFLET FOR:**

**Fipralone duo 50 mg/60 mg spot-on solution for cats  
Fipralone duo 100 mg/120 mg spot-on solution for very large cats**

**Fiprokil duo 50 mg/60 mg spot-on solution for cats  
Fiprokil duo 100 mg/120 mg spot-on solution for very large cats  
(FR)**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder and manufacturer responsible for batch release:  
ALFAMED  
13ème Rue – L.I.D.  
06517 CARROS CEDEX  
FRANCE

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fipralone duo 50 mg/60 mg spot-on solution for cats  
Fipralone duo 100 mg/120 mg spot-on solution for very large cats

Fiprokil duo 50 mg/60 mg spot-on solution for cats  
Fiprokil duo 100 mg/120 mg spot-on solution for very large cats  
(FR)

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

Each pipette contains:	Active substances		Excipients	
	Fipronil	Pyriproxyfen	BHA*	BHT**
Pipette volume (single dose unit)				
0.5 ml	50 mg	60 mg	0.1 mg	0.05 mg
1 ml	100 mg	120 mg	0.2 mg	0.10 mg

\*: Butylhydroxyanisole E320, \*\*: Butylhydroxytoluene E321

Clear, colourless to yellowish solution.

**4. INDICATION(S)**

In cats, to be used against infestations with fleas alone or in association with ticks.

Against fleas:

Treatment and prevention of infestations by fleas (*Ctenocephalides felis*). One treatment prevents further infestations for 5 weeks.

Prevention of the multiplication of fleas by preventing flea eggs developing into adult fleas for 12 weeks after application.

Against ticks:

Treatment of infestations by ticks (*Ixodes ricinus* and *Rhipicephalus turanicus*).

One treatment provides persistent acaricidal efficacy for one week.

If ticks are present at the time of application, not all ticks may be killed within 48 hours.

## 5. CONTRAINDICATIONS

Do not use in rabbits, as adverse reactions and even death could occur.

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

## 6. ADVERSE REACTIONS

Transient cosmetic effects such as wet appearance or slight scaling may occur at the application site.

According to the accumulated experience on these active ingredients within spot on pharmaceutical forms, transient cutaneous reactions at the application site (squamosis, local alopecia, pruritus, erythema, skin discolouration) and general pruritus or alopecia may be observed after use. In very rare instances, hypersalivation, reversible neurologic symptoms (hyperesthesia, depression, nervous symptoms), respiratory signs or vomiting might occur.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

For cats

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Spot-on use.

Dosage:

The minimum recommended dose is 8.3 mg fipronil/kg b.w. and 10 mg pyriproxyfen/kg b.w..

For a cat weighing at least 1 kg and up to 6 kg apply one pipette of 0.5 ml per cat.

For a cat weighing over 6 kg a recommended dose of 1 ml should be applied, which can be achieved by applying one pipette of 1 ml (or alternatively two pipettes of 0.5 ml).

Cat weight	Pipette volume	Fipronil (mg)	Pyriproxyfen (mg)
------------	----------------	---------------	-------------------

1-6 kg	0.5 ml		50	60
>6-12 kg	1 ml		100	120


Method of administration:

Remove the pipette from the overblister. Hold the pipette upright. Tap the narrow part of the pipette to ensure that the contents are within the main body of the pipette. Break the snap-off top of the spot-on pipette along the scored line.

Part the coat on the back of neck area until the skin is visible. Place the tip of the pipette on the skin and squeeze firmly several times to empty the contents evenly at one or two spots. Ensure the solution is applied to healthy skin only and avoid superficial application to the cat's hair or run off, especially in large cats (over 6 kg).



Drop stop system (the product is released only by pressing the body of the pipette).



## 9. ADVICE ON CORRECT ADMINISTRATION

For optimal control of flea and tick infestations and flea multiplication, the treatment schedule can be based on the local epidemiological situation. However, in the absence of additional safety studies, do not repeat the treatment at intervals of less than 4 weeks (see section 12. Overdose).

## 10. WITHDRAWAL PERIOD

Not applicable.

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and label after {EXP}. The expiry date refers to the last day of that month. Do not store above 30°C.

Store in a dry place.

Keep the blister pack in the outer carton in order to protect from light.

## 12. SPECIAL WARNING(S)

### Special warnings for each target species:

Wetting of the haircoat with water 2 hours before product application and twice during the claimed period of efficacy against fleas (at two week intervals against adult fleas or four week intervals against developing flea stages) was investigated during two laboratory studies. Wetting of the haircoat with water as described did not adversely affect product effectiveness.

The impact of shampooing on product effectiveness has not been investigated. If a cat requires shampooing it is recommended that this is done before applying the veterinary medicinal product.

At the beginning of the control measures, in the case of an infestation, the animal's basket, bedding and regular resting areas such as carpets and soft furnishings should be treated, with a suitable insecticide and vacuumed regularly.

To reduce environmental flea challenge, all animals living in the same household should also be treated with a suitable flea control product.

The product does not prevent ticks from attaching to animals. If the animal has been treated prior to exposure to the ticks, the ticks will be killed within 48 hours after attachment. This will usually be prior to engorgement, minimising but not excluding the risk of transmission of diseases.

Once dead, ticks will often drop off the animal. Any remaining ticks should be carefully removed, ensuring that their mouth parts are not left within the skin.

### Special precautions for use in animals:

For external use only.

Do not administer orally.

Animals should be weighed accurately prior to treatment.

The safety of the product has not been established in cats younger than 10 weeks of age or in cats weighing less than 1.0 kg.

Avoid contact with the animal's eyes. In the case of accidental eye contact, flush the eyes with water.

Care should be taken to apply the product correctly as described under section 8. Do not apply the product on wounds or damaged skin. It is important to make sure that the veterinary medicinal product is applied directly onto an area of dry skin where the animal cannot lick it off and to make sure that animals do not lick each other following treatment.

The use of the product has not been studied in sick or debilitated cats. Use in sick or debilitated animals only in accordance with a benefit/risk assessment performed by the responsible veterinarian.

In the absence of additional safety studies, do not repeat the treatment at intervals of less than 4 weeks.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The product may cause neurotoxicity. The product may be harmful if swallowed. Avoid ingestion including hand-to-mouth contact. Avoid contact with skin and mouth. Do not smoke, drink or eat during application. Wash hands after use. In the event of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician. In the case of accidental skin contact, flush with water. Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children. Keep the pipettes in the original packaging until ready for use and dispose of used pipettes immediately.

For animal treatment only.

#### Other precautions

The product may have adverse effects on painted, varnished or other household surfaces or furnishings. Allow the application site to dry before permitting contact with such materials.

#### Pregnancy and lactation:

Laboratory studies using fipronil and pyriproxyfen have not shown any evidence of teratogenic or embryotoxic effect. Studies have not been carried out with this product in pregnant and lactating queens. Use in pregnant and lactating animals only in accordance with a benefit/risk assessment performed by the responsible veterinarian.

#### Interaction with other medicinal products and other forms of interaction:

None known.

#### Overdose (symptoms, emergency procedures, antidotes):

No serious adverse effects were observed in a target animal safety study in 10-week old kittens treated with up to 5 times the maximum recommended dose 3 times at intervals of 4 weeks and with the maximum recommended dose 6 times at intervals of 4 weeks.

The risk of experiencing adverse reactions (see section 6) may however increase with overdosing, so animals should always be treated with correct pipette size according to bodyweight.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

Do not contaminate ponds, waterways or ditches with the veterinary medicinal product or empty container as this may be dangerous for fish and other aquatic organisms.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

**15. OTHER INFORMATION**

**Pyriproxyfen** is an insect growth regulator (IGR) of the class of compounds known as juvenile hormone analogues. It prevents, by contact, the emergence of adult insects by blocking the development of eggs (ovicidal effect), larvae and pupae (larvicidal effect), which are subsequently eliminated.

**Combination of fipronil and pyriproxyfen** provides an insecticidal and acaricidal activity against fleas (*Ctenocephalides felis*), ticks (*Rhipicephalus turanicus*, *Ixodes ricinus*) in addition to preventing flea eggs developing into adult fleas.

Such combination provides an integrated flea control which can be used against infestations with fleas alone or in association with ticks.

*The following information is intended for veterinary healthcare professionals:*

Incompatibilities:

None known.

Boxes of 1, 2, 3, 4, and 6 pipettes.

Not all pack sizes may be marketed.