

| Identification number of the additive | Name of the holder of authorisation | Additive (Trade name)     | Composition, chemical formula, description, analytical methods   | Species or category of animal                | Maximum age              | Minimum content      | Maximum content       | Other provisions  |                  | End of period of authorisation   | Provisional maximum residue limits (MRLs) in the relevant foodstuffs of animal origin |
|---------------------------------------|-------------------------------------|---------------------------|--|--|--------------------------|----------------------|-----------------------|---|------------------|--|---|
|                                       |                                     |                           |  |  |                          |                      |                       | mg of active substance/kg of complete feedingstuff with a moisture content of 12 %  |                  |  |   |
| 51701                                 | Huvepharma NV Belgium               | Monensin sodium (Coxidin) | <p><i>Additive composition</i></p> <p>Monensin sodium technical substance equivalent to monensin activity: 25 %</p> <p>Perlite: 15 – 20 %</p> <p>Calcium Carbonate: qs. 100 %</p> <p>Active substance</p> $\text{C}_{36}\text{H}_{61}\text{O}_{11}\text{Na}$ | <p>Chickens for fattening</p> <p>Turkeys</p> | <p>—</p> <p>16 weeks</p> | <p>100</p> <p>60</p> | <p>125</p> <p>100</p> | <p>1. Use prohibited at least 1 day before slaughter.</p> <p>2. The additive shall be incorporated in compound feed-stuffs in form of a granulated premixture.</p> <p>3. Monensin sodium shall not be mixed with other coccidiostats.</p> <p>4. Indicate in the instructions for use:</p> <p>"Dangerous for equines. This feedingstuff contains an ionophore; avoid simultaneous administration with tiamulin and monitor for possible adverse reactions when used concurrently with other medicinal substances".</p> | <p>10.6.2021</p> | <p>25 µg monensin sodium/kg of wet skin + fat</p> <p>8 µg monensin sodium/kg of wet liver, wet kidney and wet muscle</p> |   |

(<sup>1</sup>) Details of the analytical methods are available at the following address of the European Union Reference Laboratory for Feed Additives: [http://jmmn.jrc.ec.europa.eu/EURLs/EURL\\_feed\\_additives/Pages/index.aspx](http://jmmn.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx)

## ANNEX

## 'ANNEX

| Identification number of the additive   | Name of the holder of authorisation | Additive (Trade name)     | Composition, chemical formula, description, analytical method   | Species or category of animal                | Maximum age              | Minimum content mg of active substance/kg of complete feedingstuff with a moisture content of 12 % | Other provisions      | End of period of authorisation  | Provisional maximum residue limits (MRLs) in the relevant foodstuffs of animal origin  |   |
|---|-------------------------------------|---------------------------|---|--|--------------------------|--|-----------------------|---|--|---|
| <b>Coccidiostats and histomonostats</b> |                                     |                           |   |  |                          |  |                       |   |  |   |
| 51701                                   | Huvepharma NV Belgium               | Monensin sodium (Coxidin) | <p>Additive composition</p> <p>Monensin sodium technical substance equivalent to monensin activity: 25 %</p> <p>Perlite: 15 – 20 %</p> <p>Wheat bran 55 – 60 %</p> <p>Active substance</p> <p><math>C_{36}H_{61}O_{11}Na</math></p> <p>Sodium salt of polyether monocarboxylic acid produced by <i>Streptomyces dimamomensis</i> 28682, LMG S-19095 in powder form</p> <p>Factor composition</p> <p>Monensin A: not less than 90 %</p> <p>Monensin A + B: not less than 95 %</p> <p>Monensin C: 0,2 – 0,3 %</p> <p>Analytical method <sup>(1)</sup></p> <p>Method for determination of the active substance: high performance liquid chromatography (HPLC) with post-column derivatisation and UV detection <math>\lambda = 520\text{nm}</math></p> | <p>Chickens for fattening</p> <p>Turkeys</p> | <p>—</p> <p>16 weeks</p> | <p>100</p> <p>60</p>   | <p>125</p> <p>100</p> | <p>1. Use prohibited at least 1 day before slaughter.</p> <p>2. The additive shall be incorporated in compound feed-stuff in form a premixture.</p> <p>3. Maximum permitted dose of monensin sodium in complementary feedingstuffs:</p> <ul style="list-style-type: none"> <li>— 625 mg/kg for chickens for fattening,</li> <li>— 500 mg/kg for turkeys.</li> </ul> <p>4. Monensin sodium shall not be mixed with other coccidiostats.</p> <p>5. Indicate in the instructions for use:</p> <p>"Dangerous for equines. This feedingstuff contains an ionophore: avoid simultaneous administration with tiamulin and monitor for possible adverse reactions when used concurrently with other medicinal substances".</p> <p>6. Wear suitable protective clothing, gloves and eye/face protection. In case of insufficient ventilation in the premise, wear suitable respiratory equipments.</p> | <p>6.2.2017</p> <p>8 µg monensin sodium/kg of wet liver, wet kidney and wet muscle</p> | <p>25 µg monensin sodium/kg of wet skin + fat</p> |