

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Awazom 800 mg/g powder for use in drinking water for chickens, ducks and turkeys

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each gram contains:

Active substance:

Amoxicillin trihydrate 800 mg (corresponds to 697 mg of amoxicillin).

Excipients:

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for use in drinking water.

White to pale yellow powder.

4. CLINICAL PARTICULARS

4.1 Target species

Chicken (broiler, pullet, breeder), duck (broiler, breeder), turkey.

4.2 Indications for use, specifying the target species

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

4.3 Contraindications

Do not use in horses, rabbits, hamsters, gerbils and guinea pigs or any other small herbivores.

Do not use in cases of known hypersensitivity to penicillins or other β -lactam antibiotics or to any of the excipients.

Do not use in the presence of β -lactamase-producing bacteria.

4.4 Special warnings for each target species

None.

4.5 Special precautions for use

Special precautions for use in animals

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria. Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the amoxicillin and may decrease the effectiveness of the treatment.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosprins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross sensitivity to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

People with known hypersensitivity to beta-lactam antibiotics should avoid handling the product.

Handle this product with care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as skin rash, you should seek medical advice and show this warning to the physician. Swelling of the face, lips or eyes, or difficulty with breathing are more serious symptoms and require urgent medical attention.

During preparation and administration of the medicated drinking water, avoid skin and eye contact and inhalation of dust particles, as this product may be irritating. Wear impervious gloves and an appropriate dust mask (either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator conforming to European Standard EN140 with a filter to EN143) when mixing and handling the product.

In the event of eye or skin contact, rinse the affected area with large amounts of clean water.

Do not smoke, eat or drink while handling the product.

Wash hands after use.

4.6 Adverse reactions (frequency and seriousness)

Penicillins and cephalosporins may cause hypersensitivity following administration. Allergic reactions to these substances may occasionally be serious.

4.7 Use during pregnancy, lactation or lay

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

4.8 Interaction with other medicinal products and other forms of interaction

Amoxicillin exerts its bactericidal action by inhibition of bacterial cell wall synthesis during multiplication. It is therefore in principle not compatible with bacteriostatic antibiotics (e.g. tetracyclines) which inhibit multiplication. Synergism occurs with β -lactam antibiotics and aminoglycosides.

4.9 Amounts to be administered and administration route

In drinking water use.

Chickens

The recommended dosage is 15 mg amoxicillin trihydrate (equivalent to 18.8 mg veterinary medicinal product) per kg body weight for 3 days or in severe cases for 5 days.

Ducks

Recommended dosage is 20 mg amoxicillin trihydrate (equivalent to 25 mg veterinary medicinal product) per kg body weight for 3 consecutive days.

Turkeys

Recommended dosage is 15-20 mg amoxicillin trihydrate (equivalent to 18.8 to 25 mg veterinary medicinal product) per kg body weight for 3 days or in severe cases for 5 days.

For the preparation of medicated water the body weight of the animals to be treated and their actual daily water consumption should be taken into account. Consumption may vary depending on factors like species, age, state of health, breed and husbandry system (e.g. different temperature, different light regimes). In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted accordingly.

The following formula may be used to calculate the required amount of veterinary medicinal product in mg per litre of drinking water:

$$\frac{\text{x mg product per kg bodyweight per day}}{\text{mean daily water consumption (L) per animal}} \times \text{mean body weight (kg) of animals to be treated} = \text{x mg product per litre drinking water}$$

To ensure a correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

The use of suitably calibrated weighing scales is recommended for measuring the calculated amount of the product.

Preparation of medicated water should provide an amount to be consumed within the next 12 hours. Any unused medicated water should be discarded after 12 hours, and freshly medicated water for the next 12 hours should be prepared. Maximum solubility of the product in water between 5 °C and 20 °C is approximately 6 g/l. The complete dissolution of the powder should be ensured. Appearance of the product after dilution: colorless to pale yellowish solution. The proportioner setting should be changed accordingly. In target animal species, water uptake may vary due to various factors, including environmental temperature, age and type of feed.

Make sure the animals do not have access to non-medicated water during the period when the medicated water is given. When all medicated water has been consumed, turn on the normal water supply again. After the end of the medication period the

water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Not known.

4.11 Withdrawal period(s)

Chickens (meat and offal): 1 day

Ducks (meat and offal): 9 days

Turkeys (meat and offal): 5 days

The product is not authorised for use in laying birds producing eggs for human consumption and within 3 weeks of onset of laying.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: beta-lactam antibacterials, penicillins, penicillins with extended spectrum

ATCvet code: QJ01CA04

5.1 Pharmacodynamic properties

Amoxicillin is a time-dependent bactericidal antibiotic belonging to the semisynthetic penicillin group which acts by inhibiting the synthesis of bacterial cell walls during bacterial replication. It has a broad spectrum of activity against Gram positive and Gram negative bacteria, and owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

There are three main mechanisms of resistance to beta-lactams: beta-lactamase production, production of penicillin binding proteins (PBP), and decreased penetration of the outer membrane. One of the most important is the inactivation of penicillin by beta-lactamase enzymes produced by certain bacteria. These enzymes are capable of cleaving the beta-lactam ring of penicillins, making them inactive. The beta-lactamase could be encoded in chromosomal or plasmidic genes. Cross-resistance is observed between amoxicillin and other penicillins, particularly with aminopenicillins. Observed resistance rates are variable.

5.2 Pharmacokinetic particulars

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium carbonate monohydrate
Sodium citrate
Silica, colloidal anhydrous

6.2 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

6.3 Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf-life after first opening the immediate packaging:

100 g pack: 1 month.

250 g, 500 g and 1000 g packs: 2 months.

Shelf life after dilution according to directions: 12 hours.

6.4 Special precautions for storage

Once opened, the medicinal product should be stored at temperatures below 25°C.

In order to protect from moisture, store the product in the original packaging.
Once opened, keep the bags tightly closed by folding the cut edge of the bag over and securing with a clip.

6.5 Nature and composition of immediate packaging

Thermosealed bags of PET/Al/PE containing 100 g, 250 g, 500 g or 1000 g powder.

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Krka, d.d, Novo mesto
Šmarješka cesta 6
8501 Novo mesto
Slovenia

8. MARKETING AUTHORISATION NUMBER

Vm 01656/4209

9. DATE OF FIRST AUTHORISATION

09 April 2019

10. DATE OF REVISION OF THE TEXT

July 2023

A handwritten signature in black ink, appearing to read 'Dennett', written in a cursive style.

Approved: 14 July 2023