



**United Kingdom  
Veterinary Medicines Directorate  
Woodham Lane  
New Haw  
Addlestone  
Surrey KT15 3LS**

**(Reference Member State)**

**DECENTRALISED PROCEDURE**

**PUBLICLY AVAILABLE ASSESSMENT REPORT FOR A VETERINARY  
MEDICINAL PRODUCT**

**Afamed Fipronil 67 mg spot-on solution for small dogs (UK)  
Afamed Fipronil 134 mg spot-on solution for medium dogs (UK)  
Afamed Fipronil 268 mg spot-on solution for large dogs (UK)  
Afamed Fipronil 402 mg spot-on solution for very large dogs (UK)**

**Fipralone 67 mg spot-on solution for small dogs (IT) (NL)  
Fiproaline 134 mg spot-on solution for medium dogs (IT) (NL)  
Fipralone 268 mg spot-on solution for large dogs (IT) (NL)  
Fiproalone 402 mg spot-on solution for very large dogs (IT) (NL)**

**Fipromedic 67 mg spot-on solution for small dogs (FR)  
Fipromedic 134 mg spot-on solution for medium dogs (FR)  
Fipromedic 268 mg spot-on solution for large dogs (FR)  
Fipromedic 402 mg spot-on solution for very large dogs (FR)**

**PuAR correct as of 14/09/2018 when RMS was transferred to FR. Please  
contact the RMS for future updates.**

Afamed Fipronil 67 mg spot on solution for small dogs  
UK/V/0307/001/DC  
Afamed Fipronil 134 mg spot on solution for medium dogs  
UK/V/0307/002/DC  
Afamed Fipronil 268 mg spot on solution for large dogs  
UK/V/0307/003/DC  
Afamed Fipronil 402 mg spot on solution for very large dogs  
UK/V/0307/004/DC  
Francodex Sante Animale S.A.S.

Application for Decentralised Procedure  
Publicly Available Assessment Report

## MODULE 1

### PRODUCT SUMMARY

|  |  |
|--|--|
| EU Procedure number                    | UK/V/0307/001/DC<br>UK/V/0307/002/DC<br>UK/V/0307/003/DC<br>UK/V/0307/004/DC   |
| Name, strength and pharmaceutical form | Afamed Fipronil 67 mg spot on solution for small dogs<br>Afamed Fipronil 134 mg spot on solution for medium dogs<br>Afamed Fipronil 268 mg spot on solution for large dogs<br>Afamed Fipronil 402 mg spot on solution for very large dogs  |
| Applicant                              | Francodex Sante Animale S.A.S.   |
| Active substance(s)                    | Fipronil   |
| ATC Vetcode                            | QP53AX15   |
| Target species                         | Dogs   |
| Indication for use                     | Treatment of flea ( <i>Ctenocephalides</i> spp.) and tick ( <i>Dermacentor reticulatus</i> ) infestations.<br><br>Insecticidal efficacy against new infestations with adult fleas persists for up to 8 weeks. The product has a persistent acaricidal efficacy for up to 4 weeks against ticks ( <i>Rhipicephalus sanguineus</i> , <i>Ixodes ricinus</i> , <i>Dermacentor reticulatus</i> ). If ticks of some species ( <i>Rhipicephalus sanguineus</i> and <i>Ixodes ricinus</i> ) are present when the product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week. |

|  |  |
|--|--|
|  | The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) where this has been previously diagnosed by a veterinary surgeon. |
|--|--|

## MODULE 2

The Summary of Product Characteristics (SPC) for this product is available on the Heads of Medicines Agencies (veterinary) (HMA(v)) website ([www.hma.eu](http://www.hma.eu)).

## MODULE 3

### PUBLIC ASSESSMENT REPORT

|  |  |
|--|--|
| Legal basis of original application                        | Application in accordance with Article 13.3 of Directive 2001/82/EC as amended by Directive 2004/28/EC |
| Date of completion of the original decentralised procedure | 16 April 2009  |
| Concerned Member States for original procedure             | France<br>Italy<br>Netherlands   |

## I. SCIENTIFIC OVERVIEW

The product is for the treatment of flea (*Ctenocephalides* spp.) and tick (*Dermacentor reticulatus*) infestations. Insecticidal efficacy against new infestations with adult fleas persists for up to 8 weeks. The product has a persistent acaricidal efficacy for up to 4 weeks against ticks (*Rhipicephalus sanguineus*, *Ixodes ricinus*, *Dermacentor reticulatus*). If ticks of some species (*Rhipicephalus sanguineus* and *Ixodes ricinus*) are present when the product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) where this has been previously diagnosed by a veterinary surgeon.

The product is produced and controlled using validated methods and tests,

---

which ensure the consistency of the product released on the market. It has been shown that the product can be safely used in the target species; the slight reactions observed are indicated in the SPC<sup>1</sup>. The product is safe for the user,

the consumer of foodstuffs from treated animals and for the environment, when used as recommended. Suitable warnings and precautions are indicated in the SPC. The efficacy of the product was demonstrated according to the claims made in the SPC. The overall benefit/risk analysis is in favour of granting a marketing authorisation.

## II. QUALITY ASPECTS

### A. *Composition*

#### Afamed Fipronil 67 mg spot on solution for small dogs

The product contains 67 mg/pipette fipronil as active substance and butylhydroxyanisole, butylhydroxytoluene, benzyl alcohol and diethyl glycol monoethyl ether as excipients.

The container/closure system is either thermoformed pipettes or polypropylene pipettes. Thermoformed pipette is a multi-layer plastic single-dose pipette containing an extractible volume of 0.67 ml. Various presentations are available from 1 to 150 pipettes. Not all products may be marketed. The internal layers in contact with the product are made of polyacrylonitrile-methacrylate or polyethylene-ethylene vinyl alcohol-polyethylene. The white external complex is composed of polypropylene / cyclic olefine copolymer / polypropylene.<sup>2</sup> The polypropylene pipette is a white polypropylene single-dose pipette containing an extractible volume of 0.67 ml packaged in uncoloured plastic blister composed of polypropylene / cyclic olefine copolymer / polypropylene closed by heat sealing with a thermosealable lacquered aluminium foil and placed in a carton box or blister card.

The particulars of the containers and controls performed are provided and conform to the regulation.

The choice of the formulation is justified.

#### Afamed Fipronil 134 mg spot on solution for medium dogs

---

<sup>1</sup> SPC – Summary of Product Characteristics.

<sup>2</sup> A transparent pipette and new pipette shape were introduced by way of a Variation Procedure, in June and August 2011. This applies to all presentations.

---

The product contains 134 mg/pipette fipronil as active substance and butylhydroxyanisole, butylhydroxytoluene, benzyl alcohol and diethyl glycol monoethyl ether as excipients.

The container/closure system is either thermoformed pipettes or polypropylene pipettes. Thermoformed pipette is a white or transparent multi-layer plastic

single-dose pipette containing an extractible volume of 1.34 ml. The internal layers in contact with the product are made of polyacrylonitrile-methacrylate. The white external complex is composed of polypropylene / cyclic olefine copolymer / polypropylene. The polypropylene pipette is a white polypropylene single-dose pipette containing an extractible volume of 1.34 ml packaged in uncoloured plastic blister composed of polypropylene / cyclic olefine copolymer /

polypropylene closed by heat sealing with a thermosealable lacquered aluminium foil and placed in a carton box or blister card.

#### Afamed Fipronil 268 mg spot on solution for large dogs

The product contains 268 mg/pipette fipronil as active substance and butylhydroxyanisole, butylhydroxytoluene, benzyl alcohol and diethyl glycol monoethyl ether as excipients.

The container/closure system is either thermoformed pipettes or polypropylene pipettes. Thermoformed pipette is a multi-layer plastic single-dose pipette containing an extractible volume of 2.68 ml. The internal layers in contact with the product are made of polyacrylonitrile-methacrylate. The white external complex is composed of polypropylene / cyclic olefine copolymer / polypropylene. The polypropylene pipette is a white polypropylene single-dose pipette containing an extractible volume of 2.68 ml packaged in uncoloured plastic blister composed of polypropylene / cyclic olefine copolymer / polypropylene closed by heat sealing with a thermosealable lacquered aluminium foil and placed in a carton box or blister card.

#### Afamed Fipronil 402 mg spot on solution for very large dogs

The product contains 402 mg/pipette fipronil as active substance and butylhydroxyanisole, butylhydroxytoluene, benzyl alcohol and diethyl glycol monoethyl ether as excipients.

The container/closure system is either thermoformed pipettes or polypropylene pipettes. Thermoformed pipette is a multi-layer plastic single-dose pipette containing an extractible volume of 4.02 ml. The internal layers in contact with the product are made of polyacrylonitrile-methacrylate. The white external complex is composed of polypropylene / cyclic olefine copolymer /

---

polypropylene. The polypropylene pipette is a white polypropylene single-dose pipette containing an extractible volume of 4.02 ml packaged in uncoloured plastic blister composed of polypropylene / cyclic olefine copolymer / polypropylene closed by heat sealing with a thermosealable lacquered aluminium foil and placed in a carton box or blister card.

### ***B. Method of Preparation of the Product***

The product is manufactured fully in accordance with the principles of good manufacturing practice from a licensed manufacturing site.

Process validation data on the product have been presented in accordance with the relevant European guidelines.

### ***C. Control of Starting Materials***

The active substance is fipronil which is almost unabsorbed through the skin and the formulation is designed to deposit the active substance easily onto the animal.

There are four excipients used in the formulation and each has been used previously in veterinary medicines.

The active substance specification is considered adequate to control the quality of the material. Batch analytical data demonstrating compliance with this specification have been provided.

All the excipients used in the final product have monographs in the Ph. Eur. and each comply with the requirements of the current edition of the Ph. Eur.

### ***D. Specific Measures concerning the Prevention of the Transmission of Animal Spongiform Encephalopathies***

There are no substances within the scope of the TSE Guideline present or used in the manufacture of this product.

### ***E. Control on intermediate products***

Not applicable

### ***F. Control Tests on the Finished Product***

---

The finished product specification controls the relevant parameters for the pharmaceutical form. The tests in the specification, and their limits, have been justified and are considered appropriate to adequately control the quality of the product.

Satisfactory validation data for the analytical methods have been provided.

Batch analytical data from the proposed production sites have been provided demonstrating compliance with the specification.

### **G. Stability**

Stability data on the active substance have been provided in accordance with applicable European guidelines, demonstrating the stability of the active substance when stored under the approved conditions.

Stability data on the finished product have been provided in accordance with applicable European guidelines, demonstrating the stability of the product throughout its shelf life when stored under the approved conditions.

### **H. Genetically Modified Organisms**

*Not applicable.*

### **J. Other information**

#### **Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

#### **Special precautions for storage**

Store below 30°C.

Store in a dry place.

Store in the original package.

Do not remove from blister until required for use.

## **III. SAFETY AND RESIDUES ASSESSMENT (PHARMACOTOXICOLOGICAL)**

### **III.A Safety Testing**

### ***Pharmacological Studies***

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not required.

### ***Toxicological Studies***

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not required.

### ***Other Studies***

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not required.

### ***User Safety***

The applicant has made reference to the user risk assessment submitted in the Expert Report which considered dermal and oral exposure including exposure to children. The user warnings proposed are the same as those for the reference product and are considered satisfactory to address user safety.

### ***Ecotoxicity***

The applicant has provided environmental risk assessment in compliance with the relevant guideline which showed that the environmental safety of the product is acceptable.

## **IV CLINICAL ASSESSMENT (EFFICACY)**

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not submitted.

### ***IV.A Pre-Clinical Studies***

#### ***Pharmacology***



---

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not provided.

### ***Pharmacokinetics***

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not provided.

### ***Tolerance in the Target Species of Animals***

The applicant has conducted a target animal tolerance study to evaluate the local and systemic tolerance of the product. The study was conducted on dogs. The dogs were divided into different groups. The study concluded that there were no laboratory or histological changes and no clinical signs associated with the treatment were noted when the product was administered at one, three and five times the expected therapeutic dose for three applications over a period of approximately two and a half months.

### ***Resistance***

Since the application is made in accordance with Article 13(3) of Directive 2004/28/EC, on the basis of essential similarity, data on this section of the dossier were not provided.

## ***IV.B Clinical Studies***

The applicant has provided four dose confirmation studies; one for fleas (*Ctenocephalides felis*) on dogs, one for each of the following tick species (*Rhipicephalus sanguineus*, *Dermacentor reticulatus* and *Ixodes ricinus*) on dogs. The first study was conducted to determine and compare the efficacy of Afamed Fipronil spot-on solution with the reference product. The study was conducted on dogs. The study concluded that both the Afamed Fipronil spot-on solution and the reference product had a therapeutic efficacy of more than 95% at 48 hours after treatment and this persisted for up to eight weeks, when administered to dogs at the doses stated (0.67 ml for dogs of two to 10 kg bodyweight or 1.34 ml for dogs of more than 10 and up to 20 kg bodyweight).

---

The second study was conducted to confirm the efficacy of Afamed Fipronil spot-on solution against the tick *Dermacentor reticulatus* on dogs compared to the reference product when applied topically at the rate of 0.067 to 0.335 ml/kg bodyweight. The study was conducted on dogs. It was concluded that both the Afamed Fipronil spot-on solution and the reference product had an immediate (therapeutic) efficacy, at 48 hours after treatment, of more than 90% against ticks already on the dogs. This level of efficacy against *Dermacentor reticulatus*

persisted for at least four weeks, based on subsequent weekly new tick infestations.

The third study was conducted to determine and compare the efficacy of Afamed Fipronil spot-on solution with the reference product against a French strain of the tick *Rhipicephalus sanguineus*. The study was conducted on dogs. The products were applied at recommended doses. The study demonstrated that the therapeutic efficacy at 48 hours after treatment of both the Afamed Fipronil spot-on solution and the reference product was much less than 90% against ticks already on the dogs. However, by one week after treatment and for up to four weeks the persistent efficacy of both the Afamed Fipronil spot-on solution and the reference product was more than 90% against subsequent weekly new tick infestations of *Rhipicephalus sanguineus*.

The fourth study was conducted to confirm the efficacy of Afamed Fipronil spot-on solution against the tick *Ixodes ricinus* on dogs compared to the reference product when applied once topically at the recommended dose rate. In the study it was demonstrated that the Afamed Fipronil spot-on solution had a therapeutic efficacy at 48 hours after treatment of just under 90% against ticks already on the dog. However, by one week and for at least three weeks after treatment the persistent efficacy of both the Afamed Fipronil spot-on solution and the reference product was more than 90% against subsequent weekly new tick infestations of *Ixodes ricinus*.

## **V OVERALL CONCLUSION AND BENEFIT– RISK ASSESSMENT**

The data submitted in the dossier demonstrate that when the product is used in accordance with the Summary of Product Characteristics, the benefit/risk profile for the target species is favourable and the quality and safety of the product for humans and the environment is acceptable.

## **MODULE 4**

### **POST-AUTHORISATION ASSESSMENTS**

The SPC and package leaflet may be updated to include new information on the quality, safety and efficacy of the veterinary medicinal product. The current SPC is available on the Product Information Database of the Veterinary Medicines Directorate website.

[www.gov.uk/check-animal-medicine-licensed](http://www.gov.uk/check-animal-medicine-licensed)

The post-authorisation assessment (PAA) contains information on significant changes which have been made after the original procedure which are important for the quality, safety or efficacy of the product.

The PAA for this product is available on the Product Information Database of the Veterinary Medicines Directorate website.

[www.gov.uk/check-animal-medicine-licensed](http://www.gov.uk/check-animal-medicine-licensed)