



**ASSURING THE SAFETY, QUALITY AND EFFICACY
OF VETERINARY MEDICINES**

**United Kingdom
Veterinary Medicines Directorate
Woodham Lane
New Haw
Addlestone
Surrey KT15 3LS**

DECENTRALISED PROCEDURE

**PUBLICLY AVAILABLE ASSESSMENT REPORT FOR A VETERINARY
MEDICINAL PRODUCT**

Enrocare 50 mg/ml Solution for Injection for Cattle, Pigs, Dogs and Cats

**PuAR correct as of 16/04/2018 when RMS was transferred
to IE. Please contact the RMS for future updates.**

MODULE 1

PRODUCT SUMMARY

EU Procedure number	UK/V/0334/002/DC
Name, strength and pharmaceutical form	Enrocare 50 mg/ml Solution for Injection for Cattle, Pigs, Dogs and Cats
Applicant	Animalcare Limited
Active substance(s)	Enrofloxacin
ATC Vetcode	QJ01MA90
Target species	Cattle, Pigs weighing more than 25 kg, Dogs and Cats
Indication for use	<p><u>Cattle</u> Diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin (e.g. <i>pasteurellosis</i>, <i>mycoplasmosis</i>, <i>coli-bacillosis</i>, <i>coli-septicaemia</i> and <i>salmonellosis</i>) and secondary bacterial infections subsequent to viral conditions (e.g. viral pneumonia), where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.</p> <p><u>Pigs (weighing more than 25 kg)</u> Diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin (e.g. <i>pasteurellosis</i>, <i>actinobacillosis</i>, <i>mycoplasmosis</i>, <i>coli-bacillosis</i>, <i>coli-septicaemia</i> and <i>salmonellosis</i>) and multifactorial diseases such as atrophic rhinitis and enzootic pneumonia, where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.</p> <p><u>Dogs and cats</u> Treatment of bacterial infections of the alimentary, respiratory and urogenital tracts, skin, secondary wound infections and otitis externa where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.</p>

MODULE 2

The Summary of Product Characteristics (SPC) for this product is available on the Heads of Medicines Agencies (veterinary) (HMA(v)) website (www.hma.eu).

MODULE 3

PUBLIC ASSESSMENT REPORT

Legal basis of original application	Application in accordance with Article 13 (1) of Directive 2001/82/EC as amended by 2004/28/EC.
Date of completion of the original decentralised procedure	22 December 2009
Concerned Member States for original procedure	France Ireland Luxembourg Netherlands

I. SCIENTIFIC OVERVIEW

Enrocare 50 mg/ml solution for injection for cattle, pigs, dogs and cats contains the active substance enrofloxacin. The product is authorised to be used in cattle, pigs weighing more than 25 kg, dogs and cats. In cattle, the product is used in the treatment of diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin (e.g. *pasteurellosis*, *mycoplasmosis*, *coli-bacillosis*, *coli-septicaemia* and *salmonellosis*) and secondary bacterial infections subsequent to viral conditions (e.g. viral pneumonia), where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice. In pigs, it is used in the treatment of diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin (e.g. *pasteurellosis*, *actinobacillosis*, *mycoplasmosis*, *coli-bacillosis*, *coli-septicaemia* and *salmonellosis*) and multifactorial diseases such as atrophic rhinitis and enzootic pneumonia, where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice. In dogs and cats, it is used in the treatment of bacterial infections of the alimentary, respiratory and urogenital tracts, skin, secondary wound infections and otitis externa where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.

This application is a generic made in accordance with Article 13(1) of Directive 2001/82/EC. The reference product is Baytril 5% solution for injection which has been marketed in the UK since 1992.

The product is produced and controlled using validated methods and tests which ensure the consistency of the product released on the market. It has been shown that the product can be safely used in the target species and the slight reactions observed are indicated in the SPC. The product is safe for the user, the consumer of foodstuffs from treated animals and for the environment, when

used as recommended. Suitable warnings and precautions are indicated in the SPC. The efficacy of the product was demonstrated according to the claims made in the SPC.

II. QUALITY ASPECTS

A. Composition

The product contains enrofloxacin as an active substance and n-butyl alcohol, potassium hydroxide and water for injections as excipients.

The product is packaged in an amber Type 1 multidose glass vial with a grey bromobutyl rubber stopper and aluminium overseal.

The particulars of the containers and controls performed are provided and conform to the regulation.

The choice of the formulation is justified.

B. Method of Preparation of the Product

The product is manufactured fully in accordance with the principles of good manufacturing practice from a licensed manufacturing site.

Process validation data on the product have been presented in accordance with the relevant European guidelines.

C. Control of Starting Materials

The active substance enrofloxacin is an established active substance and supporting data have been provided in the form of European Drug Master File (EDMF). It is considered that the manufacturing process is adequately controlled and the active substance specification has been suitably justified.

There are three excipients used in the formulation and each has been used previously in veterinary medicines. Potassium hydroxide and water for injections have monographs in the Ph. Eur. and each complies with the requirements of the current edition of the Ph. Eur. In the absence of a monograph in the European Pharmacopoeia, the requirements of the monograph in the United States Pharmacopoeia National Formulary are applied in the raw material specification for n-butyl alcohol.

D. Specific Measures concerning the Prevention of the Transmission of Animal Spongiform Encephalopathies

A declaration from the applicant has been provided which states that Enrocare 50 mg/ml solution for injection for cattle, pigs, dogs and cats complies with the latest version of the CPMP/CVMP guidelines on TSEs. Additional declarations have also been provided by the manufactures of the active substance and

suppliers of the excipients confirming that no materials of animal origin are used during their manufacture.

E. Control on intermediate products

There are no intermediate products.

F. Control Tests on the Finished Product

The finished product specification controls the relevant parameters for the pharmaceutical form. The tests in the specification, and their limits, have been justified and are considered appropriate to adequately control the quality of the product. Satisfactory validation data for the analytical methods have been provided.

G. Stability

Stability data on the finished product have been provided in accordance with applicable European guidelines, demonstrating the stability of the product throughout its shelf life. The shelf-life of the veterinary medicinal product as packaged for sale is 3 years. An in-use shelf life of 28 days is justified.

H. Genetically Modified Organisms

Not applicable

J. Other Information

Special Precautions for Storage:

Do not store above 25°C

Do not freeze.

Keep the vial in the outer carton in order to protect from light.

Discard unused material.

III. SAFETY AND RESIDUES ASSESSMENT (PHARMACOTOXICOLOGICAL)

III.A Safety Testing

Pharmacological Studies

Since the application is made in accordance with Article 13 (1) of Directive 2001/82/EC as amended by Directive 2004/28/EC, on the basis of being a generic of a reference medicinal product, data on pharmacodynamics and pharmacokinetics are not required.

Toxicological Studies

Since the application is made in accordance with Article 13 (1) of Directive 2001/82/EC as amended by Directive 2004/28/EC, on the basis of being a generic of a reference medicinal product, data on toxicology are not required.

Other Studies

Since the application is made in accordance with Article 13 (1) of Directive 2001/82/EC as amended by Directive 2004/28/EC, on the basis of being a generic of a reference medicinal product, the applicant has not submitted any data for this section.

User Safety

The following operator warnings are included in the SPC and product literature:

- This product is an alkaline solution. Wash any splashes from skin and eyes immediately with water.
- People with known hypersensitivity to (fluoro)quinolones should avoid contact with this product.
- Do not eat, drink or smoke whilst using the product.
- In case of accidental self-injection seek medical advice immediately and show the package leaflet or label to the physician.

Ecotoxicity

The applicant provided a first phase environmental risk assessment in compliance with the relevant guideline. In accordance with the Phase I decision tree, the treatment of dogs and cats does not result in extensive exposure of the environment. Since the product is also indicated to be used in cattle and pigs, the applicant calculated PEC_{soil} values. PEC_{soil} values were calculated for all types of cattle and pigs raised intensively using the equations provided by the CVMP guideline. The default value of 50 % of animals treated for bacterial infections provided by the CVMP guideline was used. PEC_{soil} values are shown below.

Target animal	PEC_{soil} ($\mu\text{g}/\text{kg}$)
Calf	71.4
Dairy cow	40.1
Cattle 0-1 year	63.0
Cattle >2 years	72.9
Fattening pig	73.7
Sow with litter	26.2

The assessment ended at Phase I as all the PEC_{soil} values were below 100 $\mu\text{g}/\text{kg}$. The warnings and precautions as listed on the product literature are adequate to ensure safety to the environment when the product is used as directed.

III.B Residues documentation

Residue Studies

As the product has the same pharmaceutical form and the same qualitative and quantitative composition as the reference product, it was not necessary to provide any data on bioequivalence in accordance with paragraph 4b of the CVMP guidelines for the conduct of bioequivalence studies for veterinary medicinal products (EMA/CVMP/016/00-corr-FINAL).

MRLs

All the excipients are listed in table 1 of regulation (EC) No 37/2010, with the exception of water for injections.

MRLs are listed below:

	Bovine	Porcine
Muscle	100 µg/kg	100 µg/kg
Liver	300 µg/kg	200 µg/kg
Kidney	200 µg/kg	300 µg/kg
Fat	100 µg/kg	100 µg/kg
Milk	100 µg/kg	N/A

Withdrawal Periods

Calves:

Following intravenous injection: Meat and offal: 5 days.

Following subcutaneous injection: Meat and offal: 12 days.

Not authorised for use in animals producing milk for human consumption.

Pigs:

Meat and offal: 13 days.

IV CLINICAL ASSESSMENT (EFFICACY)

Pharmacology

Since the application is made in accordance with Article 13 (1) of Directive 2001/82/EC as amended by Directive 2004/28/EC, on the basis of being a generic of a reference medicinal product, data on pharmacodynamics and pharmacokinetics are not required.

Tolerance in the Target Species of Animals

As the product has the same pharmaceutical form and the same qualitative and quantitative composition as the reference product, no new target species tolerance data have been presented. This complies with exemptions specified under Article 13 (2)(b) of Directive 2001/82/EC as amended by 2004/28/EC.

Resistance

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials. Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to potential cross-resistance.

Adequate warnings and precautions appear on the product literature.

IV.B Clinical Studies

Since the application is made in accordance with Article 13 (1) of Directive 2001/82/EC as amended by Directive 2004/28/EC, on the basis of being a generic of a reference medicinal product, data on clinical efficacy are not required.

V OVERALL CONCLUSION AND BENEFIT– RISK ASSESSMENT

The data submitted in the dossier demonstrate that when the product is used in accordance with the Summary of Product Characteristics, the benefit risk profile for the target species is favourable and the quality and safety of the product for humans and the environment is acceptable.

MODULE 4

POST-AUTHORISATION ASSESSMENTS

The SPC and package leaflet may be updated to include new information on the quality, safety and efficacy of the veterinary medicinal product. The current SPC is available on the Product Information Database of the Veterinary Medicines Directorate website.

www.gov.uk/check-animal-medicine-licensed

The post-authorisation assessment (PAA) contains information on significant changes which have been made after the original procedure which are important for the quality, safety or efficacy of the product.

The PAA for this product is available on the Product Information Database of the Veterinary Medicines Directorate website.

www.gov.uk/check-animal-medicine-licensed