

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Box of the vials of 200 ml, 500 ml and 1000 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Virbamec Super 10 mg/ml, 100 mg/ml Solution for Injection Ivermectin and Clorsulon



**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Active substances

Ivermectin .....	10	mg/ml
Clorsulon .....	100	mg/ml

**3. PHARMACEUTICAL FORM**

Solution for Injection

**4. PACKAGE SIZE**

200 ml  
500 ml  
1000 ml

**5. TARGET SPECIES**

**6. INDICATION(S)**

For the treatment of mixed trematode and nematode or arthropod infestations, due to adult and immature roundworms, lungworms, warbles, mites, lice and liver fluke in cattle.

Read package leaflet before use.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous route only.

The product should be given once by subcutaneous injection at the recommended dosage level of 200 µg of ivermectin and 2 mg of clorsulon per kilogram of bodyweight. Each ml contains 10 mg of ivermectin and 100 mg of clorsulon, sufficient

to treat 50 kg of bodyweight. Subcutaneous injection only.

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

Divide doses greater than 10 ml between two injection sites.

Inject under the loose skin behind the shoulder. Use of a 17-gauge, 1/2-inch (15-20 mm) needle is suggested. The injection may be given with any standard automatic, multidose or single-dose hypodermic syringe. If using a hypodermic syringe, use a separate sterile needle to withdraw the product from the pack.

This product does not contain an antimicrobial preservative. Swab septum before removing each dose. Use a dry, sterile needle and syringe.

For 200, 500 and 1000-ml pack sizes, use only automatic-syringe equipment. Injection on animals with wet or dirty hides is not recommended.

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.

When the temperature of the product is below 5°C, difficulty in administration may be encountered due to increased viscosity. Warming the product and injection equipment to about 15°C will greatly increase the ease with which the product can be injected. Different injection sites should be used for other parenteral products.

The timing for treatment should be based on epidemiological factors and should be customised for each individual farm. A dosing programme should be established by the veterinary surgeon.

## **8. WITHDRAWAL PERIOD**

Meat and offal: 66 days.

Milk: do not use in animals producing milk for human consumption.

Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

## **9. SPECIAL WARNING(S), IF NECESSARY**

### **CONTRAINDICATIONS**

Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving. Do not use in species other than cattle as severe adverse reactions, including fatalities, may occur in dogs for example. Do not use by the intravenous or intramuscular route. Do not use in animals known to be hypersensitive to the active substance.

### **UNDESIRABLE EFFECTS**

Transitory discomfort has been observed in some cattle following subcutaneous administration. Soft-tissue swelling and/or slight pain at the injection site has also been observed. These reactions have disappeared without treatment. In case of hypersensitivity reactions a symptomatic treatment should be applied.

## USER WARNINGS

Do not smoke, drink or eat whilst handling the product. Wash hands after use. Avoid contact with skin and eyes.

Take care to avoid self-administration: the product may cause irritation and/or pain at the site of injection. In case of accidental self-injection, seek medical advice and show the label to the doctor.

## SPECIAL WARNINGS

Care should be taken to avoid the following practices because they increase the risk of the development of resistance and could ultimately result in ineffective therapy.

## OTHER PRECAUTIONS

Ivermectin is highly toxic to aquatic invertebrates. Treated cattle should not have direct access to ponds, streams or ditches for 14 days after treatment to avoid adverse effects on aquatic organisms.

## USE DURING PREGNANCY, LACTATION OR LAY

Do not use in animals producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

## SPECIAL PRECAUTIONS FOR USE

Divide doses greater than 10 ml between two injection sites to reduce occasional discomfort or site reaction. Frequent and repeated use may lead to the development of resistance.

The timing of treatment for the parasitic stages of warbles should be chosen carefully. The best time to treat against infections with *Hypoderma* is immediately after the end of the swarming of the warbles, before the larvae cause damage in the body of the animal (October to November). If larvae of *Hypoderma bovis* are killed during migration through the spine, this may induce posterior paralysis and recumbency. These reactions occur mainly when animals are treated between December and March. Avermectins may not be well tolerated in non-target species. Cases of intolerance resulting in fatalities have been reported in dogs, especially collies, old English sheep dogs and related breeds or crosses and also in turtles/tortoises.

## OVERDOSE

An acute toxic syndrome consisting of CNS signs of depression and listlessness, ataxia, recumbency and possible death occurs in cattle given SQ doses equal to 40 times the therapeutic dose for ivermectin. Treatment should be symptomatic. A toxic-syndrome dose level has not been identified in cattle for clorsulon.

## 10. EXPIRY DATE

EXP {month/year}

Shelf-life after first opening the immediate packaging: 28 days Once broached use

by:.....

## **11. SPECIAL STORAGE CONDITIONS**

Avoid the introduction of contamination during use. Should any apparent growth or discolouration occur, the product should be discarded.

Protect from light.

Store in the original container.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

EXTREMELY DANGEROUS to fish and aquatic life.

Do not contaminate ponds, waterways or ditches with the product or empty container.

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

*[Distribution category]*

FOR ANIMAL TREATMENT ONLY

To be supplied only on veterinary prescription POM-VPS

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN

## **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

VIRBAC  
1ère avenue 2065m LID  
06516 Carros  
France

## **16. MARKETING AUTHORISATION NUMBER(S)**

Vm 05653/4155

## **17. MANUFACTURER’S BATCH NUMBER**

Lot: number

Distributor: VIRBAC Ltd Suffolk IP30 9UP – UK

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**Label of vials of 200 ml, 500 ml and 1000 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Virbamec Super 10 mg/ml, 100 mg/ml, solution for injection

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Active substances:  
ivermectin 10 mg/ml  
clorsulon 100 mg/ml  
Excipient:  
propyl gallate (E310) 0.2 mg/ml.

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE**

200 ml  
500 ml  
1000 ml

**6. INDICATION(S)**

For the treatment of mixed trematode and nematode or arthropod infestations, due to adult and immature roundworms, lungworms, warbles, mites, lice and liver fluke in cattle.

Read package leaflet before use.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Inject once subcutaneously, at the dosage of 1 ml per 50 kg of bodyweight. Read package leaflet before use.

**8. WITHDRAWAL PERIOD**

Meat and offal: 66 days.

Milk: do not use in animals producing milk for human consumption.

Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

**9. SPECIAL WARNING(S), IF NECESSARY**

CONTRA-INDICATIONS, UNDESIRABLE EFFECTS, SPECIAL PRECAUTIONS FOR USE, USE DURING PREGNANCY AND LACTATION, OVERDOSE, USER WARNINGS  
Read package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}  
Shelf-life after first opening the immediate packaging: 28 days  
Once broached use by: .....

**11. SPECIAL STORAGE CONDITIONS**

Avoid the introduction of contamination during use. Should any apparent growth or discolouration occur, the product should be discarded. Protect from light. Store in the original container.

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.  
EXTREMELY DANGEROUS to fish and aquatic life.  
Do not contaminate ponds, waterways or ditches with the product or empty container.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**  
*[Distribution category]*

FOR ANIMAL TREATMENT ONLY.  
To be supplied only on veterinary prescription. POM-VPS

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

VIRBAC  
1ère avenue 2065m LID  
06516 Carros  
France

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 05653/4155

**17. MANUFACTURER'S BATCH NUMBER**

Lot: number

Distributor: VIRBAC Ltd Suffolk IP30 9UP – UK

**PACKAGE LEAFLET  
FOR:**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION  
HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder and manufacturer responsible for batch release:  
VIRBAC  
1ère avenue 2065m LID  
06516 Carros  
France

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Virbamec Super 10 mg/ml, 100  
mg/ml Solution for Injection

**3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER  
INGREDIENTS**

<u>Active substances</u>		
Ivermectin.....	10	mg/ml
Clorsulon .....	100	mg/ml

Other substance  
Propyl gallate (E310)..... 0.2mg/ml

Clear, slightly yellow and slightly viscous solution

**4. INDICATIONS**

For the treatment of mixed trematode and nematode or arthropod infestations, due to adult and immature roundworms, lungworms, warbles, mites, lice and liver fluke in cattle.

Gastro-intestinal roundworms (adult and fourth-stage larvae) *Ostertagia ostertagi* (including inhibited larval stages) *Ostertagia lyrata*  
*Haemonchus placei*  
*Trichostrongylus axei*  
*Trichostrongylus colubriformis* *Cooperia oncophora* *Cooperia punctata*  
*Cooperia pectinata*  
*Bunostomum phlebotomum*  
*Oesophagostomum radiatum*  
*Strongyloides papillosus*  
(adult) *Nematodirus*



*helvetianus* (adult)  
*Nematodirus spathiger*  
(adult)  
Lungworms (adult and fourth-stage larvae):  
*Dictyocaulus*  
*viviparus* Liver fluke  
(adult): *Fasciola*  
*hepatica*

Warbles (parasitic stages): *Hypoderma*  
*bovis* *Hypoderma*  
*lineatum* Mange mites:  
*Psoroptes bovis*  
*Sarcoptes scabiei* var. *bovis*  
Sucking lice:  
*Linognathus vituli*  
*Haematopinus*  
*eurysternus*

The product may also be used as an aid in the control of the mange mite *Chorioptes bovis*, but complete elimination may not occur.

## 5. CONTRA-INDICATIONS

Not for use in species other than cattle as severe adverse reactions, including fatalities, may occur in dogs for example.

Do not use by the intravenous or intramuscular route.

Do not use in animals known to be hypersensitive to the active substance.

## 6. ADVERSE REACTIONS

Transitory discomfort has been observed in some cattle following subcutaneous administration. Soft-tissue swelling and/or slight pain at the injection site has also been observed. These reactions have disappeared without treatment. In case of hypersensitivity reactions a symptomatic treatment should be applied

## 7. TARGET SPECIES

Cattle

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

The product should be given once by subcutaneous injection at the recommended dosage level of 200 µg of ivermectin and 2 mg of clorsulon per kilogram of bodyweight. Each ml contains 10 mg of ivermectin and 100 mg of clorsulon, sufficient to treat 50 kg of bodyweight.

Subcutaneous injection only.

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or over dosing.

Divide doses greater than 10 ml between two injection sites. Inject under the loose skin behind the shoulder. Use of a 17-gauge, ½-inch (15-20 mm) needle is suggested. The injection may be given with any standard automatic, multidose or single-dose hypodermic syringe. If using a hypodermic syringe, use a separate sterile needle to withdraw the dose from the pack.

This product does not contain an antimicrobial preservative. Swab septum before removing each dose. Use a dry, sterile needle and syringe. For 200, 500 and 1000-ml pack sizes, use only automatic-syringe equipment. Injection on animals with wet or dirty hides is not recommended.

When the temperature of the product is below 5 °C, difficulty in administration may be encountered due to increased viscosity. Warming the product and injection equipment to about 15 °C will greatly increase the ease with which the product can be injected. Different injection sites should be used for other parenteral products.

The timing for treatment should be based on epidemiological factors and should be customised for each individual farm. A dosing programme should be established by the veterinary surgeon.

## **SPECIAL WARNINGS**

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## **9. ADVICE ON CORRECT ADMINISTRATION**

## 10. WITHDRAWAL PERIOD(S)

Meat and offal : 66 days.

Milk: do not use in animals producing milk for human consumption.

Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

## 11. SPECIAL STORAGE PRECAUTIONS

Avoid the introduction of contamination during use. Should any apparent growth or discolouration occur, the product should be discarded.

When the container is broached (opened) for the first time, using the in-use shelf life which is specified on this package leaflet, the date on which any product remaining in the carton should be discarded should be worked out. This discard date should be written in the space provided.

Protect from light.

Store in the original container.

Do not use this veterinary medicinal product after the expiry date stated on the carton and vial after "EXP". The expiry date refers to the last day of that month.

Shelf-life after first opening the immediate packaging: 28 days.

## 12. SPECIAL

### WARNING(S) USER

#### WARNINGS

Do not smoke, drink or eat while handling the product. Wash hands after use.

Avoid contact with skin and eyes.

Take care to avoid self-administration: the product may cause irritation and/or pain at the site of injection. In case of accidental self-injection, seek medical advice and show the label to the doctor.

#### SPECIAL PRECAUTIONS FOR USE

Divide doses greater than 10 ml between two injection sites to reduce occasional discomfort or site reaction.

Frequent and repeated use may lead to the development of resistance.

The timing of treatment for the parasitic stages of warbles should be chosen carefully. The best time to treat against infections with *Hypoderma* is immediately after the end of the swarming of the warbles, before the larvae cause damage in the body of the animal (October to November). If larvae of *Hypoderma bovis* are killed during migration through the spine, this may induce posterior paralysis and recumbency. These reactions occur mainly when animals are treated between December and March.

Avermectins may not be well tolerated in non-target species. Cases of intolerance resulting in fatalities have been reported in dogs, especially collies, old English sheep dogs and related breeds or crosses and also in turtles/tortoises.

### **OTHER PRECAUTIONS**

Ivermectin is highly toxic to aquatic invertebrates. Treated cattle should not have direct access to ponds, streams or ditches for 14 days after treatment to avoid adverse effects on aquatic organisms.

### **USE DURING PREGNANCY, LACTATION OR LAY**

Do not use in animals producing milk for human consumption.  
Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

### **OVERDOSE**

An acute toxic syndrome consisting of CNS signs of depression and listlessness, ataxia, recumbency and possible death occurs in cattle given SQ doses equal to 40 times the therapeutic dose for ivermectin. Treatment should be symptomatic.  
A toxic-syndrome dose level has not been identified in cattle for clorsulon.

### **INTERACTIONS**

Not known.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

EXTREMELY DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the product or empty container.

### **14. PID LINK (Do not print heading)**

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

### **15. OTHER INFORMATION PRESENTATIONS**

Sizes: 200-ml, 500-ml and 1L; multidose colourless-plastic vials. Not all pack sizes may be marketed.

### **FOR ANIMAL TREATMENT ONLY KEEP OUT OF THE SIGHT AND REACH OF CHILDREN**

To be supplied only on veterinary prescription. Vm 05653/4155  
POM-VPS

Distributor: VIRBAC Ltd - Suffolk IP30 9UP – UK

Revised: April 2024  
AN: 01199/2023 & 01201/2023

Approved 11 April 2024

A handwritten signature in black ink, consisting of a stylized initial followed by the name "Hunter." with a period.