

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING WHERE THERE IS NO PACKAGE LEAFLET, i.e. Combined label and package leaflet
5 Litre Container

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Vectocert 1.25% w/v Pour-On Solution for Sheep.

2. COMPOSITION

1.25% w/v cypermethrin
0.02% Green s Dye (E142)

Clear blue non-aqueous solution.

3. PACKAGE SIZE

1 Litre, 2.5 Litres, 5 Litres

4. TARGET SPECIES

Sheep

5. INDICATIONS FOR USE

Indications for use

This veterinary medicinal product is for prevention and treatment of blowfly strike on sheep. Also used for the treatment and control of headflies, and treatment of ticks and biting lice on sheep.

6. CONTRAINDICATIONS

Contraindications

Not to be used in lambs less than one week old.

7. SPECIAL WARNINGS

Special warnings

Special warnings:

Contains a blue dye to assist with the identification of treated sheep.

Special precautions for safe use in the target species:

Take care not to apply in the sheep's eyes. This precaution should be observed particularly when treating breeds with little wool on their heads.

Toxic signs in mammals are tremors, hyperexcitability, salivation, extreme limb movements and paralysis, rarely leading to death. Usually the signs disappear rapidly and the animals recover, generally within a week. Symptomatic therapy can be given if necessary.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product is harmful if swallowed and may cause skin, eye or respiratory irritation. This veterinary medicinal product may also cause hypersensitivity (allergic) reactions. People with known hypersensitivity to cypermethrin should avoid contact with the veterinary medicinal product.

Handle this veterinary medicinal product with great care to avoid accidental exposure.

Personal protective equipment consisting of eye protection, protective clothing, rubber gloves and boots should be worn when applying the veterinary medicinal product. Wear a disposable face mask when applying as a fan-spray for the prevention of blowfly strike.

Make sure when attaching the applicator gun onto the container that the coiled retainer is secured onto the cap and the applicator.

Use in a well-ventilated area and avoid inhaling the vapour.
Do not eat, drink or smoke whilst using the veterinary medicinal product.

Wash splashes from skin and eyes immediately with plenty of clean water. If irritation persists seek medical advice immediately and show the label to the physician.
Remove contaminated clothing immediately and wash exposed skin with plenty of clean water.

Wash hands and exposed skin before eating, drinking or smoking and after work.
In case of accidental ingestion or mouth contact, immediately rinse the mouth with plenty of water and seek medical advice.

Avoid handling treated animals.

Special precautions for the protection of the environment:

As a precautionary measure sheep must be kept away from watercourses for at least one hour following treatment.

Pregnancy and lactation:

No restrictions. Pregnant ewes must be handled with care.

Interactions with other medicinal products and other forms of interaction:

The veterinary medicinal product must be applied only with the recommended applicator gun, as the product may have a detrimental effect on certain components of conventional dosing guns.

Overdose:

Toxic signs in mammals are tremors, hyperexcitability, salivation, choreoathetosis and paralysis, rarely leading to death. Usually, the signs disappear rapidly and the animals recover, generally within a week.

Major incompatibilities:

None known.

8. ADVERSE EVENTS

Adverse events

Sheep:

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details on this label, or via your national reporting system at: Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

9. DOSAGE FOR EACH TARGET SPECIES, ROUTES AND METHOD OF ADMINISTRATION

Dosage for each species, routes and method of administration

Pour-On Solution

Shake well before use.

Vectocert 1.25% w/v Pour-On must be applied only with a recommended applicator gun as the product may have a detrimental effect on certain components of standard applicator guns. The operating and maintenance instructions, including details on the use of nozzles, are supplied with each applicator gun. To ensure accurate dosage of the product, maintain applicator gun carefully.

Any surplus product remaining in the applicator gun must be expelled before the applicator gun is stored.

Vectocert Pour-On minimises mothering-up problems after treatment. Nevertheless, care must be taken not to apply the product right to the lamb's tail as a ewe recognises a young lamb by the smell of the tail area.

This product contains a blue dye to aid in the identification of treated animals. The colour is temporary and will fade from the fleece within 48 hours of application.

Blowflies


Prevention of blowfly strike

BODYWEIGHT	DOSE	DOSES PER PACK
Up to 25 kg*	20 ml	250
25–40 kg	30 ml	166
Over 40 kg	40 ml	125

*Do not administer to animals of less than 12.5 kg bodyweight.

Apply as a fan-spray to the surface of the fleece on the back and hindquarters of the sheep using the applicator gun fitted with a fan spray nozzle. Half the dose should be applied on the shoulders, back and flanks and half to the rump.

Maintain a distance between the nozzle and the fleece of approximately 20 cm. Each dose will require 2 or 3 sweeps to apply. **THE PRODUCT WILL PREVENT BLOWFLY STRIKE ONLY ON AREAS COVERED BY SPRAY.** One application will give 6–8 weeks protection. Re-treat as required.



Treatment of Blowfly Strike

Apply directly all to the affected parts at the rate of 2.5 ml per 100–150 cm² (roughly the size of a hand). The T-bar nozzle should be used. Most strikes will require 5–10 ml. Ensure that all affected parts are treated. A single application is sufficient to ensure that larvae are expelled and killed within a few hours.

Further Information


Blowflies are attracted by dirty, damp wool or open wounds. The most common site for flystrike is the rump, due to soiling caused by diarrhoea and urine staining. The efficacy of Vectocert Pour-On is reduced in the presence of soiled and dirty wool. Therefore sheep must be crutched (dagged) regularly and appropriate worm control measures employed.

Lice

5 ml per 20 kg bodyweight, up to a maximum of 20 ml.

BODYWEIGHT	DOSE	DOSES PER PACK
Up to 20 kg	5 ml	1000
21–40 kg	10 ml	500
41-60 kg	15 ml	333
Over 60kg	20 ml	250

Using the straight nozzle on the applicator gun, apply as a pin stream from the shoulders to the rump along the middle of the backline. Sheep may be treated off-shears or anytime during the year.




Ticks

Adult sheep and lambs over 10 kg: 10 ml per 20 kg bodyweight, up to a maximum of 40 ml.
Lambs under 10 kg: 5 ml followed 3 weeks later by a 10 ml application.

BODYWEIGHT	DOSE	DOSES PER PACK
Up to 10 kg	5 ml	1000
11-20 kg	10 ml	500
21-40 kg	20 ml	250
41-60 kg	30 ml	166
Over 60kg	40 ml	125

Using the straight nozzle on the applicator gun, apply as a pin stream from the crown of the head to the top of the rump.




Headflies

5 ml per animal irrespective of size.

Apply to the top of the head between the ears, using the T-bar nozzle. Take care not to apply the product in the sheep's eyes.

One application of Vectocert Pour-On before the start of the headfly season will give up to 4 weeks protection. Re-treat as required.



10. ADVICE ON CORRECT ADMINISTRATION

Advice on correct administration

Avoid treating very young lambs if the weather is unseasonably hot.

11. WITHDRAWAL PERIODS

Withdrawal periods

Meat and offal: 8 days.

Milk - Not authorised for use in animals producing milk for human consumption.

12. SPECIAL STORAGE PRECAUTIONS

Special storage precautions

Do not store above 25°C.

Protect from direct sunlight. Keep the container tightly closed.

Store away from food, drink and animal feed stuffs.

13. SPECIAL PRECAUTIONS FOR DISPOSAL

Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

The veterinary medicinal product should not enter water courses as cypermethrin may be dangerous for fish and other aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

Cypermethrin is moderately persistent and non-mobile in soil. Cypermethrin is toxic to dung insects. Long-term effects on dung insects caused by continuous or repeated use of the product cannot be excluded.

14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Classification of veterinary medicinal products

For external use only.

To be supplied only on veterinary prescription.

15. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 50146/4037

Pack sizes

1 Litre, 2.5 Litre, 5 Litre.

Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

17. CONTACT DETAILS

Contact details

Marketing authorisation holder:

Bimeda Animal Health Limited
2 / 3 / 4 Airton Close
Tallaght
Dublin 24
Ireland

Manufacturer responsible for batch release:

Pasteur Filiala Filipesti S.A.
Strada Principala Nr 944
Filipestii De Padure
107245
Romania

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder listed below.

DOWNLAND logo®

Downland Marketing Ltd.
Main Mill
Warwick Mill Business Centre
Warwick Bridge
Carlisle
CA4 8RR

18. OTHER INFORMATION

Other information

POM-VPS

19. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

20. EXPIRY DATE

Exp: {mm/yyyy}

Do not use this veterinary medicinal product after the expiry date which is stated on the label {mm/yyyy}. The expiry date refers to the last day of that month.

The product must be used within 3 months of opening the container.
Once opened use by

21. BATCH NUMBER

Lot {number}

Approved 28 May 2026
Gavin Hall