

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARTON BOX 100 ml and 250 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cadorex 300 mg/ml solution for injection for cattle, sheep and pigs
Florfenicol

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:
Florfenicol... 300 mg

3. PHARMACEUTICAL FORM

Solution for injection.

4. PACKAGE SIZE

100 ml
250 ml

5. TARGET SPECIES

Cattle, sheep and pigs.

6. INDICATION(S)

Read the package leaflet before use.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Cattle: Intramuscular or subcutaneous use
Sheep: Intramuscular use
Pigs: Intramuscular use
Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Withdrawal period:

Cattle:

Meat and offal:	by IM route:	30 days
	by SC route:	44 days

Milk: Not authorised for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Sheep:
Meat and offal: by IM route: 39 days

Milk: Not authorised for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Pigs:
Meat and offal: by IM route: 18 days

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP [month/year]
Once opened, use within 28 days.
Once broached, use by ...

11. SPECIAL STORAGE CONDITIONS

Store below 30°C.
Do not freeze.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of waste material in accordance with local requirements.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.
To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Livisto Int'l, S.L.
Av. Universitat Autònoma, 29
08290 Cerdanyola del Vallès
(Barcelona) Spain

16. MARKETING AUTHORISATION NUMBER(S)

Vm 43173/5001

17. MANUFACTURER'S BATCH NUMBER

Batch:

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

LABEL 100 ml and 250 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cadorex 300 mg/ml solution for injection for cattle, sheep and pigs
Florfenicol

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:
Florfenicol..... 300 mg

3. PHARMACEUTICAL FORM

Solution for injection.

4. PACKAGE SIZE

100 ml
250 ml

5. TARGET SPECIES

Cattle, sheep and pigs.

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Cattle: Intramuscular or subcutaneous use
Sheep: Intramuscular use
Pigs: Intramuscular use
Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Withdrawal period:

Cattle:
Meat and offal: by IM route: 30 days
 by SC route: 44 days

Milk: Not authorised for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Sheep:

Meat and offal: by IM route: 39 days

Milk: Not authorised for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Pigs:
Meat and offal: by IM route: 18 days

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP [month/year]
Once opened, use within 28 days.
Once broached, use by ...

11. SPECIAL STORAGE CONDITIONS

Store below 30°C.
Do not freeze.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of waste material in accordance with local requirements.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.
To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Livisto Int'l, S.L.
Av. Universitat Autònoma, 29
08290 Cerdanyola del Vallès
(Barcelona) Spain

16. MARKETING AUTHORISATION NUMBER(S)

Vm 43173/5001

17. MANUFACTURER'S BATCH NUMBER

Batch:

B. PACKAGE LEAFLET

PACKAGE LEAFLET FOR:
Cadorex 300 mg/ml solution for injection for cattle, sheep and pigs

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Livisto Int'l, S.L.
Av. Universitat Autònoma, 29
08290 Cerdanyola del Vallès
(Barcelona) Spain

Manufacturer responsible for batch release:

Industrial Veterinaria, S.A.
Esmeralda 19
08950 Esplugues de Llobregat (Barcelona) Spain

aniMedica GmbH
Im Südfeld 9
48308 Senden-Bösensell Germany

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cadorex 300 mg/ml solution for injection for cattle, sheep and pigs
Florfenicol

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each ml contains:

Active substance:

Florfenicol..... 300 mg

Excipients:

N-methyl pyrrolidone..... 250 mg

Clear, light yellow to straw-coloured, somewhat viscous solution, free from foreign matter.

4. INDICATION(S)

Cattle:

Diseases caused by florfenicol susceptible to bacteria: Treatment of respiratory tract infections in cattle due to *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*.

Sheep:

Treatment of ovine respiratory tract infections due to *Mannheimia haemolytica* and *Pasteurella multocida*.

Pigs:

Treatment of acute outbreaks of swine respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

5. CONTRAINDICATIONS

Do not use in adult bulls and rams intended for breeding purposes.

Do not administer to boars intended for breeding.

Do not use in case of known hypersensitivity to the active substance or to any of the excipients.

6. ADVERSE REACTIONS

Cattle:

A decrease in food consumption and transient softening of the faeces may occur very rare during the treatment period. The treated animals recover quickly and completely upon termination of treatment.

Administration of the product by the intramuscular and subcutaneous routes may cause inflammatory lesions at injection site very rare which persist for 14 days.

Anaphylactic shocks have been reported in bovines in very rare cases.

Sheep:

A decrease in food consumption may occur very rare during the treatment period. The treated animals recover quickly and completely upon termination of the treatment.

Administration of the product by the intramuscular route may cause inflammatory lesions at the injection site very rare which may persist up to 28 days. Typically, these are mild and transient.

Pigs:

Commonly observed adverse effects are transient diarrhoea and/or peri-anal and rectal erythema/oedema which may affect 50% of the animals. These effects can be observed for one week.

Under field conditions approximately 30% of treated pigs presented with pyrexia (40°C) associated with either moderate depression or moderate dyspnoea a week or more after administration of the second dose.

Transient swelling lasting up to 5 days may be observed at the site of injection very rare. Inflammatory lesions at the injection site may be seen up to 28 days.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

Alternatively you can report via your national reporting system {national system details}.

7. TARGET SPECIES

Cattle, sheep and pigs.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For intramuscular and subcutaneous use in cattle.

For intramuscular use in sheep and pigs.

For treatment

Cattle:

Intramuscular route: 20 mg of florfenicol/kg bodyweight (equivalent to 1 ml of the product/15 kg bodyweight) to be administered twice 48 hours apart using a 16 gauge needle.

Subcutaneous route: 40 mg of florfenicol/kg bodyweight (equivalent to 2 ml of the product/15 kg bodyweight) to be administered once using a 16 gauge needle. The dose volume given at any one injection site should not exceed 10 ml.

The injection should only be given in the neck.

Sheep:

20 mg of florfenicol/kg bodyweight (equivalent to 1 ml of the product/15 kg bodyweight) by intramuscular injection daily for three consecutive days. The volume administered per injection site should not exceed 4 ml.

Pigs:

15 mg of florfenicol/kg bodyweight (equivalent to 1 ml of the product/ 20 kg bodyweight) by intramuscular injection into the neck muscle twice at 48 hours intervals using a 16-gauge needle.

The volume administered per injection site should not exceed 3 ml.

For intramuscular, it is recommended to treat animals in the early stages of disease and to evaluate the response to treatment within 48 hours after the second injection. If clinical signs of respiratory disease persist 48 hours after the last injection, treatment should be changed using another formulation or another antibiotic and continued until clinical signs have resolved.

9. ADVICE ON CORRECT ADMINISTRATION

Wipe the stopper before removing each dose. Use a dry sterile needle and syringe.

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing.

As the vial should not be broached more than 25 times, the user should select the most appropriate vial size according to the target species to be treated. When treating groups of animals in one run, use a draw-off needle that has been placed in the vial stopper to avoid excess broaching of the stopper. The draw-off needle should be removed after treatment.

10. WITHDRAWAL PERIOD

Cattle:

Meat and offal:	by IM route:	30 days
	by SC route:	44 days

Milk: Not authorised for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Sheep:

Meat and offal:	by IM route:	39 days
-----------------	--------------	---------

Milk: Not authorised for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Pigs:

Meat and offal:	by IM route:	18 days
-----------------	--------------	---------

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store below 30°C.

Do not freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Shelf-life after first opening the container: 28 days

12. SPECIAL WARNING(S)

Special warnings for each target species:

None.

Special precautions for use in animals:

This medicinal product does not contain any antimicrobial preservative.

The safety of the product has not been established in sheep under 7 weeks of age.

Do not use in piglets of less than 2 kg.

Use of the product should be based on identification and susceptibility testing of the target pathogens. Use of the product should be in accordance with official, national and regional antimicrobial policies.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the florfenicol and may decrease the effectiveness of treatment with other amphenicols due to the potential for cross-resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This product may cause hypersensitivity (allergy).

People with known hypersensitivity to florfenicol, propylene glycol or polyethylene glycols should avoid contact with the veterinary medicinal product. Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. Women of childbearing age, pregnant women or women suspected of being pregnant should use the veterinary medicinal product with serious caution to avoid accidental self-injection.

Take care to avoid accidental self-injection. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Avoid skin or eye contact with the product. In case of contact with the skin or eyes, rinse the affected area immediately with plenty of clean water.

If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the physician the package leaflet or the label.

Wash hands after use.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established in cattle, sheep and pigs during pregnancy, lactation or in animals intended for breeding.

Studies in laboratory animals have not revealed any evidence of embryo- or foetotoxic potential for florfenicol. Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

Fertility:

Do not use in adult bulls, rams and boars intended for breeding (see section 5).

Interaction with other medicinal products and other forms of interaction:

None known.

Overdose (symptoms, emergency procedures, antidotes):

Cattle:

No symptoms other than those described in section 6..

Sheep:

After administration of 3 times the recommended dose or more, a transient reduction in feed and water consumption has been observed. Additional secondary effects that were noted included an increased incidence of lethargy, emaciation and loose faeces.

Head tilt was seen after administration of 5 times the recommended dose and was considered most likely a result of irritation at the injection site.

Swine:

After administration of 3 times the recommended dose or more, a reduction in feeding, hydration and weight gain has been observed.

After administration of 5 times the recommended dose or more, vomiting has also been noted.

Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

May 2023

15. OTHER INFORMATION

Package sizes:

Cardboard box containing 1 vial of 100 ml

Cardboard box containing 1 vial of 250 ml

Not all pack sizes may be marketed.

A handwritten signature in black ink, appearing to read 'Dennett', is positioned above the approval date.

Approved: 24 July 2023