## MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO PACKAGE LEAFLET, E.g. Concertina Labels. {NATURE/TYPE}

#### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Evans Vanodine International Plc Brierlay Road Walton Summit Preston PR5 8AH

## 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

MEGODINE 1:3 CONCENTRATE TEAT DIP AND TEAT SPRAY SOLUTION 2.15% w/v.

## 3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

lodine, Glycerol and Sorbitol.

## 4. PHARMACEUTICAL FORM

Concentrate for teat dip/spray solution.

Contains 2.15% w/v available iodine, 2% w/v glycerol and 18% w/v sorbitol

Effective against mastitis causing bacteria

lodine based teat dip/spray concentrate

Concentrate teat dip and spray solution 2.15% w/v

5000ppm available iodine minimum when diluted 1:3 as a teat dip or spray

40 ppm available iodine minimum when diluted 1:500 as udderwash or cluster dip

#### Product Information

An iodophor-based teat dip and teat spray which has been specially formulated to help reduce the bacterial contamination that causes mastitis in dairy herds. Reduced the risk of infection by leaving an antiseptic film of iodine adhering to the teats which have treated by dipping or spraying. Contains 2% w/v glycerol and 18% w/v sorbitol (0.5% w/v/ glycerol and 4.5% w/v sorbitol when diluted) to help keep teats soft and supple, and to aid in the prevention and healing of sores and cracks.

**TEAT DIP or SPRAY** (5000ppm available iodine)

Dilute 1 part Megodine 1:3 concentrate Teat Dip and Teat Spray Solution 2.15% w/v with 3 parts clean water and mix well.

**UDDER WASH AND CLUSTER DIP** (40ppm minimum available iodine) Dilute 1 part Megodine 1:3 concentrate Teat Dip and Teat Spray Solution 2.15% w/v with 500 parts clean water (20ml to 10 litres or 1 fl.oz.in 3 gall).

## 5. PACKAGE SIZE

5,25,60,200

Additional wording required on 200 litre label:

200 litre label: This container should only be used once and should not be returned for refilling.

#### 6. INDICATION(S)

[Optional. In case of space restriction and if the indication is clear from the name of the product, the indication should not be repeated]

## 7. CONTRAINDICATIONS

## 8. ADVERSE REACTIONS

#### 9. TARGET SPECIES

## 10. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Directions for use

TEAT DIPPING: Fill teat dip cup about two-thirds full with the prepared solution. Dip teats of every cow immediately after each has been milked, making sure that the full length of the teat is immersed. Top up with fresh solution as required. Empty and wash out cups after milking.

**TEAT SPRAYING** Immediately after milking, spray entire surface of each teat of every cow with prepared solution.

Dilute 1: 500. To help prevent cross infection, individual cloths or disposable paper towels are strongly recommended.

Dry teats before milking. Immerse clusters and agitate for at least half a minute. Rinse in clean water before use. Wash and dry udders and teats before milking.

## **11. ADVICE ON CORRECT ADMINISTRATION**

#### CONCENTRATE

The following safety phrases relate to the concentrated product only, and do not apply once the product is diluted to the working solution:

Risk of serious damage to eyes. Wear eye/face protection when preparing the dip or spray.

## **DILUTED WORKING SOLUTION**

When using as a spray, avoid working in spray mist.
Avoid contact with eyes.
In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.
If swallowed, seek medical advice immediately and show this container or label.
Hands and exposed skin should be washed after using this product.
Do not eat, drink or smoke while using the product.

Keep away from food, drink and animal feedstuffs.

#### 12. WITHDRAWAL PERIOD

Milk: Zero hours. Meat: Zero days.

#### 13. SPECIAL STORAGE PRECAUTIONS

Do not store above 25°C. Store in tightly closed original container. Protect from frost. Diluted product should not be stored above 25°C. Protect from light. Once diluted use the product within 3 months. Store diluted product only in glass or plastic containers.

#### 14. SPECIAL WARNING(S)<User Warnings>

For animal treatment only. For external use only. Keep out of the reach of children. If signs of disease persist or appear, consult your veterinary surgeon.

#### **15. EXPIRY DATE**

# 16. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Harmful to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the product or empty container. Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority. To dispose of unused product to land you must have an authorisation under the groundwater regulations 1998.

#### 17. DATE ON WHICH THE LABEL WAS LAST APPROVED

September 2009

# 18. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.

[Distribution category]

	AVM-GSL
--	---------

## 19. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

## 20. MARKETING AUTHORISATION NUMBER(S)

#### Vm <number>

Vm03940/4049

#### 21. MANUFACTURER'S BATCH NUMBER

## 22. OTHER INFORMATION>

## Approved: 09/10/2017

